SFB/CPP-14-16, TTP14-007

Prepared for submission to JHEP

Quark Mass and Field Anomalous Dimensions to $\mathcal{O}(\alpha_s^5)$

P. A. Baikov,[a] **K. G. Chetyrkin,**[b] **J. H. Kühn,**[b]

[a] *Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, 1(2), Leninskie gory, Moscow 119991, Russian Federation*

[b] *Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT), Wolfgang-Gaede-Straße 1, 726128 Karlsruhe, Germany*

E-mail: baikov@theory.sinp.msu.ru, Konstantin.Chetyrkin@kit.edu, johann.kuehn@kit.edu

Abstract:

We present the results of the first complete analytic calculation of the quark mass and field anomalous dimensions to $\mathcal{O}(\alpha_s^5)$ in QCD.

Keywords: Quantum chromodynamics, Perturbative calculations

Contents

1 Introduction

The quark masses depend on a renormalization scale. The dependence is usually referred to as "running" and is governed by the quark mass anomalous dimension, γ_m, defined as:

$$\mu^2 \frac{d}{d\mu^2} m|_{g^0,m^0} = m\gamma_m(a_s) \equiv -m \sum_{i\geq 0} \gamma_i \, a_s^{i+1}, \tag{1.1}$$

where $a_s = \alpha_s/\pi = g^2/(4\pi^2)$, g is the renormalized strong coupling constant and μ is the normalization scale in the customarily used $\overline{\text{MS}}$ renormalization scheme. Up to and including four loop level the anomalous dimension is known since long [1–5]. In this paper we will describe the results of calculation of γ_m and a related quantity — the quark field anomalous dimension — in the five-loop order.

The evaluation of the quark mass anomalous dimension with five-loop accuracy has important implications. The Higgs boson decay rate into charm and bottom quarks is proportional to the square of the respective quark mass at the scale of m_H and the uncertainty from the presently unknown 5-loop terms in the running of the quark mass is of order 10^{-3}. This is comparable to the precision advocated for experiments e.g. at TLEP [6]. Similarly, the issue of Yukawa unification is affected by precise predictions for the anomalous quark mass dimension.

The paper is organized as follows. The next section deals with the overall set-up of the calculations. Then we present our results (Section 3), and a brief discussion (Section 4) as well as a couple of selected applications (Section 5). Our short conclusions are given Section 6.

2 Technical preliminaries

To calculate γ_m one needs to find the so-called quark mass renormalization constant, Z_m, which is defined as the ratio of the bare and renormalized quark masses, viz.

$$Z_m = \frac{m^0}{m} = 1 + \sum_{i,j}^{0<j\leq i} (Z_m)_{ij} \frac{a_s^i}{\epsilon^j}. \tag{2.1}$$

Within the $\overline{\text{MS}}$ scheme [7, 8] the coefficients $(Z_m)_{ij}$ are just numbers [9]; $\epsilon \equiv 2 - D/2$ and D stands for the space-time dimension. Combining eqs. (1.1,2.1) and using the RG-invariance of of m^0, one arrives at the following formula for γ_m:

$$\gamma_m = \sum_{i \geq 0} (Z_m)_{i1} \, i \, a_s^i. \tag{2.2}$$

To find Z_m one should compute the vector and scalar parts of the quark self-energy $\Sigma_V(p^2)$ and $\Sigma_S(p^2)$. In our convention, the bare quark propagator is proportional to $\left[\not{p}\left(1 + \Sigma_V^0(p^2)\right) - m_q^0\left(1 - \Sigma_S^0(p^2)\right)\right]^{-1}$. Requiring the finiteness of the renormalized quark propagator and keeping only massless and terms linear in m_q, one arrives at the following recursive equations to find Z_m

$$Z_m Z_2 = 1 + K_\epsilon\left\{Z_m Z_2 \Sigma_S^0(p^2)\right\}, \quad Z_2 = 1 - K_\epsilon\left\{Z_2 \Sigma_V^0(p^2)\right\}, \tag{2.3}$$

where $K_\epsilon\{f(\epsilon)\}$ stands for the singular part of the Laurent expansion of $f(\epsilon)$ in ϵ near $\epsilon = 0$ and Z_2 is the quark wave function renormalization constant. Eqs. (2.3) express Z_m through massless propagator-type (that is dependent on one external momentum only) Feynman integrals (FI), denoted as *p-integrals* below.

Eqs. (2.3) require the calculation of a large number[1] the *five*-loop p-integrals to find Z_m and Z_2 to $\mathcal{O}(\alpha_s^5)$.

At present there exists no direct way to analytically evaluate five-loop p-integrals. However, according to (2.1) for a given five-loop p-integral we need to know only its *pole* part in ϵ in the limit of $\epsilon \to 0$. A proper use of this fact can significantly simplify our task. The corresponding method—so-called Infrared Rearrangement (IRR)—first suggested in [11] and elaborated further in [12–14] allows to effectively decrease number of loops to be computed by one[2]. In its initial version IRR was not really universal; it was not applicable in some (though rather rare) cases of complicated FI's. The problem was solved by elaborating a special technique of subtraction of IR divergences — the R^*-operation [15, 16]. This technique succeeds in expressing the UV counterterm of every L-loop Feynman integral in terms of divergent and finite parts of some (L-1)-loop massless propagators.

In our case $L = 5$ and, using IRR, one arrives at at around 10^5 four-loop p-integrals. These can, subsequently, be reduced to 28 four-loop masterp-integrals, which are known analytically, including their finite parts, from [17, 18] as well as numerically from [19].

[1] We have used QGRAF [10] to produce around 10^5 FI's contributing to the quark self-energy at $\mathcal{O}(\alpha_s^5)$.

[2] With the price that resulting one-loop-less p-integrals should be evaluated up to and *including* their constant part in the small ϵ-expansion.

We need, thus, to compute around 10^5 p-integrals. Their singular parts, in turn, can be algebraically reduced to only 28 master 4-loop p-integrals. The reduction is based on evaluating sufficiently many terms of the $1/D$ expansion [20] of the corresponding coefficient functions [21].

All our calculations have been performed on a SGI ALTIX 24-node IB-interconnected cluster of eight-cores Xeon computers using parallel MPI-based [22] as well as thread-based [23] versions of FORM [24].

3 Results

Our result for the anomalous dimension

$$\gamma_m = -\sum_{i=0}^{\infty} (\gamma_m)_i \, a_s^{i+1}$$

reads:

$$(\gamma_m)0 = 1, \qquad (\gamma_m)1 = \frac{1}{16}\left\{ \frac{202}{3} + n_f \left[-\frac{20}{9} \right] \right\}, \tag{3.1}$$

$$(\gamma_m)2 = \frac{1}{64}\left\{ 1249 + n_f \left[-\frac{2216}{27} - \frac{160}{3}\zeta_3 \right] + n_f^2 \left[-\frac{140}{81} \right] \right\}, \tag{3.2}$$

$$\begin{aligned}(\gamma_m)3 = \frac{1}{256}\Bigg\{ & \frac{4603055}{162} + \frac{135680}{27}\zeta_3 - 8800\,\zeta_5 \\ + \; & n_f \left[-\frac{91723}{27} - \frac{34192}{9}\zeta_3 + 880\,\zeta_4 + \frac{18400}{9}\zeta_5 \right] \\ + \; & n_f^2 \left[\frac{5242}{243} + \frac{800}{9}\zeta_3 - \frac{160}{3}\zeta_4 \right] + n_f^3 \left[-\frac{332}{243} + \frac{64}{27}\zeta_3 \right] \Bigg\}.\end{aligned} \tag{3.3}$$

$$\begin{aligned}(\gamma_m)_4 = \frac{1}{4^5}\Bigg\{ & \frac{99512327}{162} + \frac{46402466}{243}\zeta_3 + 96800\,\zeta_3^2 - \frac{698126}{9}\zeta_4 \\ & - \frac{231757160}{243}\zeta_5 + 242000\,\zeta_6 + 412720\,\zeta_7 \\ + \; n_f & \left[-\frac{150736283}{1458} - \frac{12538016}{81}\zeta_3 - \frac{75680}{9}\zeta_3^2 + \frac{2038742}{27}\zeta_4 \right. \\ & \left. + \frac{49876180}{243}\zeta_5 - \frac{638000}{9}\zeta_6 - \frac{1820000}{27}\zeta_7 \right] \\ + \; n_f^2 & \left[\frac{1320742}{729} + \frac{2010824}{243}\zeta_3 + \frac{46400}{27}\zeta_3^2 - \frac{166300}{27}\zeta_4 - \frac{264040}{81}\zeta_5 + \frac{92000}{27}\zeta_6 \right] \\ + \; & \boxed{n_f^3 \left[\tfrac{91865}{1458} + \tfrac{12848}{81}\zeta_3 + \tfrac{448}{9}\zeta_4 - \tfrac{5120}{27}\zeta_5 \right] + n_f^4 \left[-\tfrac{260}{243} - \tfrac{320}{243}\zeta_3 + \tfrac{64}{27}\zeta_4 \right]} \Bigg\}.\end{aligned} \tag{3.4}$$

Here ζ is the Riemann zeta-function ($\zeta_3 = 1.202056903\ldots$, $\zeta_4 = \pi^4/90$, $\zeta_5 = 1.036927755\ldots$, $\zeta_6 = 1.017343062\ldots$ and $\zeta_7 = 1.008349277\ldots$). Note that in four-loop order we exactly[3]

[3] This agreement can be also considered as an important check of all our setup which is completely different from the ones utilized at the four-loop calculations.

reproduce well-known results obtained in [4, 5]. The boxed terms in (3.4) are in full agreement with the results derived previously on the basis of the $1/n_f$ method in [25–27].

For completeness we present below the result for the quark field anomalous dimension $\gamma_2 = -\sum_{i=0}^{\infty} (\gamma_2)_i \, a_s^{i+1}$:

$$
\begin{aligned}
(\gamma_2)_4 = \frac{1}{4^5}\Bigg\{ & \frac{2798900231}{7776} + \frac{17969627}{864}\zeta_3 + \frac{13214911}{648}\zeta_3^2 + \frac{16730765}{864}\zeta_4 - \frac{832567417}{3888}\zeta_5 \\
& + \frac{40109575}{1296}\zeta_6 + \frac{124597529}{1728}\zeta_7 \\
+ n_f \Bigg[& -\frac{861347053}{11664} - \frac{274621439}{11664}\zeta_3 + \frac{1960337}{972}\zeta_3^2 + \frac{465395}{1296}\zeta_4 \\
& + \frac{22169149}{5832}\zeta_5 + \frac{1278475}{1944}\zeta_6 + \frac{3443909}{216}\zeta_7 \Bigg] \\
+ n_f^2 \Bigg[& \frac{37300355}{11664} + \frac{1349831}{486}\zeta_3 - \frac{128}{9}\zeta_3^2 - \frac{27415}{54}\zeta_4 - \frac{12079}{27}\zeta_5 - \frac{800}{9}\zeta_6 - \frac{1323}{2}\zeta_7 \Bigg] \\
+ n_f^3 \Bigg[& -\frac{114049}{8748} - \frac{1396}{81}\zeta_3 + \frac{208}{9}\zeta_4 \Bigg] + n_f^4 \Bigg[\frac{332}{729} - \frac{64}{81}\zeta_3 \Bigg] \Bigg\}.
\end{aligned} \tag{3.5}
$$

The above result is presented for the Feynman gauge; the coefficients $(\gamma_2)_i$ with $i \leq 3$ can be found in [28] (for the case of a general covariant gauge and SU(N) gauge group).

4 Discussion

In numerical form γ_m reads

$$
\begin{aligned}
\gamma_m = & - a_s - a_s^2 \, (4.20833 - 0.138889 n_f) \\
& - a_s^3 \left(19.5156 - 2.28412 n_f - 0.0270062 n_f^2\right) \\
& - a_s^4 \left(98.9434 - 19.1075 n_f + 0.276163 n_f^2 + 0.00579322 n_f^3\right) \\
& - a_s^5 \left(559.7069 - 143.6864 \, n_f + 7.4824 \, n_f^2 + 0.1083 \, n_f^3 - 0.000085359 \, n_f^4\right)
\end{aligned} \tag{4.1}
$$

and

$$
\begin{aligned}
\gamma_m & \underset{n_f=3}{=\!=} - a_s - 3.79167 \, a_s^2 - 12.4202 \, a_s^3 - 44.2629 \, a_s^4 - 198.907 \, a_s^5, \\
g_m & \underset{n_f=4}{=\!=} - a_s - 3.65278 \, a_s^2 - 9.94704 \, a_s^3 - 27.3029 \, a_s^4 - 111.59 \, a_s^5, \\
g_m & \underset{n_f=5}{=\!=} - a_s - 3.51389 \, a_s^2 - 7.41986 \, a_s^3 - 11.0343 \, a_s^4 - 41.8205 \, a_s^5, \\
\gamma_m & \underset{n_f=6}{=\!=} - a_s - 3.37500 \, a_s^2 - 4.83867 \, a_s^3 + 4.50817 \, a_s^4 + 9.76016 \, a_s^5.
\end{aligned} \tag{4.2}
$$

Note that significant cancellations between n_f^0 and n_f^1 terms for the values of n_f around 3 or so persist also at five-loop order. As a result we observe a moderate growth of the series in a_s appearing in the quark mass anomalous dimension at various values of active quark flavours (recall that even for scales as small as 2 GeV $a_s \equiv \frac{\alpha_s}{\pi} \approx 0.1$).

n_f	3	4	5	6
$(\gamma_m)_4^{\rm exact}$	198.899	111.579	41.807	-9.777
$(\gamma_m)_4^{\rm APAP}$ [29]	162.0	67.1	-13.7	-80.0
$(\gamma_m)_4^{\rm APAP}$ [30]	163.0	75.2	12.6	12.2
$(\gamma_m)_4^{\rm APAP}$ [31]	164.0	71.6	-4.8	-64.6

Table 1: The exact results for $(\gamma_m)_4$ together with the predictions made with the help of the original APAP method and its two somewhat modified versions.

Similar behavior shows up for γ_2:

$$\begin{aligned}\gamma_2 = &- 0.33333 a_s - a_s^2\,(-1.9583 + 0.08333\,n_f)\\ &- a_s^3\,(-10.3370 + 1.0877\,n_f - 0.01157\,n_f^2)\\ &- a_s^4\,(-53.0220 + 10.1090\,n_f - 0.27703\,n_f^2 - 0.0023\,n_f^3)\\ &- a_s^4\,(-310.0700 + 76.3260\,n_f - 4.6339\,n_f^2 + 0.0085\,n_f^3 + 0.00048\,n_f^4) \qquad (4.3)\end{aligned}$$

and

$$\begin{aligned}\gamma_2 &\underset{n_f=3}{==} - 0.33333\,a_s - 1.7083\,a_s^2 - 7.1779\,a_s^3 - 25.2480\,a_s^4 - 122.5300\,a_s^5,\\ g_2 &\underset{n_f=4}{==} - 0.33333\,a_s - 1.6250\,a_s^2 - 6.1712\,a_s^3 - 17.1610\,a_s^4 - 78.2430\,a_s^5,\\ g_2 &\underset{n_f=5}{==} - 0.33333\,a_s - 1.5417\,a_s^2 - 5.1877\,a_s^3 - 9.6824\,a_s^4 - 42.9240\,a_s^5,\\ \gamma_2 &\underset{n_f=6}{==} - 0.33333\,a_s - 1.4583\,a_s^2 - 4.2274\,a_s^3 - 2.8251\,a_s^4 - 16.4710\,a_s^5. \qquad (4.4)\end{aligned}$$

It is instructive to compare our numerical result for $(\gamma_m)_4$

$$(\gamma_m)_4 = 559.71 - 143.6\,n_f + 7.4824\,n_f^2 + 0.1083\,n_f^3 - 0.00008535\,n_f^4 \qquad (4.5)$$

with a 15 years old prediction based on the "Asymptotic Páde Approximants" (APAP) method [29] (the boxed term below was used as the input)

$$(\gamma_m)_4^{\rm APAP} = 530 - 143\,n_f + 6.67\,n_f^2 + 0.037\,n_f^3 - \boxed{0.00008535\,n_f^4} \qquad (4.6)$$

Unfortunately, this impressively good agreement does *not* survive for fixed values of n_f due to severe cancellations between different powers of n_f as one can see from the Table 1.

The solution of eq. (1.1) reads:

$$\frac{m(\mu)}{m(\mu_0)} = \frac{c(a_s(\mu))}{c(a_s(\mu_0))}, \quad c(x) = \exp\left\{\int dx' \frac{\gamma_m(x')}{\beta(x')}\right\}, \qquad (4.7)$$

$$\begin{aligned}c(x) = \quad &(x)^{\bar\gamma_0}\left\{1 + d_1 x + (d_1^2/2 + d_2)\,x^2 + (d_1^3/6 + d_1 d_2 + d_3)\,x^3\right.\\ + \;&\left.(d_1^4/24 + d_1^2 d_2/2 + d_2^2/2 + d_1 d_3 + d_4)\,x^4 + \mathcal{O}(x^5)\right\}, \qquad (4.8)\end{aligned}$$

$$d_1 = -\bar\beta_1\,\bar\gamma_0 + \bar\gamma_1, \tag{4.9}$$
$$d_2 = \bar\beta_1^2\,\bar\gamma_0/2 - \bar\beta_2\,\bar\gamma_0/2 - \bar\beta_1\,\bar\gamma_1/2 + \bar\gamma_2/2, \tag{4.10}$$
$$d_3 = -\bar\beta_1^3\,\bar\gamma_0/3 + 2\,\bar\beta_1\,\bar\beta_2\,\bar\gamma_0/3 - \bar\beta_3\,\bar\gamma_0/3 + \bar\beta_1^2\,\bar\gamma_1/3 - \bar\beta_2\,\bar\gamma_1/3 - \bar\beta_1\,\bar\gamma_2/3 + \bar\gamma_3/3, \tag{4.11}$$
$$\begin{aligned} d_4 = {} & \bar\beta_1^4\,\bar\gamma_0/4 - 3\,\bar\beta_1^2\,\bar\beta_2\,\bar\gamma_0/4 + \bar\beta_2^2\,\bar\gamma_0/4 + \bar\beta_1\,\bar\beta_3\,\bar\gamma_0/2 - \bar\beta_4\,\bar\gamma_0/4 - \bar\beta_1^3\,\bar\gamma_1/4 \\ & + \bar\beta_1\,\bar\beta_2\,\bar\gamma_1/2 - \bar\beta_3\,\bar\gamma_1/4 + \bar\beta_1^2\,\bar\gamma_2/4 - \bar\beta_2\,\bar\gamma_2/4 - \bar\beta_1\,\bar\gamma_3/4 + \bar\gamma_4/4. \end{aligned} \tag{4.12}$$

Here $\bar\gamma_i = (\gamma_m)_i/\beta_0$, $\bar\beta_i = \beta_i/\beta_0$ and

$$\beta(a_s) = -\sum_{i\geq 0} \beta_i\, a_s^{i+2} = -\beta_0 \left\{ \sum_{i\geq 0} \bar\beta_i\, a_s^{i+2} \right\}$$

is the QCD β-function. Unfortunately, the coefficient d_4 in eq. (4.12) does depend on the yet unknown *five-loop* coefficient β_4 (up to four loops the β-function is known from [14, 32–39]).

Numerically, the c-function reads:

$$c(x) \underset{n_f=3}{=\!=} x^{4/9}\, c_s(x),\;\; c(x) \underset{n_f=4}{=\!=} x^{12/25}\, c_c(x),\;\; c(x) \underset{n_f=5}{=\!=} x^{12/23}\, c_b(x),\;\; c(x) \underset{n_f=6}{=\!=} x^{4/7}\, c_t(x),$$

with

$$\begin{aligned} c_s(x) &= 1 + 0.8950\,x + 1.3714\,x^2 + 1.9517\,x^3 + (15.6982 - 0.11111\,\bar\beta_4)\,x^4, \\ c_c(x) &= 1 + 1.0141\,x + 1.3892\,x^2 + 1.0905\,x^3 + (9.1104 - 0.12000\,\bar\beta_4)\,x^4, \\ c_b(x) &= 1 + 1.1755\,x + 1.5007\,x^2 + 0.17248\,x^3 + (2.69277 - 0.13046\,\bar\beta_4)\,x^4, \\ c_t(x) &= 1 + 1.3980\,x + 1.7935\,x^2 - 0.68343\,x^3 + (-3.5130 - 0.14286\,\bar\beta_4)\,x^4. \end{aligned} \tag{4.13}$$

5 Applications

5.1 RGI mass

Eq. (4.7) naturally leads to an important concept: the RGI mass

$$m^{\rm RGI} \equiv m(\mu_0)/c(a_s(\mu_0)), \tag{5.1}$$

which is often used in the context of lattice calculations. The mass is μ and *scheme* independent; in *any* (mass-independent) scheme

$$\lim_{\mu\to\infty} a_s(\mu)^{-\bar\gamma_0}\; m(\mu) = m^{\rm RGI}.$$

The function $c_s(x)$ is used, e.g, by the **ALPHA** lattice collaboration to find the $\overline{\rm MS}$ mass of the strange quark at a lower scale, say, $m_s(2\text{ GeV})$ from the $m_s^{\rm RGI}$ mass determined from lattice simulations (see, e.g. [40]). For example, setting $a_s(\mu = 2\,\text{GeV}) = \frac{\alpha_s(\mu)}{\pi} = 0.1$, we arrive at ($h$ counts loops):

$$\begin{aligned} m_s(2\,\text{GeV}) = m_s^{\rm RGI}\,(a_s(2\,\text{GeV}))^{\frac{4}{9}}\, \Big(& 1 + 0.0895\,h^2 + 0.0137\,h^3 + 0.00195\,h^4 \\ & + (0.00157 - 0.000011\,\bar\beta_4)\,h^5 \Big) \end{aligned} \tag{5.2}$$

In order to have an idea of effects due the five-loop term in (5.2) one should make a guess about $\bar{\beta}_4$. By inspecting lower orders in

$$\beta(n_f = 3) = -\left(\frac{4}{9}\right)\left(a_s + 1.777\,a_s^2 + 4.4711\,a_s^3 + 20.990\,a_s^4 + \bar{\beta}_4\,a_s^5\right)$$

one can assume a natural estimate of $\overline{\beta}_4$ as laying in the interval $50-100$. With this choice we conclude that the (apparent) convergence of the above series is quite good even at a rather small energy scale of 2 GeV.

On the other hand, the authors of [30] estimate $\bar{\beta}_4$ in the $n_f = 3$ QCD as large as -850! With such a huge and negative value of $\bar{\beta}_4$ the five loop term in (5.2) would amount to 0.01092 and, thus, would significantly exceed the four-loop contribution (0.00195).

5.2 Higgs decay into quarks

The decay width of the Higgs boson into a pair of quarks can be written in the form

$$\Gamma(H \to \bar{f}f) = \frac{G_F\,M_H}{4\sqrt{2}\pi}\,m_f^2(\mu)\,R^S(s = M_H^2, \mu) \tag{5.3}$$

where μ is the normalization scale and R^S is the spectral density of the scalar correlator, known to α_s^4 from [41]

$$\begin{aligned} R^S(s = M_H^2, \mu = M_H) &= 1 + 5.667\,a_s + 29.147\,a_s^2 + 41.758\,a_s^3 - 825.7\,a_s^4 \\ &= 1 + 0.2041 + 0.0379 + 0.0020 - 0.00140 \end{aligned} \tag{5.4}$$

where we set $a_s = \alpha_s/\pi = 0.0360$ (for the Higgs mass value $M_H = 125$ GeV and $\alpha_s(M_Z) = 0.118$).

Expression (5.3) depends on two phenomenological parameters, namely, $\alpha_s(M_H)$ and the quark running mass m_q. In what follows we consider, for definiteness, the dominant decay mode $H \to \bar{b}b$. To avoid the appearance of large logarithms of the type $\ln \mu^2/M_H^2$ the parameter μ is customarily chosen to be around M_H. However, the starting value of m_b is usually determined at a much smaller scale (typically around 5-10 GeV [42]). The evolution of $m_b(\mu)$ from a lower scale to $\mu = M_h$ is described by a corresponding RG equation which is completely fixed by the quark mass anomalous dimension $\gamma(\alpha_s)$ and the QCD beta function $\beta(\alpha_s)$ (for QCD with $n_f = 5$). In order to match the $\mathcal{O}({\alpha_s}^4)$ accuracy of (5.4) one should know *both* RG functions β and γ_m in the five-loop approximation. Let us proceed, assuming conservatively that $0 \leq \overline{\beta}_4^{n_f=5} \leq 200$.

The value of $m_b(\mu = M_H)$ is to be obtained with RG running from $m_b(\mu = 10\,\mathrm{GeV})$ and, thus, depends on β and γ_m. Using the Mathematica package RunDec[4] [43] and eq. (4.13) we find for the shift from the five-loop term

$$\frac{\delta m_b^2(M_H)}{m_b^2(M_H)} = -1.3 \cdot 10^{-4}(\bar{\beta}_4 = 0)| - 4.3 \cdot 10^{-4}(\bar{\beta}_4 = 100)| - 7.3 \cdot 10^{-4}(\bar{\beta}_4 = 200)$$

[4] We have extended the package by including the five-loop effects to the running of α_s and quark masses.

If we set $\mu = M_H$, then the combined effect of $\mathcal{O}(\alpha_s^4)$ terms as coming from the five-loop running and four-loop contribution to R^S on

$$\Gamma(H \to \bar{b}b) = \frac{G_F\, M_H}{4\sqrt{2}\pi}\, m_f^2(M_H)\, R^S(s = M_H^2, M_H) \tag{5.5}$$

is around -2‰ (for $\bar{\beta}_4 = 100$). This should be contrasted to the parametric uncertainties coming from the input parameters $\alpha_s(M_Z) = 0.1185(6)$ [44] and $m_b(m_b) = 4.169(8)$ GeV [45] which correspond to ± 1‰ and ± 4‰ respectively.

We conclude, that the $\mathcal{O}(\alpha_s^4)$ terms in (5.4), (5.5)) are of no phenomenological relevancy at present. But, the situation could be different if the project of TLEP [6] is implemented. For instance, the uncertainty in $\alpha_s(M_Z)$ could be reduced to ±2‰ and Higgs boson branching ratios with precisions in the permille range are advertised.

6 Conclusions

We have analytically computed the anomalous dimensions of the quark mass γ_m and field γ_2 in the five loop approximation. The self-consistent description of the quark mass evolution at five loop requires the knowledge of the QCD β-function to the same number of loops. The corresponding, significantly more complicated calculation is under consideration.

K.G.C. thanks J. Gracey and members of the DESY-Zeuthen theory seminar for usefull discussions.

This work was supported by the Deutsche Forschungsgemeinschaft in the Sonderforschungsbereich/Transregio SFB/TR-9 "Computational Particle Physics". The work of P. Baikov was supported in part by the Russian Ministry of Education and Science under grant NSh-3042.2014.2.

References

[1] R. Tarrach, *The pole mass in perturbative qcd*, *Nucl. Phys.* **B183** (1981) 384.

[2] O. V. Tarasov, *Anomalous dimensions of quark masses in three loop approximation*, . JINR-P2-82-900.

[3] S. A. Larin, *The renormalization of the axial anomaly in dimensional regularization*, *Phys. Lett.* **B303** (1993) 113–118, [hep-ph/9302240].

[4] K. G. Chetyrkin, *Quark mass anomalous dimension to O(alpha(s)**4)*, *Phys. Lett.* **B404** (1997) 161–165, [hep-ph/9703278].

[5] J. A. M. Vermaseren, S. A. Larin, and T. van Ritbergen, *The 4-loop quark mass anomalous dimension and the invariant quark mass*, *Phys. Lett.* **B405** (1997) 327–333, [hep-ph/9703284].

[6] M. Bicer, H. Duran Yildiz, I. Yildiz, G. Coignet, M. Delmastro, et al., *First Look at the Physics Case of TLEP*, arXiv:1308.6176.

[7] G. 't Hooft and M. J. G. Veltman, *Regularization and Renormalization of Gauge Fields*, *Nucl. Phys.* **B44** (1972) 189–213.

[8] W. A. Bardeen, A. J. Buras, D. W. Duke, and T. Muta, *Deep inelastic scattering beyond the leading order in asymptotically free gauge theories*, *Phys. Rev.* **D18** (1978) 3998.

[9] J. C. Collins, *Normal Products in Dimensional Regularization*, *Nucl. Phys.* **B92** (1975) 477.

[10] P. Nogueira, *Automatic feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279–289.

[11] A. A. Vladimirov, *Method For Computing Renormalization Group Functions In Dimensional Renormalization Scheme*, *Theor. Math. Phys.* **43** (1980) 417.

[12] D. I. Kazakov, O. V. Tarasov, and A. A. Vladimirov, *Calculation of critical exponents by quantum field theory methods*, *Sov. Phys. JETP* **50** (1979) 521.

[13] K. G. Chetyrkin, A. L. Kataev, and F. V. Tkachov, *New Approach to Evaluation of Multiloop Feynman Integrals: The Gegenbauer Polynomial x Space Technique*, *Nucl. Phys.* **B174** (1980) 345–377.

[14] O. V. Tarasov, A. A. Vladimirov, and A. Y. Zharkov, *The gell-mann-low function of qcd in the three loop approximation*, *Phys. Lett.* **B93** (1980) 429–432.

[15] K. G. Chetyrkin and V. A. Smirnov, *R* OPERATION CORRECTED*, *Phys. Lett.* **B144** (1984) 419–424.

[16] K. G. Chetyrkin, *Corrections of order alpha(s)**3 to R(had) in pQCD with light gluinos*, *Phys. Lett.* **B391** (1997) 402–412, [hep-ph/9608480].

[17] P. A. Baikov and K. G. Chetyrkin, *Four-Loop Massless Propagators: an Algebraic Evaluation of All Master Integrals*, *Nucl. Phys.* **B837** (2010) 186–220, [arXiv:1004.1153].

[18] R. N. Lee, A. V. Smirnov, and V. A. Smirnov, *Master Integrals for Four-Loop Massless Propagators up to Transcendentality Weight Twelve*, *Nucl. Phys.* **B856** (2012) 95–110, [arXiv:1108.0732].

[19] A. V. Smirnov and M. Tentyukov, *Four Loop Massless Propagators: a Numerical Evaluation of All Master Integrals*, *Nucl. Phys.* **B837** (2010) 40–49, [arXiv:1004.1149].

[20] P. A. Baikov, *A practical criterion of irreducibility of multi-loop feynman integrals*, *Phys. Lett.* **B634** (2006) 325–329, [hep-ph/0507053].

[21] P. A. Baikov, *Explicit solutions of the 3–loop vacuum integral recurrence relations*, *Phys. Lett.* **B385** (1996) 404–410, [hep-ph/9603267].

[22] M. Tentyukov et al., *ParFORM: Parallel Version of the Symbolic Manipulation Program FORM*, cs/0407066.

[23] M. Tentyukov and J. A. M. Vermaseren, *The multithreaded version of FORM*, hep-ph/0702279.

[24] J. A. M. Vermaseren, *New features of form*, math-ph/0010025.

[25] A. Palanques-Mestre and P. Pascual, *The 1/n-f expansion of the gamma and beta functions in qed*, *Commun. Math. Phys.* **95** (1984) 277.

[26] M. Ciuchini, S. E. Derkachov, J. Gracey, and A. Manashov, *Computation of quark mass anomalous dimension at O(1 / N**2(f)) in quantum chromodynamics*, *Nucl.Phys.* **B579** (2000) 56–100, [hep-ph/9912221].

[27] M. Ciuchini, S. E. Derkachov, J. Gracey, and A. Manashov, *Quark mass anomalous dimension at O(1/N(f)**2) in QCD*, *Phys.Lett.* **B458** (1999) 117–126, [hep-ph/9903410].

[28] K. G. Chetyrkin and A. Retey, *Renormalization and running of quark mass and field in the*

regularization invariant and MS-bar schemes at three and four loops, *Nucl. Phys.* **B583** (2000) 3–34, [hep-ph/9910332].

[29] J. R. Ellis, I. Jack, D. Jones, M. Karliner, and M. Samuel, *Asymptotic Pade approximant predictions: Up to five loops in QCD and SQCD*, *Phys.Rev.* **D57** (1998) 2665–2675, [hep-ph/9710302].

[30] V. Elias, T. G. Steele, F. Chishtie, R. Migneron, and K. B. Sprague, *Pade improvement of QCD running coupling constants, running masses, Higgs decay rates, and scalar channel sum rules*, *Phys.Rev.* **D58** (1998) 116007, [hep-ph/9806324].

[31] A. Kataev and V. Kim, *Higgs boson decay into bottom quarks and uncertainties of perturbative QCD predictions*, arXiv:0804.3992.

[32] D. J. Gross and F. Wilczek, *Ultraviolet behavior of non-abelian gauge theories*, *Phys. Rev. Lett.* **30** (1973) 1343–1346.

[33] H. D. Politzer, *Reliable perturbative results for strong interactions?*, *Phys. Rev. Lett.* **30** (1973) 1346–1349.

[34] W. E. Caswell, *Asymptotic behavior of nonabelian gauge theories to two loop order*, *Phys. Rev. Lett.* **33** (1974) 244.

[35] D. R. T. Jones, *Two loop diagrams in yang-mills theory*, *Nucl. Phys.* **B75** (1974) 531.

[36] E. Egorian and O. V. Tarasov, *Two loop renormalization of the qcd in an arbitrary gauge*, *Theor. Math. Phys.* **41** (1979) 863–867.

[37] S. A. Larin and J. A. M. Vermaseren, *The three loop qcd beta function and anomalous dimensions*, *Phys. Lett.* **B303** (1993) 334–336, [hep-ph/9302208].

[38] T. van Ritbergen, J. A. M. Vermaseren, and S. A. Larin, *The four-loop beta function in quantum chromodynamics*, *Phys. Lett.* **B400** (1997) 379–384, [hep-ph/9701390].

[39] M. Czakon, *The four-loop QCD beta-function and anomalous dimensions*, *Nucl. Phys.* **B710** (2005) 485–498, [hep-ph/0411261].

[40] **ALPHA** Collaboration, M. Della Morte et al., *Non-perturbative quark mass renormalization in two-flavor qcd*, *Nucl. Phys.* **B729** (2005) 117–134, [hep-lat/0507035].

[41] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Scalar correlator at $\mathcal{O}(\alpha_s^4)$, Higgs decay into b- quarks and bounds on the light quark masses*, *Phys. Rev. Lett.* **96** (2006) 012003, [hep-ph/0511063].

[42] K. Chetyrkin, J. Kühn, A. Maier, P. Maierhofer, P. Marquard, et al., *Charm and Bottom Quark Masses: An Update*, *Phys.Rev.* **D80** (2009) 074010, [arXiv:0907.2110].

[43] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, *RunDec: A Mathematica package for running and decoupling of the strong coupling and quark masses*, *Comput. Phys. Commun.* **133** (2000) 43–65, [hep-ph/0004189].

[44] **Particle Data Group** Collaboration, J. Beringer et al., *Review of Particle Physics (RPP)*, *Phys.Rev.* **D86** (2012) 010001.

[45] A. A. Penin and N. Zerf, *Bottom Quark Mass from Υ Sum Rules to $\mathcal{O}(\alpha_s^3)$*, arXiv:1401.7035.