

Towards predictive flavour models in SUSY SU(5) GUTs with doublet-triplet splitting

Stefan Antusch^{*,† 1}, Ivo de Medeiros Varzielas^{* 2}, Vinzenz Maurer^{* 3},
Constantin Sluka^{* 4}, Martin Spinrath^{‡ 5},

** Department of Physics, University of Basel,
Klingelbergstr. 82, CH-4056 Basel, Switzerland*

*† Max-Planck-Institut für Physik (Werner-Heisenberg-Institut),
Föhringer Ring 6, D-80805 München, Germany*

*‡ Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology
Engesserstraße 7, D-76131 Karlsruhe, Germany*

Abstract

We discuss how the double missing partner mechanism solution to the doublet-triplet splitting problem in four-dimensional supersymmetric SU(5) Grand Unified Theories (GUTs) can be combined with predictive models for the quark-lepton Yukawa coupling ratios at the GUT scale. It is argued that towards this goal a second SU(5) breaking Higgs field in the adjoint representation is very useful and we discuss all possible renormalizable superpotentials with two adjoint Higgs fields and calculate the constraints on the GUT scale and effective triplet mass from a two-loop gauge coupling unification analysis. Two explicit flavour models with different predictions for the GUT scale Yukawa sector are presented, including shaping symmetries and a renormalizable messenger sector. Towards calculating the rates for proton decay induced by the exchange of colour triplets, the required Clebsch-Gordan coefficients for their couplings are calculated for the possible dimension five and six operators. They are provided in detailed tables in the appendix, together with additional helpful material for GUT flavour model building.

¹E-mail: stefan.antusch@unibas.ch

²E-mail: ivo.de@unibas.ch

³E-mail: vinzenz.maurer@unibas.ch

⁴E-mail: constantin.sluka@unibas.ch

⁵E-mail: martin.spinrath@kit.edu

1 Introduction

Grand Unified Theories (GUTs) offer one of the most attractive extensions of the Standard Model of particle physics (SM), unifying three of the four fundamental forces of nature. Towards a more fundamental theory of nature it would, however, be desirable to also explain the pattern of fermion masses and mixing with its plenty of parameters. Hence, successful models of flavour in GUTs need to address two main issues, firstly, they have to achieve a working mechanism for GUT symmetry breaking, including sufficient suppression of proton decay without unacceptably large fine-tuning, and secondly, they should provide correct predictions for the flavour observables, such as the Yukawa coupling ratios and mixing angles.

Existing models which manage to naturally suppress proton decay either predict the unrealistic quark-lepton Yukawa relation $Y_e = Y_d^T$ (e.g. [1]), which may only be viable in the presence of extensive uncontrolled higher order corrections (e.g. [2]) rendering the model non-predictive, or the experimentally disfavoured combination of the Georgi-Jarlskog relations $y_\mu = -3y_s$ and $y_e = \frac{1}{3}y_d$ [3] (as e.g. in [4]), or rely on linear combinations of GUT Yukawa operators (e.g. [5]), which again implies the loss of predictivity. Furthermore, there exists a large number of GUT models which focus on the flavour sector, but do not include the Higgs potential. So while there are several existing models focusing on one of these two concerns, we are not aware of any work to date capable of resolving the two challenges in full detail (and without invoking extra space-time dimensions) in a predictive setup, also given the present rather precise experimental data.

In this paper we employ the framework of supersymmetric (SUSY) GUTs, as in the minimal supersymmetric extension of the SM (MSSM) the gauge couplings unify to a surprising precision. The scale where they unify (about 10^{16} GeV) is also large enough to sufficiently suppress proton decay from dimension six GUT operators [6]. However, the introduction of SUSY in combination with GUTs leads to the so-called “Doublet-Triplet Splitting (DTS) Problem”. In a SUSY SU(5) GUT model the minimal embedding of the Higgs fields contains additional colour triplets which lead to baryon number violating operators with larger contributions than in non-SUSY GUTs. Therefore the colour triplets have to be very heavy to suppress proton decay sufficiently or their couplings to the MSSM fields should be very strongly suppressed. In SU(5) in four dimensions, a proposed solution to the DTS problem is the “missing partner mechanism” (MPM) [7, 8] or its improved version, the “Double Missing Partner Mechanism” (DMPM) [9] which we will review briefly in the next section. In the existing models referred to above, either the MPM or the DMPM is applied.

Since GUTs not only unify the forces of the SM into a single GUT force, but also the fermions into joint GUT representations, they are indeed a promising starting point for addressing the flavour puzzle. More specifically, GUTs are capable of predicting the ratios between the Yukawa couplings of quarks and leptons at the GUT scale. After their renormalization group evolution to low energies (including supersymmetric 1-loop threshold corrections), these predictions can be compared to the experimental data for quark and lepton masses. As well known, the prediction of minimal SU(5) for the charged

lepton and down-type quark Yukawa matrix, the relation $Y_e = Y_d^T$ already mentioned above, is strongly disfavoured by the experimental results on the fermion masses. But also the ubiquitous proposal for more realistic ratios, the above mentioned Georgi-Jarlskog relations, obtained from the introduction of a 45-dimensional Higgs representation of SU(5) and certain assumptions about the Yukawa textures [3], are disfavoured by the recently improved data [10].

To arrive at experimentally favoured predictions for GUT scale Yukawa coupling ratios, we employ an alternative approach [11] involving higher-dimensional operators which contain a GUT breaking Higgs field. Due to this, new Clebsch-Gordan (CG) factors appear as Yukawa coupling ratios, with interesting associated implications for the masses and mixing of the fermions, cf. [12, 13] and [14, 15, 16, 17]. The main goal of this paper is to show how GUT flavour models featuring these promising quark-lepton mass relations can be combined with a version of the DMPM for solving the DTS problem.

As explicit examples, we will present two models with these properties that are “UV complete” in terms of messenger fields and employ sets of discrete Abelian symmetries (referred to as shaping symmetries) such that only the desired effective GUT operators are generated when the heavy degrees of freedom are integrated out. The two models predict the GUT scale quark lepton Yukawa ratios but are not yet predictive for the fermion mixing parameters (although the experimentally observed values can be fitted by both of the models), so we only view them as existence proofs which show that DTS and predictive Yukawa coupling ratios can indeed be combined. The strategies discussed here, however, provide the tools for the construction of more ambitious GUT models of flavour, which should finally also predict the quark and lepton mixings and CP phases (and include the observed neutrino masses).

The paper is organised as follows: We will start with a brief review of the MPM and DMPM and the recently proposed alternative Yukawa coupling ratios. We will also discuss the implications of replacing the 75-dimensional representation used in the MPM and DMPM with an adjoint 24-dimensional representation of SU(5). This choice is particularly well suited towards combining the DMPM with the novel CG factors. In section 3 we discuss the impact of the additional fields on gauge coupling unification and on their implications for the colour triplet masses. We will especially focus here on the case of superpotentials with two adjoint Higgs representations. We then address the Yukawa sector in section 4, where we describe the above-mentioned two predictive example models. Before we summarise and conclude in section 6, in section 5 we briefly comment on proton decay, showing it is under control in the proposed class of models. We present additional helpful material for model building in the appendices.

2 Strategy

In this section we present the general strategy that will be implemented in two example models. We begin our presentation with the review of the MPM and DMPM and the origin of the CG coefficients (coming from higher dimensional operators) used in our models.

Afterwards we discuss a modification of the DMPM with a GUT Higgs field in the adjoint representation and follow that with the actual DMPM realization implemented in our models, where a second Higgs field in the adjoint representation is added.

Throughout this section, for illustrative purposes, we consider that the bounds on proton decay rate require the effective mass of the colour triplets to be of at least $M_T^{\text{eff}} \approx 10^{17}$ GeV [18].

2.1 The missing partner mechanism

The basic idea of the missing partner mechanism (MPM) is the introduction of two new superfields Z_{50} and \bar{Z}_{50} in $\mathbf{50}$ and $\bar{\mathbf{50}}$ representations of SU(5). The decomposition of a $\mathbf{50}$ of SU(5) under the SM gauge group does not contain an SU(2) doublet, but it includes an SU(3) triplet. Thus, using the 50-plets to generate an effective mass term keeps the electroweak doublets massless, while the colour triplets acquire masses of the order of the GUT scale. The superpotential for the MPM is given by¹

$$W_{\text{MPM}} = \bar{H}_5 H_{75} Z_{50} + \bar{Z}_{50} H_{75} H_5 + M_{50} Z_{50} \bar{Z}_{50} , \quad (2.1)$$

where H_{75} is a superfield transforming in the $\mathbf{75}$ representation of SU(5), which contains a SM singlet. When H_{75} gets a vacuum expectation value (VEV) SU(5) is broken to the SM gauge group². With the triplet mass contribution from $\langle H_{75} \rangle$ denoted by V , the mass matrices of the Higgs fields H_5 , \bar{H}_5 and Z_{50} , \bar{Z}_{50} are given by

$$m_D = 0 , \quad m_T = \begin{pmatrix} 0 & V \\ V & M_{50} \end{pmatrix} , \quad (2.2)$$

for the doublet and triplet components D and T of H_5 and Z_{50} , respectively. The dangerous terms for dimension five proton decay are obtained from the Yukawa couplings

$$W_{\text{Yuk}} = \mathcal{T}_i \mathcal{F}_j \bar{H}_5 + \mathcal{T}_i \mathcal{T}_j H_5 , \quad (2.3)$$

where the families of the MSSM matter superfields are embedded in the standard way in \mathcal{T}_i and \mathcal{F}_j , transforming as $\mathbf{10}$ and $\bar{\mathbf{5}}$ of SU(5), respectively. To calculate the effective dimension five proton decay operators all Higgs triplets from 5- and 50-dimensional representations have to be integrated out, but only the triplets in the 5-dimensional representations dominantly couple to matter. We denote the triplet mass eigenvalues with \tilde{M}_1 and \tilde{M}_2 , and the corresponding mass eigenstates as \tilde{T}_1 and \tilde{T}_2 , respectively. The triplets that couple to matter are given by the combinations

$$T^{(5)} = \sum_i U_{1i}^* \tilde{T}_i , \quad \bar{T}^{(5)} = \sum_i V_{1i} \tilde{\bar{T}}_i , \quad (2.4)$$

¹For simplicity we omit most order one coefficients in the superpotentials, except where they are relevant to the discussion.

² H_{75} can be replaced by the effective combination H_{24}^2/Λ , where H_{24} is the usual GUT-breaking Higgs field in the $\mathbf{24}$ representation of SU(5) [1], see also section 2.5.

where U and V are unitary matrices defined by $m_T = U m_T^{\text{diag}} V^\dagger$. Integrating out the triplet mass eigenstates \tilde{T}_i leads to the effective dimension five operator for proton decay, which is proportional to the inverse of the “effective triplet mass”

$$(M_T^{\text{eff}})^{-1} := U_{1i}^* \left(m_T^{\text{diag}} \right)_{ij}^{-1} V_{1j} = U_{1i}^* \left(m_T^{\text{diag}} \right)_{ij}^{-1} V_{j1}^T = (m_T^{-1})_{11}. \quad (2.5)$$

With the masses given in eq. (2.2), the effective triplet mass is thus³

$$M_T^{\text{eff}} = (m_T^{-1})_{11}^{-1} = -\frac{V^2}{M_{50}}. \quad (2.6)$$

Since the doublets obtain no mass terms, the splitting of doublet mass and effective triplet mass is achieved. Using $M_T^{\text{eff}} \gtrsim 10^{17}$ GeV together with a GUT-scale value of $V \approx 10^{16}$ GeV one obtains an upper bound for $M_{50} \lesssim 10^{15}$ GeV. Having the large representations **50** and $\overline{\mathbf{50}}$ enter the Renormalization Group Equations (RGEs) at this low mass scale, however, leads to the break down of perturbativity just above the GUT scale. Thus, the MPM solves the DTS problem – but trades it for SU(5) becoming non-perturbative much below the Planck scale M_{Pl} .

2.2 The double missing partner mechanism

This trade-off can be avoided in the double missing partner mechanism (DMPM), where the number of Higgs fields in **5**, $\overline{\mathbf{5}}$, **50** and $\overline{\mathbf{50}}$ representations gets doubled [9]. The fields H_5 and \bar{H}_5 couple to the matter fields \mathcal{F}_i and \mathcal{T}_i , whereas H'_5 and \bar{H}'_5 do not. The superpotential for the DMPM is given by

$$\begin{aligned} W_{\text{DMPM}} = & \bar{H}_5 H_{75} Z_{50} + \bar{Z}_{50} H_{75} H'_5 + \bar{H}'_5 H_{75} Z'_{50} + \bar{Z}'_{50} H_{75} H_5 \\ & + M_{50} Z_{50} \bar{Z}_{50} + M'_{50} Z'_{50} \bar{Z}'_{50} \\ & + \mu' H'_5 \bar{H}'_5. \end{aligned} \quad (2.7)$$

The mass matrices of the doublet and triplet components of the Higgs fields H_5 , H'_5 , Z_{50} , Z'_{50} and their corresponding barred fields after H_{75} gets a VEV V are given by

$$m_D = \begin{pmatrix} 0 & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} 0 & 0 & 0 & V \\ 0 & \mu' & V & 0 \\ V & 0 & M_{50} & 0 \\ 0 & V & 0 & M'_{50} \end{pmatrix}. \quad (2.8)$$

While the Higgs doublets coupling to matter remain massless, the second pair of Higgs doublets contained in H'_5 and \bar{H}'_5 has mass μ' . The improvement of the DMPM compared to the MPM can be seen from the effective triplet mass M_T^{eff}

$$M_T^{\text{eff}} = (m_T^{-1})_{11}^{-1} = -\frac{V^4}{\mu' M_{50} M'_{50}}. \quad (2.9)$$

³In the text when we quote numbers for M_T^{eff} we will always refer to its modulus.




Figure 1: MPM diagram with an external S field generating the mass term for the 50-dimensional messengers after getting a VEV.

The same effective triplet mass of $M_T^{\text{eff}} \approx 10^{17}$ GeV can now be obtained while keeping high masses $M_{50} \approx M'_{50} \approx 10^{18}$ GeV, provided the heavier doublet pair has a (relatively) small mass $\mu' \approx 10^{11}$ GeV. With the large representations of SU(5) having high masses, the perturbativity of the model can be preserved up to (almost) the Planck scale.

2.3 Planck-scale suppressed operators

The philosophy we follow in this paper is to consider all Planck-scale suppressed operators allowed by the symmetries.

The superpotentials in eq. (2.1) and eq. (2.7) include mass terms for the 50-dimensional messengers. As such, one cannot use symmetries to forbid non-renormalizable Planck-scale suppressed operators such as $H_5 H_{75}^2 \bar{H}_5 / M_{\text{Pl}}$ (for the MPM, and $H_5 H_{75}^2 \bar{H}'_5 / M_{\text{Pl}}$ and $H'_5 H_{75}^2 \bar{H}_5 / M_{\text{Pl}}$ for the DMPM). These Planck-scale suppressed operators do not involve the 50-dimensional messengers and therefore generate dangerously large contributions to the masses of the doublets contained in the 5-dimensional representations, effectively spoiling the mechanism.

Given our philosophy, we must forbid these operators through a shaping symmetry. The MPM and the DMPM can then be restored by adding a singlet field S , responsible for giving mass to the 50-dimensional superfields through couplings of the form $S Z_{50} \bar{Z}_{50}$, $S Z'_{50} \bar{Z}'_{50}$ and a VEV $\langle S \rangle \neq 0$, as seen in the diagram in figure 1 (note S is not acting as an external field but generating the mass term). The non-trivial charge of S under a shaping symmetry forbids the dangerous Planck-suppressed operators.

We will generalise this strategy of generating masses for the messenger fields through an additional singlet field in sections 2.5 and 4, to avoid similarly Planck-scale suppressed operators spoiling the DMPM or the predictions for the Yukawa coupling ratios.

2.4 Yukawa coupling ratios

The problem of non-viable GUT predictions for the fermion masses in the minimal SU(5) model, such as $Y_e = Y_d^T$, can be solved through effective Yukawa couplings generated from higher dimensional operators. When the higher dimensional operators contain a GUT




Figure 2: Supergraphs generating effectively Yukawa couplings upon integrating out the pair of messenger fields R and \bar{R} .

AB	CD	R	$(Y_e)_{ji}/(Y_d)_{ij}$
$H_{24} \mathcal{F}$	$\mathcal{T} \bar{H}_{45}$	$\bar{\mathbf{45}}$	$-\frac{1}{2}$
$H_{24} \mathcal{F}$	$\mathcal{T} \bar{H}_5$	$\bar{\mathbf{5}}$	$-\frac{3}{2}$
$H_{24} \mathcal{T}$	$\mathcal{F} \bar{H}_5$	$\mathbf{10}$	6

Table 1: CG factors for the dimension five effective operators $W \supset (AB)_R(CD)_{\bar{R}}$. See the main text, Figure 2 and [11, 21] for more details.

breaking Higgs field, new ratios between the Yukawa couplings of down-type quarks and charged leptons can emerge once the GUT symmetry gets spontaneously broken [11].

Due to the introduction of extra SU(5) non-singlet fields, which participate in higher dimensional operators, there are in general several ways to construct invariants, namely multiple ways to contract the SU(5) indices. In the general case, such effective operators are not predictive and introduce an arbitrary, linear combination of several CG factors.

This issue is generic in flavour models and can be resolved by constructing a specific UV completion of the effective operators, see [19, 20] for mixing in lepton models, [11, 21] for GUT relations, and for applications in (GUT) flavour models, e.g. [15, 16, 17]. By introducing pairs of heavy messengers in the SU(5) representations \mathbf{R} and $\bar{\mathbf{R}}$, the renormalizable couplings associated with the effective operators are specified and a unique contraction of SU(5) indices of the effective operator is obtained simply by integrating out the messenger fields.

In Figure 2 and Table 1, we briefly review the topology of the diagram and field combinations that lead to the CG factors $(Y_e)_{ji}/(Y_d)_{ij} = -\frac{1}{2}, -\frac{3}{2}$ and 6 for SU(5) GUTs, which have been shown to be useful for flavour model building [11, 12, 13]. To generate these relations we use a GUT breaking Higgs field H_{24} in the adjoint representation of SU(5), and Higgs fields \bar{H}_5 and \bar{H}_{45} in the $\bar{\mathbf{5}}$ and $\bar{\mathbf{45}}$ representation. The factor $-\frac{1}{2}$ is, for instance, generated by the coupling of H_{24} and \mathcal{F} to a messenger field transforming as $\mathbf{45}$, while its partner $\bar{\mathbf{45}}$ couples to \mathcal{T} and \bar{H}_{45} . In our models in section 4 the same CG factor $-\frac{1}{2}$ is obtained from a dimension six operator where \bar{H}_{45} acts as heavy messenger field coupling to H_{24} and \bar{H}_5 .

So far we have discussed CG factors between the MSSM Yukawa couplings. For an analysis of proton decay, the CG factors to the Yukawa couplings of the colour Higgs triplets also play an important role. In Appendix A we present an extensive discussion of CG factors in SU(5), including those.

2.5 The double missing partner mechanism with an adjoint

As we discussed in the previous subsection, the CG factors we want to combine with the DMPM require the GUT breaking Higgs field to be in the adjoint representation of SU(5), $\mathbf{24}$. This motivates us to replace the Higgs field H_{75} needed for the DMPM with




Figure 3: Supergraphs generating the non-diagonal entries of the triplet mass matrix.

the effective combination H_{24}^2/Λ [1], which at the renormalizable level can be obtained by integrating out heavy messenger fields in the $\mathbf{45}$ and $\bar{\mathbf{45}}$ representations of $SU(5)$ [5]. To replace the H_{75} in the MPM, we have to introduce a set of messenger fields X_{45} , \bar{X}_{45} , Y_{45} and \bar{Y}_{45} . For the DMPM, we also need to add a second set X'_{45} , \bar{X}'_{45} , Y'_{45} and \bar{Y}'_{45} . In Figure 3 we show the supergraphs generating the non-diagonal entries of the triplet mass matrix in the DMPM with an adjoint H_{24} .

In analogy with the discussion in section 2.3, we avoid direct mass terms in order to forbid dangerous Planck-suppressed operators that would generate universal mass contributions for Higgs doublets and triplets. The messenger pairs $X_{45}\bar{X}_{45}$, $Y_{45}\bar{Y}_{45}$, $Z_{50}\bar{Z}_{50}$ and their corresponding primed versions obtain masses from the VEV of a singlet field S , charged under an additional shaping symmetry. We specify these symmetries for two example models in section 4.

One may wonder if some of these heavy 45-dimensional messengers could be the same, so that the number of fields in the spectrum would be reduced while preserving the structure of the mechanism. But if either $X_{45} \equiv Y_{45}$ or $X'_{45} \equiv Y'_{45}$, it can be seen from Figure 3 that supergraphs without the $Z_{50}\bar{Z}_{50}$ mass insertion would be allowed, spoiling the splitting of doublets and triplets and generating large non-diagonal entries in m_D . In turn, if either $X_{45} \equiv X'_{45}$, $Y_{45} \equiv Y'_{45}$ or $Z_{50} \equiv Z'_{50}$, the DMPM is reduced to the MPM, reintroducing the issue of perturbativity. Finally, an identification of $X_{45} \equiv Y'_{45}$ would allow diagrams bypassing the 50-dimensional fields, generating unwanted mass term for both doublet and triplet components of H_5 , \bar{H}_5 , and thus a too large μ -term.

With this messenger superfield content, we carefully checked that no dangerous Planck-

suppressed operators spoil the mechanism. The renormalizable superpotential is:

$$\begin{aligned}
W_{\text{DMPM}24} = & \bar{H}_5 H_{24} X_{45} + \bar{X}_{45} H_{24} Z_{50} + \bar{Z}_{50} H_{24} Y_{45} + \bar{Y}_{45} H_{24} H'_5 \\
& + \bar{H}'_5 H_{24} X'_{45} + \bar{X}'_{45} H_{24} Z'_{50} + \bar{Z}'_{50} H_{24} Y'_{45} + \bar{Y}'_{45} H_{24} H_5 \\
& + S X_{45} \bar{X}_{45} + S Y_{45} \bar{Y}_{45} + S Z_{50} \bar{Z}_{50} + S X'_{45} \bar{X}'_{45} + S Y'_{45} \bar{Y}'_{45} + S Z'_{50} \bar{Z}'_{50} \\
& + \mu' H'_5 \bar{H}'_5 .
\end{aligned} \tag{2.10}$$

After H_{24} and S obtain their VEVs and integrating out the 45-dimensional messenger fields, we find the mass matrices for the Higgs doublets and triplets to be

$$m_D = \begin{pmatrix} 0 & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} 0 & 0 & 0 & -\frac{V^2}{\langle S \rangle} \\ 0 & \mu' & -\frac{V^2}{\langle S \rangle} & 0 \\ -\frac{V^2}{\langle S \rangle} & 0 & \langle S \rangle & 0 \\ 0 & -\frac{V^2}{\langle S \rangle} & 0 & \langle S \rangle \end{pmatrix}, \tag{2.11}$$

which one can easily compare to the ones of eq. (2.8) in section 2.5. Here V is defined by $\langle H_{24} \rangle \equiv V \text{diag}(1, 1, 1, -\frac{3}{2}, -\frac{3}{2})$. Integrating out the heavy 50-dimensional fields in a next step, the mass matrices become

$$m_D = \begin{pmatrix} 0 & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} 0 & -\frac{V^4}{\langle S \rangle^3} \\ -\frac{V^4}{\langle S \rangle^3} & \mu' \end{pmatrix}. \tag{2.12}$$

Thus, the doublets in the pair $H_5 \bar{H}_5$ stay massless, while the doublet pair in $H'_5 \bar{H}'_5$ is heavy. Using only H_5 and \bar{H}_5 for the Yukawa couplings of the SM fermions, the effective triplet component mass relevant for proton decay is given by

$$M_T^{\text{eff}} = (m_T^{-1})_{11}^{-1} = -\frac{V^8}{\langle S \rangle^6 \mu'}. \tag{2.13}$$

Now the requirement of $M_T^{\text{eff}} \approx 10^{17}$ GeV can be obtained with $\langle S \rangle \approx 10^{18}$ GeV and $V \approx 2 \cdot 10^{16}$ GeV leading to $\mu' \approx \mathcal{O}(100)$ TeV. Therefore, the effective triplet mass can be large enough to stabilize the proton, while the large SU(5) representations used in the DMPM can be heavy enough to keep the theory perturbative up to the Planck scale.

When H_{24} is uncharged under additional symmetries, having μ' several orders of magnitude smaller than V requires μ' to arise from the spontaneous breakdown of a shaping symmetry, to avoid the term $\langle H_{24} \rangle H'_5 \bar{H}'_5$, which would give rise to a much too small effective triplet mass. Note that the effective triplet mass can be expressed in terms of the mass eigenstates of doublet and triplet components as $M_T^{\text{eff}} = -\frac{\tilde{M}_1 \tilde{M}_2}{\mu'}$.

2.6 Introducing a second adjoint field

We have seen that the DMPM with an adjoint GUT breaking Higgs instead of a **75** already solves the DTS problem while providing the necessary building block for the desirable CG factors for flavour model building. To conclude this section, we will argue why it is compelling to further introduce a second adjoint Higgs field:

- In the minimal SUSY SU(5) model [22], the single GUT breaking **24** contains an SU(2) triplet component and an SU(3) octet component with equal masses. Demanding gauge coupling unification, the mass of the Higgs colour triplets is required to be about 10^{15} GeV [6], ruling out this model due to proton decay. Non-renormalizable operators in the GUT breaking superpotential can split the **24** component masses, allowing a higher effective triplet mass [23] (see also section 3). An additional **24** can be used to realize this non-renormalizable superpotential in a renormalizable way.
- It turns out that the introduction of an additional **24** is not just a UV-completion of the non-renormalizable superpotential of [23]. When both adjoints have approximately the same mass and therefore the second **24** is not integrated out, the additional colour octet and electroweak triplet in the spectrum lead to more freedom for the GUT scale and effective triplet mass. In the following section we discuss all possible renormalizable superpotentials with two adjoints and their impact on M_{GUT} and M_T^{eff} from a gauge coupling unification analysis.
- A renormalizable superpotential for one **24** requires it to be uncharged under shaping symmetries in order for it to obtain a VEV. However, such a shaping symmetry charge is vital in the type of flavour models considered here, to avoid unwanted admixtures of additional CG factors involving less insertions of H_{24} . With a second **24**, the adjoint fields can acquire non-vanishing VEVs even when charged under shaping symmetries.

These features of renormalizable superpotentials for two adjoints are presented in detail in the next section.

3 Grand unification and the effective triplet mass

In GUT extensions of the SM it is quite common to have additional fields below the GUT scale that modify the RGE running, as it is the case for the class of models in this paper. Therefore one has to study the impact of the additional fields on the running of the gauge couplings and especially study their unification. The modified unification condition for the gauge couplings at one-loop reads

$$\frac{1}{\alpha_u} = \frac{1}{\alpha_i} - \frac{1}{2\pi} \left(b_i^{(\text{SM})} \log \frac{M_{\text{SUSY}}}{M_Z} + b_i^{(\text{MSSM})} \log \frac{M_{\text{GUT}}}{M_{\text{SUSY}}} + \sum_f b_i^{(f)} \log \frac{M_{\text{GUT}}}{M_f} \right), \quad (3.1)$$

where $i = 1, 2, 3$ labels the SM gauge interaction and f labels the additional superfields (compared to the MSSM), with masses M_f and β coefficients $b_i^{(f)}$. The one-loop β -function coefficients for the SM are $b_i^{(\text{SM})} = (41/10, -19/6, -7)$ and for the MSSM $b_i^{(\text{MSSM})} = (33/5, 1, -3)$. The SUSY scale M_{SUSY} is defined here as the scale where we make the transition from the SM β coefficients to the MSSM ones. The α_i are defined at low energies $\alpha_i \equiv \alpha_i(M_Z)$ while α_u is the unified gauge coupling at the GUT scale $\alpha_u \equiv \alpha_i(M_{\text{GUT}})$. The GUT scale M_{GUT} is defined here as the scale where the last SU(5)

multiplet is completed, in other words the scale where all three one-loop β coefficients for the SM gauge couplings become equal.

From now on, we assume that the heaviest incomplete SU(5) multiplets to enter the RGE running are the leptoquark vector bosons, such that the GUT scale corresponds to their mass $M_{\text{GUT}} = M_V$. While other cases can certainly arise, we focus on this option because it is quite common in our setup to have heavy leptoquark vector bosons, and furthermore we verified that in this case the triplet mass can be made very heavy as well.

In addition to the MSSM field content, the DMPM introduces one additional pair of SU(2)-doublets, $D^{(5)}$ and $\bar{D}^{(5)}$ and two additional pairs of SU(3)-triplets, $T_i^{(5)}$ and $\bar{T}_i^{(5)}$, $i = 1, 2$. They enter the β -functions with the coefficients $b_i^{(5,D)} = (3/5, 1, 0)$ for the Dirac pair of doublets and $b_i^{(5,T)} = (2/5, 0, 1)$ per Dirac pair of triplet and anti-triplet.

Furthermore, we use two SU(5) breaking GUT Higgs fields H_{24} and H'_{24} in the adjoint representation. They contain one SM-singlet component each, one SU(2)-triplet, $T^{(24)}$, with $b_i^{(24,T)} = (0, 2, 0)$, an SU(3)-octet, $O^{(24)}$ with $b_i^{(24,O)} = (0, 0, 3)$ and a leptoquark superfield pair, $L^{(24)}$ with $b_i^{(24,L)} = (5, 3, 2)$. Since one leptoquark superfield pair is eaten up during the breaking of SU(5), we are left with two triplets with masses $M_{T_1^{(24)}}$ and $M_{T_2^{(24)}}$, two octets with masses $M_{O_1^{(24)}}$ and $M_{O_2^{(24)}}$ and one leptoquark superfield pair with mass $M_{L^{(24)}}$.

For convenience we define the geometric means of the masses $M_{T^{(5)}}^2 = M_{T_1^{(5)}} M_{T_2^{(5)}}$ for the colour triplets and analogously $M_{T^{(24)}}^2 = M_{T_1^{(24)}} M_{T_2^{(24)}}$, $M_{O^{(24)}}^2 = M_{O_1^{(24)}} M_{O_2^{(24)}}$ for the components of H_{24} and H'_{24} .

Having this at hand, we can solve eq. (3.1) for $M_{D^{(5)}}$, $M_{T^{(5)}}$ and M_{GUT} ,⁴

$$\begin{aligned} \log M_{D^{(5)}} &= \frac{15\pi}{4\alpha_1} - \frac{17\pi}{4\alpha_2} - \frac{3\pi}{2\alpha_3} + \frac{59}{3} \log M_Z \\ &+ \frac{2\pi}{\alpha_u} + \frac{3}{2} \log M_{L^{(24)}} - \frac{17}{2} \log M_{T^{(24)}} - \frac{9}{2} \log M_{O^{(24)}} - \frac{43}{6} \log M_{\text{SUSY}} , \end{aligned} \quad (3.2)$$

$$\begin{aligned} \log M_{T^{(5)}} &= \frac{35\pi}{24\alpha_1} - \frac{7\pi}{8\alpha_2} - \frac{19\pi}{12\alpha_3} + \frac{119}{12} \log M_Z \\ &+ \frac{\pi}{\alpha_u} + \frac{3}{4} \log M_{L^{(24)}} - \frac{7}{4} \log M_{T^{(24)}} - \frac{19}{4} \log M_{O^{(24)}} - \frac{19}{6} \log M_{\text{SUSY}} , \end{aligned} \quad (3.3)$$

$$\begin{aligned} \log M_{\text{GUT}} &= \frac{5\pi}{12\alpha_1} - \frac{\pi}{4\alpha_2} - \frac{\pi}{6\alpha_3} + \frac{11}{6} \log M_Z \\ &+ \frac{1}{2} \log M_{L^{(24)}} - \frac{1}{2} \log M_{T^{(24)}} - \frac{1}{2} \log M_{O^{(24)}} - \frac{1}{3} \log M_{\text{SUSY}} . \end{aligned} \quad (3.4)$$

For the study of proton decay, it is more convenient to instead solve eq. (3.1) for the GUT scale gauge coupling α_u and the effective triplet mass $M_T^{\text{eff}} = M_{T^{(5)}}^2 / M_{D^{(5)}}$, which gives the suppression of the dimension 5 proton decay operators (cf. the discussion in

⁴In the following, “log” of a mass is to be understood as the natural logarithm of the mass divided by one common mass scale, e.g. $\log m \equiv \log(m/\text{GeV})$.

section 2.5). Then we get the relations

$$\frac{\pi}{\alpha_u} = -\frac{43\pi}{24\alpha_1} + \frac{15\pi}{8\alpha_2} + \frac{11\pi}{12\alpha_3} - \frac{197}{20} \log M_Z + \frac{3}{5} \log M_{DT} \quad (3.5)$$

$$- \frac{3}{4} \log M_{L^{(24)}} + \frac{15}{4} \log M_{T^{(24)}} + \frac{11}{4} \log M_{O^{(24)}} + \frac{7}{2} \log M_{\text{SUSY}} ,$$

$$\log M_T^{\text{eff}} = -\frac{5\pi}{6\alpha_1} + \frac{5\pi}{2\alpha_2} - \frac{5\pi}{3\alpha_3} + \frac{1}{6} \log M_Z \quad (3.6)$$

$$+ 5 \log M_{T^{(24)}} - 5 \log M_{O^{(24)}} + \frac{5}{6} \log M_{\text{SUSY}} ,$$

$$\log M_{\text{GUT}} = \frac{5\pi}{12\alpha_1} - \frac{\pi}{4\alpha_2} - \frac{\pi}{6\alpha_3} + \frac{11}{6} \log M_Z \quad (3.7)$$

$$+ \frac{1}{2} \log M_{L^{(24)}} - \frac{1}{2} \log M_{T^{(24)}} - \frac{1}{2} \log M_{O^{(24)}} - \frac{1}{3} \log M_{\text{SUSY}} ,$$

where we have introduced the mass $M_{DT}^3 = M_{D^{(5)}}^2 M_{T^{(5)}}$. As one can see only α_u depends on M_{DT} , which is due to the fact that doublets and colour triplets together form a complete representation of SU(5). Thus, following eq. (3.1), one can see that a simultaneous rescaling $M_{D^{(5)}} \rightarrow q^2 M_{D^{(5)}}$ and $M_{T^{(5)}} \rightarrow q M_{T^{(5)}}$ leaves the GUT scale invariant and only shifts α_u , while $M_T^{\text{eff}} \propto q^0$ remains unchanged and $M_{DT} \propto q$ parametrises this rescaling. Further interdependencies between α_u , M_T^{eff} and M_{GUT} are then implicit via their shared dependence on the other masses.

Thus, unification implies that the effective triplet mass follows the relation

$$M_T^{\text{eff}} = \exp\left(\frac{5}{6}\pi\left(\frac{3}{\alpha_2} - \frac{2}{\alpha_3} - \frac{1}{\alpha_1}\right)\right) M_Z^{1/6} M_{\text{SUSY}}^5 \left(\frac{M_{T^{(24)}}}{M_{O^{(24)}}}\right)^5$$

$$= 2.5_{-0.8}^{+0.6} \cdot 10^{17} \text{ GeV} \left(\frac{M_{\text{SUSY}}}{1 \text{ TeV}}\right)^{\frac{5}{6}} \left(\frac{M_{T^{(24)}}}{M_{O^{(24)}}}\right)^5 , \quad (3.8)$$

while the GUT scale is given by

$$M_{\text{GUT}} = 1.37_{-0.05}^{+0.05} \cdot 10^{16} \text{ GeV} \left(\frac{M_{\text{SUSY}}}{1 \text{ TeV}}\right)^{-\frac{1}{3}} \left(\frac{M_{L^{(24)}}}{10^{16} \text{ GeV}}\right)^{\frac{1}{2}} \left(\frac{M_{T^{(24)}} M_{O^{(24)}}}{(10^{16} \text{ GeV})^2}\right)^{-\frac{1}{2}} . \quad (3.9)$$

For completeness, the unified gauge coupling is given by

$$\frac{1}{\alpha_u} = 24.58 \pm 0.06 + \frac{7}{2\pi} \ln \frac{M_{\text{SUSY}}}{1 \text{ TeV}} + \frac{3}{5\pi} \ln \frac{M_{DT}}{10^{14} \text{ GeV}} \quad (3.10)$$

$$- \frac{3}{4\pi} \ln \frac{M_{L^{(24)}}}{10^{16} \text{ GeV}} + \frac{15}{4\pi} \ln \frac{M_{T^{(24)}}}{10^{16} \text{ GeV}} + \frac{11}{4\pi} \ln \frac{M_{O^{(24)}}}{10^{16} \text{ GeV}} .$$

For these numbers, we have used the experimental values and uncertainties for the gauge couplings found in [10]. Note that for all three quantities the resulting uncertainty is dominated by the experimental error on α_s . In the following we will not quote any errors on the masses anymore since the relative uncertainty changes only negligibly for the different

superpotentials and for two-loop running. The reference scale 10^{14} GeV is chosen due to the fact that $M_{DT} = 10^{14}$ GeV and $M_{T(5)} = 10^{16}$ GeV implies $M_T^{\text{eff}} = 10^{19}$ GeV.

Since the effective triplet mass M_T^{eff} receives significant two-loop contributions (cf., for instance, [6]), we have also implemented a numerical two-loop RGE analysis using the following procedure. We start with SM values for the gauge and Yukawa couplings [24] at M_Z , run up to to a scale of 1 TeV with the full two-loop SM RGEs and match the SM to the MSSM (including $\overline{\text{MS}}$ to $\overline{\text{DR}}$ scheme conversion). From there we run and match using full two-loop MSSM RGEs and one-loop gauge coupling threshold corrections⁵ while step-by-step including all additional multiplets at their mass scale via their contributions to the one- and two-loop gauge coupling RGEs, see Appendix B. The Yukawa couplings of the Higgs colour triplets are well approximated by using the Yukawa couplings of the corresponding doublets. We do not take into account any other Yukawa couplings. We verified this approximation numerically and the results for the colour triplet masses are barely affected.

3.1 Superpotentials with two adjoints of SU(5)

In this section, we systematically study all superpotentials with two adjoints that can break SU(5) to the SM gauge group. We find only four possibilities with non-vanishing VEVs and masses. Classified based on their symmetry, they are:

- (a) $W = M_{24} \text{tr} H_{24}^2 + M'_{24} \text{tr} H_{24}'^2 + \kappa' \text{tr} H_{24} H_{24}'^2 + \lambda \text{tr} H_{24}^3$,
 \mathbb{Z}_2 symmetry where H_{24} is uncharged and H_{24}' charged.
- (b) $W = \tilde{M}_{24} \text{tr} H_{24} H_{24}' + \lambda \text{tr} H_{24}^3 + \lambda' \text{tr} H_{24}'^3$,
 \mathbb{Z}_3 symmetry, where H_{24} has charge 2 and H_{24}' charge 1.
- (c) $W = \tilde{M}_{24} \text{tr} H_{24} H_{24}' + \lambda \text{tr} H_{24}^3 + \kappa' \text{tr} H_{24} H_{24}'^2$,
 \mathbb{Z}_4^R symmetry where H_{24} has a charge of 2 (with $q_\theta = 1$) and H_{24}' is uncharged.
- (d) The trivial case with both fields only charged under SU(5) and all (non-linear) terms allowed. We will not consider this case any further.

Since we are dealing with two adjoint Higgs fields, it is convenient to define a quantity $\tan \beta_V$ similar to $\tan \beta$ of the MSSM, so that

$$\langle H_{24} \rangle = V_1 e^{i\phi_1} \text{diag}(1, 1, 1, -3/2, -3/2), \quad (3.11)$$

$$\langle H_{24}' \rangle = V_2 e^{i\phi_2} \text{diag}(1, 1, 1, -3/2, -3/2), \quad (3.12)$$

with $V_1, V_2 > 0$ and $\tan \beta_V = V_1/V_2$.

3.1.1 Superpotential (a)

We will begin our discussion with superpotential (a) which turns out to be the most complicated case since it has the most parameters. As it contains two mass parameters,

⁵When one integrates out particles at a threshold scale equal to their mass, these threshold corrections vanish, as can be seen in [25].




Figure 4: The effective colour triplet mass M_T^{eff} (left) and GUT scale M_{GUT} (right) in GeV at one-loop (upper) and two-loop (lower) order as resulting from superpotential (a) for $M_{\text{SUSY}} = 1$ TeV, $M = 10^{15}$ GeV and $\bar{\phi} = 0$. Note the different colour coding between left and right. For illustration, the white strips denote areas with light $M_{T(24)}$ or $M_{O(24)}$ ($< 10^{13}$ GeV). Such relatively low values for these components can arise either from cancellation between terms, or from a generic suppression due to small parameters, cf. eqs. (3.14).

we introduce a second angle β_M and mean mass $M > 0$ such that $M_{24} = Me^{i\alpha_1} \sin \beta_M$ and $M'_{24} = Me^{i\alpha_2} \cos \beta_M$. The vacuum expectation values are given by

$$V_1 e^{i\phi_1} = \frac{4M'_{24}}{\kappa'} \quad \text{and} \quad V_2 e^{2i\phi_2} = 4M'_{24} \frac{2M_{24}\kappa' - 3M'_{24}\lambda}{\kappa'^3}, \quad (3.13)$$

which can as well be expressed in terms of a ratio of the coupling constants involved, $3\lambda/\kappa' = 2e^{i(\alpha_1 - \alpha_2)} \tan \beta_M - e^{-2i(\phi_1 - \phi_2)} \cot^2 \beta_V$. For the geometric means of the masses of the additional fields compared to the MSSM, we find

$$M_{T^{(24)}}^2 = 5M^2 \cos \beta_M \sqrt{(2 \cos \beta_M - 3 \sin \beta_M \tan^2 \beta_V)^2 + \Delta}, \quad (3.14a)$$

$$M_{O^{(24)}}^2 = 5M^2 \cos \beta_M \sqrt{(3 \cos \beta_M - 2 \sin \beta_M \tan^2 \beta_V)^2 + \Delta}, \quad (3.14b)$$

$$M_{L^{(24)}}^2 = \frac{1}{4} M^2 \frac{\cos^2 \beta_M}{\sin^4 \beta_V}, \quad (3.14c)$$

with $\Delta = 12 \sin(2\beta_M) \cot^2(\beta_V) \sin^2 \bar{\phi}$ and $\bar{\phi} = (\alpha_1 - \alpha_2)/2 + \phi_1 - \phi_2$. Note that not only the geometric mean masses, but also the mass eigenvalues themselves only depend on this phase combination $\bar{\phi}$ and are invariant under $\bar{\phi} \rightarrow \bar{\phi} + \pi$.

The effective triplet mass as of eq. (3.8) is heaviest if the phase $\bar{\phi}$ is 0, π or 2π , since then the ratio $M_{T^{(24)}}/M_{O^{(24)}}$ is not bounded from above (or below), which allows for the maximal range for M_T^{eff} . Thus, in the following, we choose $\bar{\phi}$ and thus Δ to vanish.

The resulting plots for M_T^{eff} and for M_{GUT} are shown in Fig. 4 for $M_{\text{SUSY}} = 1$ TeV, $M = 10^{15}$ GeV and $\bar{\phi} = 0$, including a comparison between one- and two-loop results.

3.1.2 Superpotential (b) and (c)

These two superpotentials have only one massive parameter $\tilde{M}_{24} = Me^{i\alpha}$ and hence the analytic results become much less cumbersome. The vacuum solutions are given by

$$V_1 e^{i\phi_1} = \frac{2}{3} \frac{Me^{i\alpha}}{\sqrt[3]{\lambda^2 \lambda'}} \quad \text{and} \quad V_2 e^{i\phi_2} = \frac{2}{3} \frac{Me^{i\alpha}}{\sqrt[3]{\lambda \lambda'^2}}, \quad (3.15)$$

up to a \mathbb{Z}_3 symmetry transformation for superpotential (b) and

$$V_1 e^{i\phi_1} = \frac{1}{\sqrt{3}} \frac{Me^{i\alpha}}{\sqrt{\lambda \kappa'}} \quad \text{and} \quad V_2 e^{i\phi_2} = \frac{Me^{i\alpha}}{\kappa'}, \quad (3.16)$$

up to a minus sign in V_1 for superpotential (c). In other words, the couplings fulfill the relation $\lambda'/\lambda = e^{3i(\phi_2 - \phi_1)} \tan^3 \beta_V$ for superpotential (b) and $\kappa'/\lambda = 3e^{2i(\phi_1 - \phi_2)} \tan^2 \beta_V$ for superpotential (c).

For the geometric means of the masses of the colour triplets, octets and the mass of the left-over leptoquark superfield in H_{24} and H'_{24} , we find

$$M_{T^{(24)}}^2 = \frac{35}{4} M^2, \quad M_{O^{(24)}}^2 = \frac{15}{4} M^2 \quad \text{and} \quad M_{L^{(24)}}^2 = \frac{1}{\sin^2(2\beta_V)} M^2, \quad (3.17)$$

for superpotential (b) and

$$M_{T^{(24)}}^2 = \frac{5}{4} M^2, \quad M_{O^{(24)}}^2 = \frac{5}{4} M^2 \quad \text{and} \quad M_{L^{(24)}}^2 = \frac{1}{4 \sin^2(2\beta_V)} M^2, \quad (3.18)$$

for superpotential (c). Note that in both cases also all mass eigenvalues turn out to be phase independent. Therefore, applying eq. (3.8) unification of the gauge couplings on one-loop implies

$$M_T^{\text{eff, 1-loop}} = 2.5 \cdot 10^{17} \text{ GeV} \left(\frac{M_{\text{SUSY}}}{1 \text{ TeV}} \right)^{\frac{5}{6}} \times \begin{cases} \left(\frac{7}{3}\right)^{\frac{5}{2}} \approx 8.3 & \text{(b)} \\ 1 & \text{(c)} \end{cases}, \quad (3.19)$$

and

$$M_{\text{GUT}}^{\text{1-loop}} = \frac{1.37 \cdot 10^{16} \text{ GeV}}{\sqrt{|\sin 2\beta_V|}} \left(\frac{M_{\text{SUSY}}}{1 \text{ TeV}} \right)^{-\frac{1}{3}} \left(\frac{M}{10^{15} \text{ GeV}} \right)^{-\frac{1}{2}} \times \begin{cases} \sqrt{\frac{8}{\sqrt{21}}} \approx 1.32 & \text{(b)} \\ 2 & \text{(c)} \end{cases}. \quad (3.20)$$

Assuming the same parameters, we find an almost ten times heavier effective triplet mass in superpotential (b) than in (c) and hence we focus on this case in our second example model later.

At two-loop, we find the following approximate behaviour for the masses

$$M_T^{\text{eff, 2-loop}} = \left(\frac{M_{\text{SUSY}}}{1 \text{ TeV}} \right)^{0.74} \cdot \begin{cases} 5.4 \cdot 10^{16} \text{ GeV} \left(\frac{M}{10^{15} \text{ GeV}} \right)^{-0.15} & \text{(b)} \\ 7.1 \cdot 10^{15} \text{ GeV} \left(\frac{M}{10^{15} \text{ GeV}} \right)^{-0.19} & \text{(c)} \end{cases}, \quad (3.21)$$

and

$$M_{\text{GUT}}^{\text{2-loop}} = |\sin 2\beta_V|^{-0.48} \left(\frac{M_{\text{SUSY}}}{1 \text{ TeV}} \right)^{-0.41} \cdot \begin{cases} 2.95 \cdot 10^{16} \text{ GeV} \left(\frac{M}{10^{15} \text{ GeV}} \right)^{-0.62} & \text{(b)} \\ 4.89 \cdot 10^{16} \text{ GeV} \left(\frac{M}{10^{15} \text{ GeV}} \right)^{-0.64} & \text{(c)} \end{cases}. \quad (3.22)$$

The dependence on other parameters is very small. As we can see, at two-loop, having $M_T^{\text{eff}} \gtrsim 10^{17} \text{ GeV}$ requires $M_{\text{SUSY}} \gtrsim 2.3 \text{ TeV}$ and 35 TeV for superpotential (b) and (c) respectively.

There are a few more comments in order. There is a claim [26] that there is no four-dimensional (SUSY) GUT which has an R-symmetry and solves the DTS problem. This seems to be in conflict with our superpotentials, but it is not, because the R-symmetry is absent (a,b,d) or broken spontaneously at the GUT scale (c), which was not considered therein. Furthermore, superpotential (c) is particularly interesting for model building purposes because R-symmetries are very popular in flavour models with non-Abelian family symmetries (and spontaneous CP violation). We will discuss this in more detail in appendix D.

4 Flavour Models with DMPM

In this section we combine the DMPM (featuring two adjoints of $SU(5)$) with a predictive GUT flavour model for the quark-lepton Yukawa ratios at the GUT scale. In particular, we implement CG factors as given in [11]. Two examples with different Yukawa matrix structures are presented: in the first model we construct diagonal down-type quark and charged lepton Yukawa matrices Y_d and Y_e , with all mixing originating from the up-type quark Yukawa matrix Y_u . The second model realizes the attractive feature of the Cabibbo mixing angle θ_C originating from Y_d . Both models are providing existence proofs that successful DTS and experimentally viable predictions for the GUT scale Yukawa coupling ratios can indeed be realised simultaneously in one model.

Let us be more specific on the predictions made by the two models: Due to the CG factors in the down-type quark and charged lepton sector, $\frac{y_\tau}{y_b}$, $\frac{y_\mu}{y_s}$ and $\frac{y_e}{y_d}$ are predicted at the GUT scale. To confront them with the experimental data, the RG running to low energies has to be performed, including in particular supersymmetric 1-loop threshold corrections [27] when the MSSM is matched to the SM. These threshold corrections can have a sizeable impact on the low energy values of the Yukawa couplings (and thus the fermion masses), depending on the sparticle spectrum and $\tan\beta$. So the predictions here are two-fold: Firstly, the predictions for the Yukawa ratios at the GUT scale imply constraints on the SUSY breaking parameters, which may be tested at future collider searches if SUSY is found. Secondly, the ratios $\frac{y_d}{y_s}$ and $\frac{y_e}{y_\mu}$ are not affected by RG running and by the SUSY threshold corrections (as long as the first two families of sfermions are almost degenerate in mass as commonly assumed). They can be directly used to constrain GUT models. A particularly useful quantity in this context is indeed the double ratio

$$\left| \frac{y_\mu y_d}{y_s y_e} \right| = 10.7_{-0.8}^{+1.8}, \quad (4.1)$$

which can be checked directly at the GUT scale [24]. In our models we will use the CG factors $\frac{y_\tau}{y_b} = -\frac{3}{2}$, $\frac{y_\mu}{y_s} = 6$ and $\frac{y_e}{y_d} = -\frac{1}{2}$. This leads to $\left| \frac{y_\mu y_d}{y_s y_e} \right| = 12$, which is in good agreement with the phenomenological value. On the other hand, the ubiquitous CG factors $y_\mu = -3y_s$ and $y_e = \frac{1}{3}y_d$, known as Georgi-Jarlskog relations [3], would give $\left| \frac{y_\mu y_d}{y_s y_e} \right| = 9$ and deviate from the central value 10.7 by more than two sigma.

In this section we explicitly construct only the Yukawa matrices of the up- and down-type quarks and charged leptons. Adding one of the ubiquitous mechanisms to generate neutrino masses and lepton mixing angles would be straightforward. However, we do not consider neutrinos in this paper, since they are not directly relevant for the discussion of proton decay, doublet-triplet splitting and the CG factors between Y_d and Y_e .

4.1 A model with diagonal Y_d and Y_e Yukawa matrices

We now turn to our first model featuring diagonal down-type quark and charged lepton Yukawa matrices Y_d and Y_e .⁶ In this case all the mixing in the quark sector has to come exclusively from the up-type quark Yukawa matrix Y_u . Explicitly, we have the following structure for the Yukawa matrices

$$Y_d = \begin{pmatrix} y_d & 0 & 0 \\ 0 & y_s & 0 \\ 0 & 0 & y_b \end{pmatrix}, Y_e = \begin{pmatrix} -\frac{1}{2}y_d & 0 & 0 \\ 0 & 6y_s & 0 \\ 0 & 0 & -\frac{3}{2}y_b \end{pmatrix}, Y_u = \begin{pmatrix} y_{11} & y_{12} & y_{13} \\ y_{12} & y_{22} & y_{23} \\ y_{13} & y_{23} & y_{33} \end{pmatrix}. \quad (4.2)$$

An approach to flavour (GUT) model building with diagonal Y_e (and Y_d) has been discussed recently in [28].

We introduce flavon fields θ_1 , θ_2 , θ_3 and θ_4 that obtain a VEV and generate the hierarchical structure of the Yukawa matrices. After the flavon fields, H_{24} and H'_{24} obtain their VEVs, the Yukawa matrices of eq. (4.2) originate from the following effective superpotentials

$$W_u = \frac{1}{\Lambda^4} H_5 \mathcal{T}_1 \mathcal{T}_1 \theta_1^2 \theta_2^2 + \frac{1}{\Lambda^3} H_5 \mathcal{T}_1 \mathcal{T}_2 \theta_1^2 \theta_2 + \frac{1}{\Lambda^2} H_5 \mathcal{T}_1 \mathcal{T}_3 \theta_1 \theta_2 + \frac{1}{\Lambda^2} H_5 \mathcal{T}_2 \mathcal{T}_2 \theta_1^2 + \frac{1}{\Lambda} H_5 \mathcal{T}_2 \mathcal{T}_3 \theta_1 + H_5 \mathcal{T}_3 \mathcal{T}_3, \quad (4.3)$$

$$W_d = \frac{1}{\langle S' \rangle} (H'_{24} \mathcal{F}_3)_{\bar{5}} (\bar{H}_5 \mathcal{T}_3)_5 + \frac{\theta_3}{\langle S' \rangle^2} (H'_{24} \mathcal{T}_2)_{10} (\bar{H}_5 \mathcal{F}_2)_{\bar{10}} + \frac{\theta_4}{\langle S' \rangle^2 \langle S \rangle} (H'_{24} \mathcal{F}_1)_{\bar{45}} (\mathcal{T}_1 H_{24} \bar{H}_5)_{45}, \quad (4.4)$$

where we do not show order one coefficients, and denote the different messenger masses generating W_u by a generic Λ . However, keep in mind that this is just for the sake of simplicity and different entries in the Yukawa matrix should be understood as independent parameters. The ratios of flavon VEVs and messenger masses is small of about 0.01 – 0.1. For a list of all fields including their charges under the additional discrete shaping symmetries, see Tables 2, 3, 4, and 5.

The adjoint H'_{24} required to construct the desired CG factors must be charged under shaping symmetries. We will therefore implement superpotential (a). Note that it leaves the second adjoint H_{24} uncharged, which could in principle lead to a problem. A direct mass term of messenger fields $M_i Z_i \bar{Z}_i$ would in this case always show a up with a term of the form $H_{24} Z_i \bar{Z}_i$. Such a contribution would inevitably spoil the desired CG factors [21] between Y_d and Y_e^T as long as the mass and the adjoint VEV are not very hierarchical. To avoid this and still generate the desired operators the masses of the messenger fields that give rise to W_d in eq. (4.4) originate from the VEVs of the fields S and S' charged under the shaping symmetry (but with different charges than H'_{24})⁷.

⁶The matrices are diagonal in the preferred basis where the different fermion generations have well defined symmetry assignments, cf. Table 2.

⁷The VEV of the charged singlet S gives masses to the heavy messengers of the DMPM (see section 2.5).

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_9	\mathbb{Z}_2
H_5	$\mathbf{5}$
\bar{H}_5	$\bar{\mathbf{5}}$.	2	.	.	.	1	2	.
\mathcal{T}_1	$\mathbf{10}$.	.	3	.	6	.	.	1
\mathcal{T}_2	$\mathbf{10}$	6	.	.	1
\mathcal{T}_3	$\mathbf{10}$	1
\mathcal{F}_1	$\bar{\mathbf{5}}$	1	1	1	2	1	1	2	1
\mathcal{F}_2	$\bar{\mathbf{5}}$	1	.	.	2	1	.	2	1
\mathcal{F}_3	$\bar{\mathbf{5}}$	1	2	.	1	.	6	7	1

Table 2: SU(5) representations and charges under discrete shaping symmetries of the MSSM fields and colour triplets of the model presented in subsection 4.1. A dot denotes charge zero.

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_9	\mathbb{Z}_2
H_{24}	$\mathbf{24}$
H'_{24}	$\mathbf{24}$	1
S	$\mathbf{1}$.	3	.	.	.	2	.	.
S'	$\mathbf{1}$.	.	.	1
θ_1	$\mathbf{1}$	1	.	.	.
θ_2	$\mathbf{1}$.	.	1
θ_3	$\mathbf{1}$.	2	.	.	.	6	5	.
θ_4	$\mathbf{1}$	5	.

Table 3: SU(5) representations and charges under discrete shaping symmetries of the superfields obtaining VEVs at around the GUT scale of the model presented in subsection 4.1. A dot denotes charge zero.




Figure 5: Supergraphs leading to the effective superpotential W_d of eq. (4.4) when the heavy messenger fields get integrated out in the model presented in subsection 4.1.

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_9	\mathbb{Z}_2
H'_5	5	.	3	.	.	.	5	7	.
\bar{H}'_5	$\bar{5}$.	1	.	.	.	6	.	.
X_{45}	45	.	2	.	.	.	6	7	.
\bar{X}_{45}	$\bar{45}$.	3	.	.	.	6	2	.
Y_{45}	45	3	7	.
\bar{Y}_{45}	$\bar{45}$.	1	.	.	.	2	2	.
Z_{50}	50	.	1	.	.	.	1	7	.
\bar{Z}_{50}	$\bar{50}$	4	2	.
X'_{45}	45	.	3	.	.	.	1	.	.
\bar{X}'_{45}	$\bar{45}$.	2	.	.	.	4	.	.
Y'_{45}	45	.	1	.	.	.	5	.	.
\bar{Y}'_{45}	$\bar{45}$
Z'_{50}	50	.	2	.	.	.	3	.	.
\bar{Z}'_{50}	$\bar{50}$.	3	.	.	.	2	.	.

Table 4: SU(5) representations and charges under discrete shaping symmetries of the fields in the DMPM sector of the model presented in subsection 4.1. A dot denotes charge zero.

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_4	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_9	\mathbb{Z}_2
$Z_{5,1}$	5	.	2	.	3	.	1	2	1
$\bar{Z}_{5,1}$	$\bar{5}$.	2	.	.	.	6	7	1
$Z_{10,1}$	10	1	.	.	3	6	.	.	1
$\bar{Z}_{10,1}$	$\bar{10}$	1	.	.	.	1	.	.	1
$Z_{10,2}$	10	1	2	.	2	6	6	5	1
$\bar{Z}_{10,2}$	$\bar{10}$	1	2	.	1	1	1	4	1
$Z_{45,1}$	45	.	3	3	2	6	6	7	1
$\bar{Z}_{45,1}$	$\bar{45}$.	1	1	1	1	1	2	1
$Z_{45,2}$	45	.	3	3	3	6	6	2	1
$\bar{Z}_{45,2}$	$\bar{45}$.	1	1	.	1	1	7	1
$Z_{10,3}$	10	1	.	.	1
$Z_{10,4}$	10	.	.	1	.	1	.	.	1
Z_1	1	5	.	.	.
Z_2	1	.	.	3	.	5	.	.	.

Table 5: SU(5) representations and charges under discrete shaping symmetries of the flavon and flavour messenger fields of the model presented in subsection 4.1. Note that the messengers $Z_{5,1}\bar{Z}_{5,1}$, $Z_{10,1}\bar{Z}_{10,1}$, $Z_{10,2}\bar{Z}_{10,2}$, $Z_{45,1}\bar{Z}_{45,1}$ and $Z_{45,2}\bar{Z}_{45,2}$ have no direct mass term, but get their masses through the VEVs of S and S' . The other messenger fields have direct mass terms, so their corresponding barred field is omitted in this table. A dot denotes charge zero.




Figure 6: Supergraphs leading to the effective superpotential W_u of eq. (4.3) when the heavy messenger fields are integrated out in the model presented in subsection 4.1. Note that there are three supergraphs contributing to the superpotential term generating y_{12} . A detailed discussion of the messenger sector is presented in the appendix C.

The full messenger sector can be read off from the supergraphs presented in Figures 5 and 6, see also Table 5. After the heavy messenger fields get integrated out, the effective superpotentials W_d and W_u are obtained.

The predictivity of the down-type quarks and charged lepton Yukawa couplings has already been discussed above. Note that the hierarchical structure is enforced due to the use of higher order effective operators in W_d . In a small angle approximation the leading order estimates for the eigenvalues of Y_u and the mixing angles are given as

$$y_u \approx y_{11} - \frac{y_{12}^2}{y_{22}}, \quad y_c \approx y_{22}, \quad y_t \approx y_{33}, \quad \theta_C \approx \frac{y_{12}}{y_{22}}, \quad \theta_{23} \approx \frac{y_{23}}{y_{33}}, \quad \theta_{13} \approx \frac{y_{13}}{y_{33}}. \quad (4.5)$$

Phenomenology requires that all parameters y_{ij} of Y_u are independent, which needs to be carefully considered in the construction of the messenger sector, as discussed in Appendix C.

In section 2.5 we argued that for the case of an uncharged H_{24} , as it appears in the selected superpotential (a), the mass term for the additional, five-dimensional Higgs fields must come from the VEV of some singlet field. In our model an effective μ' term is generated from a higher-dimensional operator and with an even higher suppression, there is also a μ -term for the Higgs fields coupling to matter,

$$W_5^{\text{eff}} = \mu H_5 \bar{H}_5 + \mu' H_5' \bar{H}_5', \quad (4.6)$$

where $\mu' \equiv \langle \theta_3 \rangle^4 / M_{\text{Pl}}^3$ and $\mu \equiv \langle \theta_3 \rangle \langle \theta_4 \rangle^4 / M_{\text{Pl}}^4$.⁸ The mass matrices for the doublet and triplet components are then given by

$$m_D = \begin{pmatrix} \mu & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} \mu & 0 & 0 & \frac{V^2}{\langle S \rangle} \\ 0 & \mu' & \frac{V^2}{\langle S \rangle} & 0 \\ \frac{V^2}{\langle S \rangle} & 0 & \langle S \rangle & 0 \\ 0 & \frac{V^2}{\langle S \rangle} & 0 & \langle S \rangle \end{pmatrix}. \quad (4.7)$$

After the heavy 50-dimensional fields are integrated out and SU(5) gets spontaneously broken, the mass matrices for the doublet and triplet components of the Higgs fields H_5 , H_5' and their corresponding barred fields are given by

$$m_D = \begin{pmatrix} \mu & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} \mu & -\frac{V^4}{\langle S \rangle^3} \\ -\frac{V^4}{\langle S \rangle^3} & \mu' \end{pmatrix}. \quad (4.8)$$

The effective triplet mass is then given by

$$M_T^{\text{eff}} = (m_T^{-1})_{11}^{-1} = \mu - \frac{V^8}{\langle S \rangle^6 \mu'} \approx -\frac{V^8 M_{\text{Pl}}^3}{\langle S \rangle^6 \langle \theta_3 \rangle^4}. \quad (4.9)$$

⁸Alternatively, we checked that a UV-complete generation of these μ and μ' operators via messenger fields would be possible. However, the necessary messenger fields are not included in the model and it turns out that the masses are already generated by Planck-scale suppressed effective operators.

Let us give an explicit example for the scales involved in the model: Because of perturbativity the mass of the 50-dimensional Higgs fields has to be almost at the Planck scale. We therefore assume $\langle S \rangle \sim 10^{-1} M_{\text{Pl}}$. Using the known values of the Yukawa couplings, we can estimate the values of the relevant masses of our model. At the GUT scale with $\tan \beta = 30$ the Yukawa couplings are approximately given by [24]

$$y_d \approx 1.6 \cdot 10^{-4}, \quad y_s \approx 3 \cdot 10^{-3}, \quad \text{and } y_b \approx 0.18. \quad (4.10)$$

In our example model these Yukawa couplings are (up to order one couplings) given by the operators

$$y_d \sim \frac{\langle H_{24} \rangle \langle H'_{24} \rangle \langle \theta_4 \rangle}{\langle S' \rangle^2 \langle S \rangle}, \quad y_s \sim \frac{\langle H'_{24} \rangle \langle \theta_3 \rangle}{\langle S' \rangle^2} \quad \text{and } y_b \sim \frac{\langle H'_{24} \rangle}{\langle S' \rangle}. \quad (4.11)$$

Then we find from eqs. (4.9)–(4.11) for the effective triplet mass, the μ -term, the mass of the additional, heavy Higgs fields and the geometric mean of the physical masses of the colour triplets

$$M_T^{\text{eff}} \approx 2 \cdot 10^{18} \text{ GeV}, \quad \mu \approx \mathcal{O}(100) \text{ GeV}, \quad \mu' \approx \mathcal{O}(10) \text{ TeV}, \quad m_T \approx 2 \cdot 10^{11} \text{ GeV}. \quad (4.12)$$

Although these numbers are only estimates, which neglect order one couplings, they illustrate the model's features: the DTS problem is solved with a large effective triplet mass, therefore the proton decay rate is suppressed and the fermion mass ratios are realistic. The μ -term of the size of the electroweak scale emerges from Planck scale suppressed operators. Note that the heavier, additional Higgs doublet fields are light enough to be in reach of detection by the LHC or its successors. Also the DMPM does not suffer from any dangerous Planck-scale suppressed operators, due to the charge assignment of the singlet field S .

4.2 A model with θ_C from Y_d

We now turn to our second example model, which realises the attractive feature of θ_C emerging dominantly from the down-type quark mixing $\theta_C \approx \theta_{12}^d$. The Yukawa matrices are given by the following structure

$$Y_d = \begin{pmatrix} 0 & y_{d,12} & 0 \\ y_{d,21} & y_s & 0 \\ 0 & 0 & y_b \end{pmatrix}, \quad Y_e = \begin{pmatrix} 0 & 6y_{d,21} & 0 \\ -\frac{1}{2}y_{d,12} & 6y_s & 0 \\ 0 & 0 & -\frac{3}{2}y_b \end{pmatrix}, \quad Y_u = \begin{pmatrix} y_{11} & y_{12} & 0 \\ y_{12} & y_{22} & y_{23} \\ 0 & y_{23} & y_{33} \end{pmatrix}. \quad (4.13)$$

A similar structure [16, 17] has been used to explain the relation $\theta_{13}^{\text{PMNS}} = \theta_C / \sqrt{2}$ via charged lepton corrections, and a right-handed quark unitarity triangle [29]. From the matrices we can see that we want to couple both \mathcal{F}_1 and \mathcal{F}_2 to \mathcal{T}_2 but still distinguish both fields from each other to forbid $(Y_d)_{11}$. This suggests to use at least a \mathbb{Z}_3 symmetry and hence we will use superpotential (b) from section 3.1 where both adjoints are charged.




Figure 7: Supergraphs leading to the effective superpotential W_d of eq. (4.15) when the heavy messenger fields get integrated out in the model presented in subsection 4.2.

The effective superpotentials that lead to the desired Yukawa matrices after integrating out heavy messenger fields and breaking of the GUT gauge group are

$$W_u = \frac{1}{\Lambda^2} H_5 \mathcal{T}_1 \mathcal{T}_1 \theta_5^2 + \frac{1}{\Lambda^2} H_5 \mathcal{T}_1 \mathcal{T}_2 \theta_2^2 + \frac{1}{\Lambda^2} H_5 \mathcal{T}_2 \mathcal{T}_2 \theta_1^2 + \frac{1}{\Lambda} H_5 \mathcal{T}_2 \mathcal{T}_3 \theta_1 + H_5 \mathcal{T}_3 \mathcal{T}_3, \quad (4.14)$$

$$W_d = \frac{1}{\langle S' \rangle} (H'_{24} \mathcal{F}_3)_5 (\bar{H}_5 \mathcal{T}_3)_5 + \frac{\theta_3}{\langle S' \rangle^2} (H_{24} \mathcal{T}_2)_{10} (\bar{H}_5 \mathcal{F}_1)_{\bar{10}} + \frac{\theta_4}{\langle S' \rangle^2} (H'_{24} \mathcal{T}_2)_{10} (\bar{H}_5 \mathcal{F}_2)_{\bar{10}} \\ + \frac{1}{\langle S \rangle^2} (H_{24} \mathcal{F}_2)_{\bar{45}} (\mathcal{T}_1 H_{24} \bar{H}_5)_{45}, \quad (4.15)$$

where again VEVs of singlet fields appear in the denominators by virtue of messenger masses generated by them. Note that in comparison to Y_u of the previous example model of eq. (4.2), the vanishing $(Y_u)_{13}$ element requires to introduce an additional flavon field θ_5 . The supergraphs that generate these effective operators are shown in Figure 7 and 8. A complete list of all fields including their charges and representations is given in Tables 6, 7, 8 and 9.

In a small angle approximation the mixing angles and Yukawa couplings are given by

$$y_d \approx \frac{y_{d,12} y_{d,21}}{y_s}, \quad \theta_C \approx \theta_{12}^d \approx \frac{y_{d,12}}{y_s}, \\ y_u \approx y_{11} - \frac{y_{12}^2}{y_{22}}, \quad y_c \approx y_{22}, \quad y_t \approx y_{33}, \quad \theta_{23} \approx \frac{y_{23}}{y_{33}}, \quad \theta_{13} \approx \frac{y_{12}}{y_{22}} \theta_{23}, \quad (4.16)$$

with y_s and y_b given as parameters of Y_d in eq. (4.13). Thus the Yukawa matrices can fit the experimental values without tension.

We continue now by discussing additional details concerning the use of the DMPM in this model. Like in the previous model, Planck suppressed operators generate mass terms

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_3	\mathbb{Z}_4	\mathbb{Z}_5	\mathbb{Z}_6	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_2
H_5	5
\bar{H}_5	$\bar{5}$.	2	.	1	4	6	6	.
\mathcal{T}_1	10	6	5	1
\mathcal{T}_2	10	1	.	1
\mathcal{T}_3	10	1
\mathcal{F}_1	$\bar{5}$.	.	.	3	2	5	6	1
\mathcal{F}_2	$\bar{5}$.	2	.	3	.	2	3	1
\mathcal{F}_3	$\bar{5}$.	1	3	4	.	1	1	1

Table 6: SU(5) representations and charges under discrete shaping symmetries of the MSSM fields and colour triplets of the model presented in subsection 4.2. A dot denotes charge zero.

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_3	\mathbb{Z}_4	\mathbb{Z}_5	\mathbb{Z}_6	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_2
H_{24}	24	4	.	.	.
H'_{24}	24	2	.	.	.
S	1	1	2	.	2
S'	1	.	.	3
θ_1	1	6	.	.
θ_2	1	1	.
θ_3	1	.	1	2	1	2	2	2	.
θ_4	1	.	2	2	1	.	5	5	.
θ_5	1	3	1	2	.

Table 7: SU(5) representations and charges under discrete shaping symmetries of the superfields obtaining VEVs at around the GUT scale of the model presented in subsection 4.2. A dot denotes charge zero.

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_3	\mathbb{Z}_4	\mathbb{Z}_5	\mathbb{Z}_6	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_2
H'_5	5	1	1	.	.	4	1	1	.
\bar{H}'_5	$\bar{5}$	1	.	.	1	2	.	.	.
X_{45}	45	.	1	.	4	4	1	1	.
\bar{X}_{45}	$\bar{45}$	1	.	.	4	2	6	6	.
Y_{45}	45	.	2	.	3	2	1	1	.
\bar{Y}_{45}	$\bar{45}$	1	2	.	.	4	6	6	.
Z_{50}	50	1	.	.	1	.	1	1	.
\bar{Z}_{50}	$\bar{50}$.	1	.	2	.	6	6	.
X'_{45}	45	1	.	.	4
\bar{X}'_{45}	$\bar{45}$.	1	.	4
Y'_{45}	45	1	1	.	3	4	.	.	.
\bar{Y}'_{45}	$\bar{45}$	2	.	.	.
Z'_{50}	50	.	2	.	1	2	.	.	.
\bar{Z}'_{50}	$\bar{50}$	1	2	.	2	4	.	.	.

Table 8: SU(5) representations and charges under discrete shaping symmetries of the fields in the DPM sector of the model presented in subsection 4.2. A dot denotes charge zero.




Figure 8: Supergraphs leading to the effective superpotential W_u of eq. (4.14) when the heavy messenger fields get integrated out.

	SU(5)	\mathbb{Z}_2	\mathbb{Z}_3	\mathbb{Z}_4	\mathbb{Z}_5	\mathbb{Z}_6	\mathbb{Z}_7	\mathbb{Z}_7	\mathbb{Z}_2
Z_5	5	.	2	1	1	4	6	6	1
\bar{Z}_5	$\bar{\mathbf{5}}$.	1	.	4	2	1	1	1
$Z_{10,1}$	10	.	1	.	1	.	3	2	1
$\bar{Z}_{10,1}$	$\bar{\mathbf{10}}$.	2	1	4	.	4	5	1
$Z_{10,2}$	10	.	.	1	.	4	1	.	1
$\bar{Z}_{10,2}$	$\bar{\mathbf{10}}$	2	6	.	1
$Z_{10,3}$	10	.	2	.	1	2	6	5	1
$\bar{Z}_{10,3}$	$\bar{\mathbf{10}}$.	1	1	4	4	1	2	1
$Z_{10,4}$	10	.	.	1	.	2	1	.	1
$\bar{Z}_{10,4}$	$\bar{\mathbf{10}}$	4	6	.	1
$Z_{45,1}$	45	.	1	.	2	2	5	4	1
$\bar{Z}_{45,1}$	$\bar{\mathbf{45}}$	1	.	.	1	4	2	3	1
$Z_{10,5}$	10	6	.	1
$Z_{10,6}$	10	1	2	1
Z_1	1	2	.	.
Z_2	1	5	.
Z_3	1	5	3	.

Table 9: SU(5) representations and charges under discrete shaping symmetries of the flavon and flavour messenger fields of the model presented in subsection 4.2. Note that the messengers $Z_{5,1}\bar{Z}_{5,1}$, $Z_{10,1}\bar{Z}_{10,1}$, $Z_{10,2}\bar{Z}_{10,2}$, $Z_{10,3}\bar{Z}_{10,3}$, $Z_{10,4}\bar{Z}_{10,4}$ and $Z_{45,1}\bar{Z}_{45,1}$ have no direct mass term, but get their masses through VEVs of S and S' . The other messenger fields have direct mass terms, so their corresponding barred field is not shown in the table. A dot denotes charge zero.

for the five-dimensional Higgs representations

$$W_5^{\text{eff}} = \mu H_5 \bar{H}_5 + \mu' H'_5 \bar{H}'_5, \quad (4.17)$$

where $\mu = \langle \theta_3 \rangle^4 / M_{\text{Pl}}^3$ and $\mu' = \langle \theta_4 \rangle^4 / M_{\text{Pl}}^3$. Note that although they appear at the same order, a modest hierarchy $\langle \theta_3 \rangle < \langle \theta_4 \rangle$, as we have in our model, will sufficiently split their masses. Furthermore, after integrating out the 45-dimensional messengers of the DMPM, the mass matrices for doublet and triplet components of the 5- and 50-dimensional superfields are given by

$$m_D = \begin{pmatrix} \mu & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} \mu & 0 & 0 & \frac{V^2}{\langle S \rangle} \\ 0 & \mu' & \frac{V^2}{\langle S \rangle} & 0 \\ \frac{V^2}{\langle S \rangle} & 0 & \langle S \rangle & 0 \\ 0 & \frac{V^2}{\langle S \rangle} & 0 & \langle S \rangle \end{pmatrix}, \quad (4.18)$$

which after the heavy 50-dimensional fields get integrated out become

$$m_D = \begin{pmatrix} \mu & 0 \\ 0 & \mu' \end{pmatrix}, \quad m_T = \begin{pmatrix} \mu & -\frac{V^4}{\langle S \rangle^3} \\ -\frac{V^4}{\langle S \rangle^3} & \mu' \end{pmatrix}. \quad (4.19)$$

This leads to an effective triplet mass of

$$M_T^{\text{eff}} = (m_T^{-1})_{11}^{-1} = \mu - \frac{V^8}{\langle S \rangle^6 \mu'} \approx -\frac{V^8 M_{\text{Pl}}^3}{\langle S \rangle^6 \langle \theta_4 \rangle^4}. \quad (4.20)$$

As for the first example model we can estimate the values of the relevant masses in an explicit example from the known Yukawa couplings. In a small angle approximation the down-type Yukawa couplings are given by

$$y_d \sim \frac{\langle H_{24} \rangle^2 \langle H_{24} \rangle \langle \theta_3 \rangle}{\langle S \rangle^2 \langle S' \rangle^2} \frac{1}{y_s} \approx 1.6 \cdot 10^{-4} \quad \text{with} \quad y_s \sim \frac{\langle H'_{24} \rangle \langle \theta_4 \rangle}{\langle S' \rangle^2} \approx 3 \cdot 10^{-3} \quad (4.21)$$

and

$$y_b \sim \frac{\langle H'_{24} \rangle}{\langle S' \rangle} \approx 0.18, \quad (4.22)$$

where the numerical values for the Yukawa couplings taken from [24] are valid for $\tan \beta = 30$ and order one coefficients have been neglected. For the effective triplet mass, the μ -term, the mass of the additional, heavier Higgs doublets and the geometrical mean of the physical masses of the colour triplets the following estimates emerge

$$M_T^{\text{eff}} \approx 2 \cdot 10^{18} \text{ GeV}, \quad \mu \approx \mathcal{O}(300) \text{ GeV}, \quad \mu' \approx \mathcal{O}(10) \text{ TeV}, \quad m_T \approx 2 \cdot 10^{11} \text{ GeV}. \quad (4.23)$$

Thus, also in this model, proton decay can be suppressed by a large effective triplet mass. DTS is achieved through the DMPM and the light doublet gets its mass from an Planck-suppressed operator. The Yukawa sector of the model features viable fermion masses and quark mixing angles.

5 Proton Decay

We have split our discussion of proton decay into two parts. In the first part we comment on proton decay induced by dimension five operators in the superpotential. These operators are usually considered to be more dangerous for the validity of any SUSY GUT model. We will argue why we are more predictive than ordinary models and still should be able to evade current experimental bounds. In the second part we will comment on dimension six proton decay operators. These appear also in non-SUSY GUTs, but are considered not as dangerous in SUSY GUTs due to the usually higher unification scale [6].

5.1 Proton Decay from Dimension Five Operators

We will adopt for this section a notation similar to the SLHA convention [30], for more details see appendix A. The superpotential describing the couplings of the matter fields to the heavy colour triplet reads

$$W_T = \epsilon_{\alpha\beta} \left(-\frac{1}{2} (Y_{qq})_{ij} \epsilon_{abc} T^a Q_i^{\alpha b} Q_j^{\beta c} + (Y_{ql})_{ij} \bar{T}^a Q_i^{\alpha a} L_j^{\beta} \right) \\ + (Y_{ue})_{ij} T^a \bar{U}_i^a \bar{E}_j - (Y_{ud})_{ij} \epsilon_{abc} \bar{T}^a \bar{U}_i^b \bar{D}_j^c + M_T T^a \bar{T}^a . \quad (5.1)$$

In many SU(5) models at least for the first two generations it was assumed that the Yukawa couplings are generated or significantly corrected by some set of higher-dimensional operators, see for instance, [23, 31, 32]. But, usually, there is no control over which operator is the dominant one and hence it is not possible to calculate how strong exactly the heavy triplets couple to the MSSM fields. In our setup this is not the case. For every entry of the Yukawa matrix we have specified the operator with only very small corrections, if any. Therefore we know how strong the MSSM fields couple to the heavy triplets if we know the MSSM Yukawa couplings at the GUT scale. To be more precise the new Yukawa couplings are related to the MSSM ones only via CG coefficients which are fixed by the gauge structure of the underlying operator. We find for our first example model

$$Y_d = \text{Diag}(y_d, y_s, y_b) , \quad Y_e = \text{Diag}\left(-\frac{1}{2}y_d, 6y_s, -\frac{3}{2}y_b\right) , \quad (5.2)$$

$$Y_{ql} = \text{Diag}\left(y_d, y_s, -\frac{3}{2}y_b\right) , \quad Y_{ud} = \text{Diag}\left(\frac{2}{3}y_d, -4y_s, y_b\right) , \quad (5.3)$$

$$Y_{qq} = Y_u , \quad Y_{ue} = Y_u , \quad (5.4)$$

where the structure of Y_u can be read off from eq. (4.2). For the second model we find

$$Y_d = \begin{pmatrix} 0 & y_{d,12} & 0 \\ y_{d,21} & y_s & 0 \\ 0 & 0 & y_b \end{pmatrix} , \quad Y_e^T = \begin{pmatrix} 0 & -\frac{1}{2}y_{d,12} & 0 \\ 6y_{d,21} & 6y_s & 0 \\ 0 & 0 & -\frac{3}{2}y_b \end{pmatrix} , \quad (5.5)$$

$$Y_{ql} = \begin{pmatrix} 0 & y_{d,12} & 0 \\ y_{d,21} & y_s & 0 \\ 0 & 0 & -\frac{3}{2}y_b \end{pmatrix} , \quad Y_{ud} = \begin{pmatrix} 0 & \frac{2}{3}y_{d,12} & 0 \\ -4y_{d,21} & -4y_s & 0 \\ 0 & 0 & y_b \end{pmatrix} , \quad (5.6)$$

$$Y_{qq} = Y_u, \quad Y_{ue} = Y_u, \quad (5.7)$$

where the structure of Y_u can be read off from eq. (4.13).⁹

The dimension five operators which violate baryon number after integrating out the triplets read

$$W_{\mathcal{B}} = \frac{1}{M_T^{\text{eff}}} \left[\frac{1}{2} Y_{qq}^{ij} Y_{ql}^{mn} Q_i Q_j Q_m L_n + Y_{ue}^{ij} Y_{ud}^{mn} \bar{U}_i \bar{E}_j \bar{U}_m \bar{D}_n \right], \quad (5.8)$$

where we have suppressed ϵ -tensors. The first operator is called the $LLLL$ operator and the second one the $RRRR$ operator. To make definite predictions for the proton decay rate one would have to take the RGE evolution of these operators to low energies into account and dress the operators with a closed loop including MSSM particles to calculate the decay rate of the proton, see e.g. [34] and references therein. This goes clearly beyond the scope of this paper. Before we give some more qualitative statements about what we expect for the proton decay rate in comparison to other models we want to argue first that the operators in eq. (5.8) together with M_T^{eff} from eq. (2.13) give the dominant contribution.

Inside the additional higher dimensional representations used in the DMPM there are additional colour triplets which could in principle give the same operators like in eq. (5.8) but with a weaker suppression. We have explicitly checked that the DMPM by itself is safe: the additional colour triplets from 5-, 45- and 50-dimensional representations can only mediate operators that are suppressed compared to the leading contribution by one or more powers of H_{24}/M_{Pl} and H'_{24}/M_{Pl} .

We have also checked for each of our models that no other Yukawa matrix entries (through components in the messengers) lead to a lower M_T^{eff} . This is, of course, model dependent and has to be checked for any specific model.

We now turn to a qualitative discussion of dimension five proton decay in the considered class of models. Firstly, we want to point out that especially in the first model one could expect some decay modes of the proton to be suppressed because Y_{ql} and Y_{ud} are diagonal in flavour space. Therefore, decays which need a flavour transition in these matrices would be suppressed. This reinforces the statement that our setup is more predictive than conventional ones, due to the better control over the flavour structure of all the Yukawa matrices governing proton decay.

Secondly, introducing a second adjoint in a renormalizable way gives us enough freedom to enhance the effective triplet mass M_T^{eff} to comfortable levels, cf. section 3. In the case of superpotential (b), we get a mass of roughly $5 \cdot 10^{16}$ GeV for a SUSY scale of 1 TeV which increases with increasing M_{SUSY} , see section 3.1.2. For superpotential (a) the situation is more involved and depends on the specific GUT scale, and the effective triplet mass can be easily above 10^{18} GeV.

To conclude the discussion of the dimension five operators mediating proton decay we also want to stress that in our setup we need only a moderate value of $\tan \beta \approx 25$. As it was

⁹GUT textures for proton decay, without fully constructed models, have been considered, for example, in [32, 33].

pointed out by Lucas and Raby [35] especially the contribution from the $RRRR$ operator is enhanced by $\tan^2 \beta$ for large $\tan \beta$ which poses a challenge for many GUT scenarios which rely on $\tan \beta \approx 50$. Our setup with $y_\tau/y_b = \frac{3}{2}$ and $\tan \beta \approx 25$ has therefore a suppression of proton decay via the $RRRR$ operator by a factor of four compared to these models.

In summary, together with the large effective triplet mass in the double missing partner mechanism we hence expect the proton decay rate to be sufficiently small but possibly in the reach of the next generation of proton decay experiments.

5.2 Proton Decay from Dimension Six Operators

For completeness, we will now briefly discuss as well proton decay from dimension six operators. There is no substantial difference in our models compared to other SUSY SU(5) models since the gauge structure is exactly the same.

To make the discussion a little bit more analogous to the previous discussion we define an effective GUT scale

$$M_{\text{GUT}}^{\text{eff}} = \frac{M_{\text{GUT}}}{\sqrt{\alpha_u}}, \quad (5.9)$$

where α_u is the unified gauge coupling at the GUT scale and $M_{\text{GUT}} \equiv M_V$ is the mass of the leptoquark vector bosons, cf. the discussion in section 3. Just integrating out the heavy leptoquarks in the tree-level diagrams governing proton decay, and from dimensional analysis, we can estimate the lifetime of the proton to be

$$\Gamma_p \approx \alpha_u^2 \frac{m_p^5}{M_V^4} = \frac{m_p^5}{(M_{\text{GUT}}^{\text{eff}})^4}, \quad (5.10)$$

where m_p is the proton mass and we have neglected RGE effects and order one coefficients from nuclear matrix elements and such. The most stringent bound on the proton lifetime is $\tau(p \rightarrow \pi^0 e^+) > 8.2 \cdot 10^{33}$ years [10] which yields an effective GUT scale of about $M_{\text{GUT}}^{\text{eff}} \gtrsim 2 \cdot 10^{16} \text{GeV}$. In section 3 we have seen that the GUT scale can be easily above 10^{16} GeV and $\sqrt{\alpha_u} \approx 1/5$ such that the proton decay rate from dimension six operators is sufficiently small but possibly in the reach of the next generation of proton decay experiments.

6 Summary and Conclusions

In this work we have discussed how the double missing partner mechanism solution to the doublet-triplet splitting problem in four-dimensional supersymmetric SU(5) Grand Unified Theories can be combined with predictive models featuring novel predictions for the quark-lepton Yukawa coupling ratios at the GUT scale.

We have argued that towards this goal a second SU(5) breaking Higgs field in the adjoint representation is very useful. We systematically discussed all possible renormalizable superpotentials with two adjoint Higgs fields, also calculating the corresponding constraints on the GUT scale and effective triplet mass from a two-loop gauge coupling unification analysis. We found that the effective mass of the colour triplet can easily be raised enough

to avoid problems with proton decay (more than feasible with standard non-renormalizable Higgs potentials with only one adjoint GUT Higgs field).

We have constructed two explicit flavour models with different predictions for the GUT scale Yukawa sector. A set of shaping symmetries and a renormalizable messenger sector for the models is presented, which guarantees that only the desired effective GUT operators are generated when the heavy degrees of freedom are integrated out. In addition, we also include all possible effective Planck scale suppressed operators consistent with our symmetries, and make sure that they do not spoil our results. The models stay perturbative until close to the Planck-scale, such that our predictions do not suffer from large uncertainties due to these Planck-scale suppressed operators. They serve as existence proofs that predictive models for the GUT scale quark-lepton mass relations can be combined successfully with the DMPM solution for solving the DTS problem.

We also provide several useful appendices for GUT flavour model building: For instance, one appendix contains the Clebsch-Gordan coefficients for the couplings of the colour triplets, which are required for calculating the rates for proton decay induced by their exchange. We provide detailed tables with the Clebsch-Gordan coefficients for the possible dimension five and six GUT Yukawa operators. We also discuss there how one can use GUT Higgs potentials for flavour model building, where a (discrete) R-symmetry is broken spontaneously around the GUT scale. R-symmetries are a helpful ingredient of many flavour models, especially when they include non-Abelian family symmetries.

In summary, we have demonstrated that four-dimensional supersymmetric SU(5) GUTs with successful doublet-triplet splitting can be combined with predictive models featuring promising predictions for the quark-lepton Yukawa coupling ratios at the GUT scale. We have provided the tools for the construction of even more ambitious GUT models of flavour with additional non-Abelian family symmetries, as well as towards the calculation of the predictions for the rates of the various nucleon decay channels in such models by systematically providing the required Clebsch-Gordan coefficients.

Acknowledgements

This work is supported by the Swiss National Science Foundation. We thank Borut Bajc and David Emmanuel-Costa for useful discussions.

Appendix

A Yukawa Coupling Ratios including Colour Triplets

The MSSM superpotential is given by

$$W = \epsilon_{\alpha\beta} \left((Y_e)^{ij} H_d^\alpha L_i^\beta \bar{E}_j + (Y_d)^{ij} H_d^\alpha Q_i^{\beta a} \bar{D}_j^a + (Y_u)^{ij} H_d^\beta Q_i^{\alpha a} \bar{U}_j^a + \mu H_u^\alpha H_d^\beta \right), \quad (\text{A.1})$$

where i, j are generation indices, $\epsilon_{\alpha\beta}$ the Levi-Civita tensor ($\epsilon_{12} = 1$), α, β are SU(2) indices and a, b and c are SU(3) indices¹⁰. Adding a pair of colour triplets T and \bar{T} , we get the additional terms

$$W_T = \epsilon_{\alpha\beta} \left(-\frac{1}{2} (Y_{qq})_{ij} \epsilon_{abc} T^a Q_i^{\alpha b} Q_j^{\beta c} + (Y_{ql})_{ij} \bar{T}^a Q_i^{\alpha a} L_j^{\beta} \right) + (Y_{ue})_{ij} T^a \bar{U}_i^a \bar{E}_j - (Y_{ud})_{ij} \epsilon_{abc} \bar{T}^a \bar{U}_i^b \bar{D}_j^c + M_T T^a \bar{T}^a, \quad (\text{A.2})$$

where ϵ_{abc} is the three indices Levi-Civita tensor (with $\epsilon_{123} = 1$).

Extending the SM gauge group to SU(5), we embed the MSSM superfields in a 5-plet H_5 , $\bar{5}$ -plets \bar{H}_5 and \mathcal{F}_i , and 10-plets \mathcal{T}_i as in

$$H_5 = (T^r \quad T^g \quad T^b \quad H_u^0 \quad -H_u^+)^T, \quad (\text{A.3})$$

$$\bar{H}_5 = (\bar{T}^r \quad \bar{T}^g \quad \bar{T}^b \quad H_d^- \quad -H_d^0), \quad (\text{A.4})$$

$$\mathcal{F}_i = (\bar{D}_i^r \quad \bar{D}_i^g \quad \bar{D}_i^b \quad E_i \quad -\nu_i), \quad (\text{A.5})$$

$$\mathcal{T}_i = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -\bar{U}_i^b & \bar{U}_i^g & -U_i^r & -D_i^r \\ \bar{U}_i^b & 0 & -\bar{U}_i^r & -U_i^g & -D_i^g \\ -\bar{U}_i^g & \bar{U}_i^r & 0 & -U_i^b & -D_i^b \\ U_i^r & U_i^g & U_i^b & 0 & -\bar{E}_i \\ D_i^r & D_i^g & D_i^b & \bar{E}_i & 0 \end{pmatrix}, \quad (\text{A.6})$$

where r, g, b are the SU(3) colours and U, D and ν, E are the components of SU(2)-doublets Q and L .¹¹ We can write down the renormalizable superpotential terms

$$W = (Y_{TF})_{ij} \mathcal{T}_i^{ab} (\mathcal{F}_j)_a (\bar{H}_5)_b + \frac{1}{2} (Y_{TT})_{ij} \epsilon_{abcde} \mathcal{T}_i^{ab} \mathcal{T}_j^{cd} H_5^e + \mu_5 H_5^a (\bar{H}_5)_a, \quad (\text{A.7})$$

where now a, b, c, d, e are SU(5)-indices and ϵ_{abcde} is the respective Levi-Civita tensor. From the embedding of the MSSM fields, one obtains the minimal SU(5) GUT scale relations

$$\mu = M_T = \mu_5, \quad (\text{A.8})$$

$$Y_d = Y_e^T = Y_{ql} = Y_{ud} = \frac{1}{\sqrt{2}} Y_{TF}, \quad (\text{A.9})$$

$$Y_u = Y_u^T = Y_{qq} = Y_{ue} = 2 Y_{TT}. \quad (\text{A.10})$$

The relation $Y_d = Y_e^T$ is highly disfavoured as was already discussed in section 2.4. The conventional approach is to add a 45-dimensional Higgs representation which generates a relative factor of -3 between the Yukawa couplings of the charged leptons and down-type quarks [3]. In this work we instead focus on an approach where the ratios between Yukawa couplings are fixed by the CG coefficients of higher-dimensional operators where in addition an adjoint Higgs representation of SU(5) is added [11, 21]. This approach was briefly reviewed in section 2.4 and for a thorough discussion we refer the interested reader to the original papers, see also [36].

¹⁰This definition coincides with the definitions of [30].

¹¹Likewise $H_d = (H_d^0 \quad H_d^-)^T$ and $H_u = (H_u^+ \quad H_u^0)^T$.




Figure 9: Supergraphs generating Yukawa couplings upon integrating out messengers fields in representation R, \bar{R} , etc.

In this appendix we will extend the previous discussions to include also the relative CG coefficients to the triplets. In [11, 21] only Y_d , Y_e and Y_u were discussed while here we will also discuss in detail the implications of this approach for Y_{qt} , Y_{ud} , Y_{qq} , Y_{ue} . The list of the resulting ratios for dimension 4 and 5 operators with Higgs fields in a 5- and 45-dimensional representation can be found in Tabs. 10 and 11, where the labels for the representations is defined by figure 9. The corresponding results for dimension six operators are given in Tabs. 12 - 15. The ones used in the example models are marked with a “ \rightarrow ” in the tables.

There are a few comments in order. First, note that several topologies involving a 45-dimensional messenger field exhibit a free parameter, simply because the tensor product $45 \otimes 24$ contains two 45-dimensional representations. Hence, there are two operators possibly giving two different ratios so that any ratio is possible depending on the coefficients of the two operators. For these cases we write x in the tables.

We want to mention as well that, unlike at the renormalizable level, the up-type quark Yukawa and related matrices do not have to be symmetric or antisymmetric. Consider, for example, the operator $(H_{24}\mathcal{T}_1)_{10}(H_5\mathcal{T}_2)_{\bar{10}}$. Due to the symmetries and messenger content the operator $(H_{24}\mathcal{T}_2)_{10}(H_5\mathcal{T}_1)_{\bar{10}}$ could be forbidden. In this case we find $(Y_u)_{12}/(Y_u)_{21} = -4$. Hence we have adopted the following notation for the ratios in the tables for the Yukawa couplings related to Y_u

$$(Y_u)_{ij} : (Y_u)_{ji} : (Y_{qq})_{ij} : (Y_{ue})_{ij} : (Y_{ue})_{ji} = a : b : c : d : e , \quad (\text{A.11})$$

which reduces for the diagonal entries of the Yukawa matrices to

$$(Y_u)_{ii} : (Y_{qq})_{ii} : (Y_{ue})_{ii} = (a + b) : c : (d + e) . \quad (\text{A.12})$$

The ratios related to Y_d do not have this extra complication since none of them could be expected to be symmetric or anti-symmetric in the first place.

If the considered model contains Higgs fields in 5- and 45-dimensional representations¹², there are two Higgs doublet pairs in the spectrum and care should be taken that unification is still possible. One solution is mixing both and making one linear combination heavy

¹²One could also imagine using 45-dimensional Higgs fields exclusively. However, this severely exacerbates the doublet-triplet splitting problem as now one has to split the doublets from even more component fields that can generate proton decay operators.

	AB	CD	R	$(Y_d)_{ij} : (Y_e)_{ji} : (Y_{ql})_{ij} : (Y_{ud})_{ij}$
		$\mathcal{F}_j \mathcal{T}_i \bar{H}_5$	—	1 : 1 : 1 : 1
→	$H_{24} \mathcal{T}_i$	$\mathcal{F}_j \bar{H}_5$	10	1 : 6 : 1 : -4
	$H_{24} \mathcal{T}_i$	$\mathcal{F}_j \bar{H}_5$	15	1 : 0 : -1 : 0
	$H_{24} \bar{H}_5$	$\mathcal{F}_j \mathcal{T}_i$	$\bar{5}$	1 : 1 : $-\frac{2}{3}$: $-\frac{2}{3}$
	$H_{24} \bar{H}_5$	$\mathcal{F}_j \mathcal{T}_i$	$\bar{45}$	1 : -3 : -2 : 2
→	$H_{24} \mathcal{F}_j$	$\mathcal{T}_i \bar{H}_5$	$\bar{5}$	1 : $-\frac{3}{2}$: $-\frac{3}{2}$: 1
	$H_{24} \mathcal{F}_j$	$\mathcal{T}_i \bar{H}_5$	$\bar{45}$	1 : $\frac{3}{2}$: $-\frac{1}{2}$: -1
		$\mathcal{F}_j \mathcal{T}_i \bar{H}_{45}$	—	1 : -3 : $\sqrt{3}$: $-\sqrt{3}$
	$H_{24} \mathcal{T}_i$	$\mathcal{F}_j \bar{H}_{45}$	10	1 : -18 : $\sqrt{3}$: $4\sqrt{3}$
	$H_{24} \mathcal{T}_i$	$\mathcal{F}_j \bar{H}_{45}$	40	1 : 0 : $-\frac{\sqrt{3}}{2}$: $-\frac{\sqrt{3}}{2}$
	$H_{24} \mathcal{T}_i$	$\mathcal{F}_j \bar{H}_{45}$	175	1 : $\frac{36}{23}$: $-\frac{19\sqrt{3}}{23}$: $-\frac{16\sqrt{3}}{23}$
	$H_{24} \bar{H}_{45}$	$\mathcal{F}_j \mathcal{T}_i$	$\bar{5}$	1 : 1 : $-\frac{2}{\sqrt{3}}$: $-\frac{2}{\sqrt{3}}$
	$H_{24} \bar{H}_{45}$	$\mathcal{F}_j \mathcal{T}_i$	$\bar{45}$	1 : -3 : x : $-x$
	$H_{24} \mathcal{F}_j$	$\mathcal{T}_i \bar{H}_{45}$	$\bar{5}$	1 : $\frac{9}{2}$: $-\frac{3\sqrt{3}}{2}$: $-\sqrt{3}$
	$H_{24} \mathcal{F}_j$	$\mathcal{T}_i \bar{H}_{45}$	$\bar{45}$	1 : $-\frac{1}{2}$: $-\frac{\sqrt{3}}{2}$: $-\frac{1}{\sqrt{3}}$
	$H_{24} \mathcal{F}_j$	$\mathcal{T}_i \bar{H}_{45}$	$\bar{70}$	1 : $\frac{9}{4}$: $-\frac{3\sqrt{3}}{4}$: $-\sqrt{3}$

Table 10: Y_{TF} -like CG ratios for the dimension 4 operator and effective dimension 5 operators $W \supset (AB)_R(CD)_{\bar{R}}$ (involving 5- and 45-dimensional Higgs fields) corresponding to the the left diagram in figure 9. Note that one combination has a free parameter x due to the ambiguity of the index contraction. See main text for more details.

	AB	CD	R	$(Y_u)_{ij} : (Y_u)_{ji} : (Y_{qq})_{ij} : (Y_{ue})_{ij} : (Y_{ue})_{ji}$
\rightarrow	$\mathcal{T}_i \mathcal{T}_j H_5$	—	—	1 : 1 : 1 : 1 : 1
	$H_{24} H_5$	$\mathcal{T}_i \mathcal{T}_j$	5	1 : 1 : $-\frac{2}{3}$: $-\frac{2}{3}$: $-\frac{2}{3}$
	$H_{24} H_5$	$\mathcal{T}_i \mathcal{T}_j$	45	1 : -1 : 0 : -2 : 2
	$H_{24} \mathcal{T}_i$	$\mathcal{T}_j H_5$	10	1 : -4 : 1 : -4 : 6
	$H_{24} \mathcal{T}_i$	$\mathcal{T}_j H_5$	40	1 : $\frac{1}{2}$: $-\frac{1}{2}$: -1 : 0
	$\mathcal{T}_i \mathcal{T}_j H_{45}$	—	—	1 : -1 : 0 : $\sqrt{3}$: $-\sqrt{3}$
	$H_{24} \mathcal{T}_i$	$\mathcal{T}_j H_{45}$	10	1 : 4 : 0 : $-4\sqrt{3}$: $-6\sqrt{3}$
	$H_{24} \mathcal{T}_i$	$\mathcal{T}_j H_{45}$	15	1 : 0 : $-\frac{\sqrt{3}}{2}$: 0 : 0
	$H_{24} \mathcal{T}_i$	$\mathcal{T}_j H_{45}$	40	1 : $-\frac{7}{2}$: $\frac{3\sqrt{3}}{2}$: $-\sqrt{3}$: 0
	$H_{24} \mathcal{T}_i$	$\mathcal{T}_j H_{45}$	175	1 : $\frac{16}{19}$: $-\frac{21\sqrt{3}}{38}$: $-\frac{16\sqrt{3}}{19}$: $-\frac{12\sqrt{3}}{19}$
	$H_{24} H_{45}$	$\mathcal{T}_i \mathcal{T}_j$	5	1 : 1 : $-\frac{2}{\sqrt{3}}$: $-\frac{2}{\sqrt{3}}$: $-\frac{2}{\sqrt{3}}$
	$H_{24} H_{45}$	$\mathcal{T}_i \mathcal{T}_j$	45	1 : -1 : 0 : x : $-x$
	$H_{24} H_{45}$	$\mathcal{T}_i \mathcal{T}_j$	50	0 : 0 : 1 : -2 : -2

Table 11: Y_{TT} -like CG ratios for the dimension 4 operator and effective dimension 5 operators $W \supset (AB)_R (CD)_R$ (involving 5- and 45-dimensional Higgs fields) corresponding to the the left diagram in figure 9. Note that one combination has a free parameter x due to the ambiguity of the index contraction. See main text for more details.

AB	C	DE	R_1, R_2	$(Y_d)_{ij} : (Y_e)_{ji} : (Y_{ql})_{ij} : (Y_{ud})_{ij}$
$\mathcal{T}_i \bar{H}_5$	\mathcal{F}_j	$H_{24} H_{24}$	5, 1	1 : 1 : 1 : 1
$\mathcal{T}_i \bar{H}_5$	\mathcal{F}_j	$H_{24} H_{24}$	5, 24	1 : $-\frac{3}{2}$: $-\frac{3}{2}$: 1
$\mathcal{T}_i \bar{H}_5$	\mathcal{F}_j	$H_{24} H_{24}$	45, 24	1 : $\frac{3}{2}$: $-\frac{1}{2}$: -1
$\mathcal{T}_i \bar{H}_5$	\mathcal{F}_j	$H_{24} H_{24}$	45, 75	1 : -3 : 1 : -1
$H_{24} \bar{H}_5$	H_{24}	$\mathcal{F}_j \mathcal{T}_i$	$\bar{5}, 5$	1 : 1 : $\frac{4}{9}$: $\frac{4}{9}$
$H_{24} \bar{H}_5$	H_{24}	$\mathcal{F}_j \mathcal{T}_i$	$\bar{5}, 45$	1 : -3 : $\frac{4}{3}$: $-\frac{4}{3}$
$H_{24} \bar{H}_5$	H_{24}	$\mathcal{F}_j \mathcal{T}_i$	$\overline{45}, 5$	1 : 1 : $\frac{4}{3}$: $\frac{4}{3}$
$H_{24} \bar{H}_5$	H_{24}	$\mathcal{F}_j \mathcal{T}_i$	$\overline{45}, 45$	1 : -3 : x : $-x$
$H_{24} \bar{H}_5$	H_{24}	$\mathcal{F}_j \mathcal{T}_i$	$\overline{70}, 5$	1 : 1 : $\frac{8}{9}$: $\frac{8}{9}$
$H_{24} \bar{H}_5$	H_{24}	$\mathcal{F}_j \mathcal{T}_i$	$\overline{70}, 45$	1 : -3 : $\frac{8}{3}$: $-\frac{8}{3}$
$\mathcal{F}_j \mathcal{T}_i$	\bar{H}_5	$H_{24} H_{24}$	5, 1	1 : 1 : 1 : 1
$\mathcal{F}_j \mathcal{T}_i$	\bar{H}_5	$H_{24} H_{24}$	5, 24	1 : 1 : $-\frac{2}{3}$: $-\frac{2}{3}$
$\mathcal{F}_j \mathcal{T}_i$	\bar{H}_5	$H_{24} H_{24}$	45, 24	1 : -3 : -2 : 2
$\mathcal{F}_j \mathcal{T}_i$	\bar{H}_5	$H_{24} H_{24}$	45, 75	1 : -3 : 1 : -1
$H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\bar{5}, \bar{5}$	1 : $-\frac{3}{2}$: 1 : $-\frac{2}{3}$
$H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\bar{5}, \overline{45}$	1 : $\frac{3}{2}$: $\frac{1}{3}$: $\frac{2}{3}$
$H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\overline{45}, \bar{5}$	1 : $\frac{9}{2}$: 3 : 2
$\rightarrow H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\overline{45}, \overline{45}$	1 : $-\frac{1}{2}$: 1 : $\frac{2}{3}$
$H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\overline{45}, \overline{70}$	1 : $\frac{9}{4}$: $\frac{3}{2}$: 2
$H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\overline{70}, \overline{45}$	1 : $\frac{3}{2}$: $\frac{2}{3}$: $\frac{4}{3}$
$H_{24} \bar{H}_5$	\mathcal{T}_i	$H_{24} \mathcal{F}_j$	$\overline{70}, \overline{70}$	1 : $\frac{3}{4}$: 1 : $\frac{2}{3}$
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{10}, 10$	1 : 36 : 1 : 16
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{10}, 15$	1 : 0 : 1 : 0
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{10}, 40$	1 : 0 : 1 : 1
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{10}, 175$	1 : $\frac{72}{61}$: 1 : $\frac{64}{61}$
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{15}, 10$	1 : 0 : -1 : 0
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{15}, 15$	1 : 0 : -1 : 0
$\mathcal{F}_j \bar{H}_5$	H_{24}	$H_{24} \mathcal{T}_i$	$\overline{15}, 175$	1 : 0 : -1 : 0

Table 12: Y_{TF} -like CG ratios for the effective dimension 6 operators $W \supset (AB)_{R_1} C(DE)_{R_2}$ corresponding to the the right diagram in figure 9. Note one combination has a free parameter x . See main text for more details.

AB	C	DE	R_1, R_2	$(Y_d)_{ij} : (Y_e)_{ji} : (Y_{ql})_{ij} : (Y_{ud})_{ij}$
$H_{24} \mathcal{F}_j$	H_{24}	$\mathcal{T}_i \bar{H}_5$	$\bar{5}, 5$	$1 : \frac{9}{4} : \frac{9}{4} : 1$
$H_{24} \mathcal{F}_j$	H_{24}	$\mathcal{T}_i \bar{H}_5$	$\bar{5}, 45$	$1 : -\frac{9}{4} : \frac{3}{4} : -1$
$H_{24} \mathcal{F}_j$	H_{24}	$\mathcal{T}_i \bar{H}_5$	$\overline{45}, 5$	$1 : \frac{3}{4} : \frac{3}{4} : 1$
$H_{24} \mathcal{F}_j$	H_{24}	$\mathcal{T}_i \bar{H}_5$	$\overline{45}, 45$	$1 : x : -\frac{x}{3} : -1$
$H_{24} \mathcal{F}_j$	H_{24}	$\mathcal{T}_i \bar{H}_5$	$\overline{70}, 5$	$1 : \frac{9}{8} : \frac{9}{8} : 1$
$H_{24} \mathcal{F}_j$	H_{24}	$\mathcal{T}_i \bar{H}_5$	$\overline{70}, 45$	$1 : -\frac{9}{8} : \frac{3}{8} : -1$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\bar{5}, 10$	$1 : -9 : -\frac{3}{2} : -4$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\bar{5}, 15$	$1 : 0 : \frac{3}{2} : 0$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\overline{45}, 10$	$1 : 9 : -\frac{1}{2} : 4$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\overline{45}, 40$	$1 : 0 : 1 : 1$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\overline{45}, 175$	$1 : \frac{18}{19} : \frac{23}{38} : \frac{16}{19}$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\overline{70}, 15$	$1 : 0 : \frac{3}{4} : 0$
$H_{24} \mathcal{F}_j$	\bar{H}_5	$H_{24} \mathcal{T}_i$	$\overline{70}, 175$	$1 : \frac{9}{7} : \frac{33}{28} : \frac{8}{7}$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\bar{5}, 10$	$1 : 6 : -\frac{2}{3} : \frac{8}{3}$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\bar{5}, 15$	$1 : 0 : \frac{2}{3} : 0$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\overline{45}, 10$	$1 : -18 : -2 : -8$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\overline{45}, 40$	$1 : 0 : 1 : 1$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\overline{45}, 175$	$1 : \frac{36}{23} : \frac{38}{23} : \frac{32}{23}$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\overline{70}, 15$	$1 : 0 : \frac{4}{3} : 0$
$H_{24} \bar{H}_5$	\mathcal{F}_j	$H_{24} \mathcal{T}_i$	$\overline{70}, 175$	$1 : \frac{12}{11} : \frac{28}{33} : \frac{32}{33}$
$\mathcal{F}_j \bar{H}_5$	\mathcal{T}_i	$H_{24} H_{24}$	$\overline{10}, 1$	$1 : 1 : 1 : 1$
$\mathcal{F}_j \bar{H}_5$	\mathcal{T}_i	$H_{24} H_{24}$	$\overline{10}, 24$	$1 : 6 : 1 : -4$
$\mathcal{F}_j \bar{H}_5$	\mathcal{T}_i	$H_{24} H_{24}$	$\overline{10}, 75$	$1 : -3 : 1 : -1$
$\mathcal{F}_j \bar{H}_5$	\mathcal{T}_i	$H_{24} H_{24}$	$\overline{15}, 24$	$1 : 0 : -1 : 0$

Table 13: Continuation of table 12: Y_{TF} -like CG ratios for the effective dimension 6 operators $W \supset (AB)_{R_1} C(DE)_{R_2}$ corresponding to the the right diagram in figure 9. Note another combination with a free parameter x . See main text for more details.

AB	C	DE	R_1, R_2	$(Y_u)_{ij} : (Y_u)_{ji} : (Y_{qq})_{ij} : (Y_{ue})_{ij} : (Y_{ue})_{ji}$
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	10, 10	1 : 1 : $-\frac{1}{4}$: 6 : 6
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	10, 40	1 : -8 : -1 : 0 : -12
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	15, 40	1 : 0 : 1 : 0 : 0
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	40, 10	1 : $-\frac{1}{8}$: $\frac{1}{8}$: $\frac{3}{2}$: 0
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	40, 15	0 : 1 : 1 : 0 : 0
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	40, 175	1 : $\frac{23}{32}$: $\frac{19}{32}$: $\frac{3}{4}$: 0
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	175, 40	1 : $\frac{32}{23}$: $\frac{19}{23}$: 0 : $\frac{24}{23}$
$H_{24}\mathcal{T}_i$	H_5	$H_{24}\mathcal{T}_j$	175, 175	1 : 1 : $\frac{41}{40}$: $\frac{6}{5}$: $\frac{6}{5}$
$\mathcal{T}_i\mathcal{T}_j$	H_5	$H_{24}H_{24}$	$\bar{5}, 1$	1 : 1 : 1 : 1 : 1
$\mathcal{T}_i\mathcal{T}_j$	H_5	$H_{24}H_{24}$	$\bar{5}, 24$	1 : 1 : $-\frac{2}{3}$: $-\frac{2}{3}$: $-\frac{2}{3}$
$\mathcal{T}_i\mathcal{T}_j$	H_5	$H_{24}H_{24}$	$\overline{45}, 24$	1 : -1 : 0 : -2 : 2
$\mathcal{T}_i\mathcal{T}_j$	H_5	$H_{24}H_{24}$	$\overline{45}, 75$	1 : -1 : 0 : 1 : -1
$\mathcal{T}_i\mathcal{T}_j$	H_5	$H_{24}H_{24}$	$\overline{50}, 75$	0 : 0 : 1 : -2 : -2
\mathcal{T}_iH_5	\mathcal{T}_j	$H_{24}H_{24}$	$\overline{10}, 1$	1 : 1 : 1 : 1 : 1
\mathcal{T}_iH_5	\mathcal{T}_j	$H_{24}H_{24}$	$\overline{10}, 24$	1 : $-\frac{1}{4}$: $-\frac{1}{4}$: $-\frac{3}{2}$: 1
\mathcal{T}_iH_5	\mathcal{T}_j	$H_{24}H_{24}$	$\overline{10}, 75$	1 : -1 : -1 : 3 : 1
\mathcal{T}_iH_5	\mathcal{T}_j	$H_{24}H_{24}$	$\overline{40}, 24$	1 : 2 : -1 : 0 : -2
\mathcal{T}_iH_5	\mathcal{T}_j	$H_{24}H_{24}$	$\overline{40}, 75$	1 : -1 : $\frac{1}{2}$: 0 : -2

Table 14: Y_{TT} -like CG ratios for the effective dimension 6 operators $W \supset (AB)_{R_1}C(DE)_{R_2}$ corresponding to the the right diagram in figure 9.

AB	C	DE	R_1, R_2	$(Y_u)_{ij} : (Y_u)_{ji} : (Y_{qq})_{ij} : (Y_{ue})_{ij} : (Y_{ue})_{ji}$
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	10, 5	1 : -4 : $-\frac{2}{3}$: $\frac{8}{3}$: -4
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	10, 45	1 : 4 : 0 : 8 : 12
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	15, 45	1 : 0 : 1 : 0 : 0
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	40, 5	1 : $\frac{1}{2}$: $\frac{1}{3}$: $\frac{2}{3}$: 0
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	40, 45	1 : $-\frac{7}{2}$: -3 : 2 : 0
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	40, 70	1 : $\frac{1}{2}$: $\frac{2}{3}$: $\frac{4}{3}$: 0
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	175, 45	1 : $\frac{16}{19}$: $\frac{21}{19}$: $\frac{32}{19}$: $\frac{24}{19}$
$H_{24}\mathcal{T}_i$	\mathcal{T}_j	$H_{24}H_5$	175, 70	1 : $\frac{8}{7}$: $\frac{20}{21}$: $\frac{16}{21}$: $\frac{8}{7}$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{5}, 5$	1 : 1 : $\frac{4}{9}$: $\frac{4}{9}$: $\frac{4}{9}$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{5}, 45$	1 : 1 : $\frac{4}{3}$: $\frac{4}{3}$: $\frac{4}{3}$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{5}, 70$	1 : 1 : $\frac{8}{9}$: $\frac{8}{9}$: $\frac{8}{9}$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{45}, 5$	1 : -1 : 0 : $\frac{4}{3}$: $-\frac{4}{3}$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{45}, 45$	1 : -1 : 0 : x : $-x$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{45}, 70$	1 : -1 : 0 : $\frac{8}{3}$: $-\frac{8}{3}$
$\mathcal{T}_i\mathcal{T}_j$	H_{24}	$H_{24}H_5$	$\bar{50}, 45$	0 : 0 : 1 : -2 : -2
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	10, $\bar{10}$	1 : 16 : 1 : 16 : 36
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	10, $\bar{40}$	1 : -2 : $-\frac{1}{2}$: 4 : 0
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	15, $\bar{10}$	1 : 0 : 1 : 0 : 0
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	40, $\bar{10}$	1 : 1 : 1 : 1 : 0
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	40, $\bar{40}$	1 : x : $-\frac{1}{2}$: $-2x$: 0
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	175, $\bar{10}$	1 : $\frac{64}{61}$: 1 : $\frac{64}{61}$: $\frac{72}{61}$
$H_{24}\mathcal{T}_i$	H_{24}	\mathcal{T}_jH_5	175, $\bar{40}$	1 : 4 : $-\frac{1}{2}$: -8 : 0

Table 15: Continuation of table 14: Y_{TT} -like CG ratios for the effective dimension 6 operators $W \supset (AB)_{R_1}C(DE)_{R_2}$ corresponding to the the right diagram in figure 9. Note two combinations have a free parameter x . See main text for more details.

while one stays at the electroweak scale. The simplest term generating such a mixing is

$$W \supset H_{24} H_5 \bar{H}_{45} \propto H_u H_d^{45} - \frac{2}{\sqrt{3}} T \bar{T}^{45}, \quad (\text{A.13})$$

where we suppressed any additional MSSM multiplets in \bar{H}_{45} . Since a $\overline{45}$ contains more potentially dangerous MSSM multiplets, it is natural to have the heavy linear combination be predominantly in the $\overline{45}$. Then it is possible to treat \bar{H}_{45} like a messenger field and the renormalizable operator $\mathcal{FT}\bar{H}_{45}$ turns into the non-renormalizable operator $(\mathcal{FT})_{45}(H_{24}\bar{H}_5)_{\overline{45}}$, cf. table 10. Analogous limits can be deduced trivially. If the approximation $m_{45} \gg \langle H_{24} \rangle$ does not hold, one has to take into account the full mass matrix for the Higgs doublets including the term in equation (A.13).

B Two-loop RGEs in Extensions to the MSSM

The renormalization group equations for gauge couplings at two-loop are given by [37]

$$\mu \frac{d}{d\mu} g_a = \frac{g_a^3}{16\pi^2} b_a + \frac{g_a^3}{(16\pi^2)^2} \left(\sum_{b=1}^3 B_{ab} g_b^2 - \sum_f C_a^f \text{tr}(Y_f^\dagger Y_f) \right), \quad (\text{B.1})$$

in the $\overline{\text{DR}}$ renormalization scheme, where μ is the renormalization scale and f runs over all Yukawa coupling matrices. In the MSSM, the beta function coefficients are given by (in GUT normalisation for g_1)

$$b_a = \begin{pmatrix} \frac{33}{5} \\ 1 \\ -3 \end{pmatrix}, \quad B_{ab} = \begin{pmatrix} \frac{199}{25} & \frac{27}{5} & \frac{88}{5} \\ \frac{9}{5} & 25 & 24 \\ \frac{11}{5} & 9 & 14 \end{pmatrix}, \quad (\text{B.2})$$

and

$$C_a^{u,d,e} = \begin{pmatrix} \frac{26}{5} & \frac{14}{5} & \frac{18}{5} \\ 6 & 6 & 2 \\ 4 & 4 & 0 \end{pmatrix}. \quad (\text{B.3})$$

where the first column stands for u , the second column for d and the third for e . Additional colour triplet and weak doublet pairs, as those contained in $\mathbf{5}, \bar{\mathbf{5}}$ representations, contribute (per pair) at one-loop with

$$b_a^{(5,T)} = \begin{pmatrix} \frac{2}{5} \\ 0 \\ 1 \end{pmatrix}, \quad b_a^{(5,D)} = \begin{pmatrix} \frac{3}{5} \\ 1 \\ 0 \end{pmatrix}, \quad (\text{B.4})$$

and at two-loop with

$$B_{ab}^{(5,T)} = \begin{pmatrix} \frac{8}{75} & 0 & \frac{32}{15} \\ 0 & 0 & 0 \\ \frac{4}{15} & 0 & \frac{34}{3} \end{pmatrix}, \quad B_{ab}^{(5,D)} = \begin{pmatrix} \frac{9}{25} & \frac{9}{5} & 0 \\ \frac{3}{5} & 7 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (\text{B.5})$$

SU(2) triplets, SU(3) octets and leptoquark superfields¹³ from an adjoint contribute (per chiral superfield) at one-loop with

$$b_a^{(24,T)} = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, b_a^{(24,O)} = \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix}, b_a^{(24,L)} = \begin{pmatrix} \frac{5}{2} \\ \frac{3}{2} \\ 1 \end{pmatrix}, \quad (\text{B.6})$$

and at two-loop with

$$B_{ab}^{(24,T)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 24 & 0 \\ 0 & 0 & 0 \end{pmatrix}, B_{ab}^{(24,O)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 54 \end{pmatrix}, B_{ab}^{(24,L)} = \begin{pmatrix} \frac{25}{6} & \frac{15}{2} & \frac{40}{3} \\ \frac{5}{2} & \frac{21}{2} & 8 \\ \frac{5}{3} & 3 & \frac{34}{3} \end{pmatrix}. \quad (\text{B.7})$$

Additional Yukawa couplings between SM fermion superfields and a pair of colour triplet/anti-triplet contribute with

$$C_a^{qq,ue,ql,ud} = \begin{pmatrix} \frac{6}{5} & \frac{28}{5} & \frac{14}{5} & \frac{24}{5} \\ 6 & 0 & 6 & 0 \\ 6 & 2 & 4 & 6 \end{pmatrix}. \quad (\text{B.8})$$

In our numerical analysis we have assumed that $Y_{qq} = Y_{ue} = Y_u$ and $Y_{ql} = Y_{ud} = Y_d$ (as motivated by minimal SU(5)). We have checked that this approximation changes our results only negligibly.

C Discussion of messenger fields

We discuss now the messenger fields appearing in Figure 6 of the model presented in section 4.1. There are three supergraphs generating the superpotential term y_{12} . The renormalizable superpotential corresponding to Figure 6 is

$$\begin{aligned} W \supset & \gamma_1 H_5 \mathcal{T}_1 Z_{10,4} + \gamma_2 H_5 \mathcal{T}_2 Z_{10,3} + \gamma_3 \mathcal{T}_3 \theta_1 \bar{Z}_{10,3} + \gamma_\theta \theta_1^2 Z_1 \\ & + \lambda_1 \theta_2 Z_2 \bar{Z}_1 + \lambda_{10} \theta_2 Z_{10,3} \bar{Z}_{10,4} \\ & + \eta_1 \mathcal{T}_1 \bar{Z}_{10,3} \bar{Z}_2 + \eta_2 \mathcal{T}_2 \bar{Z}_{10,4} \bar{Z}_2 + \eta'_2 \mathcal{T}_2 \bar{Z}_{10,3} \bar{Z}_1 \\ & + M_1 Z_1 \bar{Z}_1 + M_2 Z_2 \bar{Z}_2 + M_{10,3} Z_{10,3} \bar{Z}_{10,3} + M_{10,4} Z_{10,4} \bar{Z}_{10,4}, \end{aligned} \quad (\text{C.1})$$

where we explicitly denote the coupling constants at the ends of a diagram with γ_x , coupling constants in the middle of the diagrams with λ_x if they involve θ_j and η_i if they involve \mathcal{T}_i and messenger masses with M_x . After Z_x and \bar{Z}_x are integrated out and the flavons θ_i

¹³Note that leptoquark superfields can only appear in Dirac pairs due to their charges.

obtain VEVs, the elements of the up-type Yukawa matrix Y_u of eq. (4.2) are given by

$$\begin{aligned}
y_{11} &= \gamma_1 \lambda_{10} \eta_1 \lambda_1 \gamma_\theta \frac{\langle \theta_1 \rangle^2 \langle \theta_2 \rangle^2}{M_1 M_2 M_{10,3} M_{10,4}} , \\
y_{12} &= \gamma_2 \eta_1 \lambda_1 \gamma_\theta \frac{\langle \theta_1 \rangle^2 \langle \theta_2 \rangle}{M_1 M_2 M_{10,3}} + \gamma_1 \eta_2 \lambda_1 \gamma_\theta \frac{\langle \theta_1 \rangle^2 \langle \theta_2 \rangle}{M_1 M_2 M_{10,4}} + \gamma_1 \lambda_{10} \eta'_2 \gamma_\theta \frac{\langle \theta_1 \rangle^2 \langle \theta_2 \rangle}{M_1 M_{10,3} M_{10,4}} , \\
y_{22} &= \gamma_2 \eta'_2 \gamma_\theta \frac{\langle \theta_1 \rangle^2}{M_1 M_{10,3}} , \\
y_{13} &= \gamma_1 \lambda_{10} \gamma_3 \frac{\langle \theta_1 \rangle \langle \theta_2 \rangle}{M_{10,3} M_{10,4}} , \\
y_{23} &= \gamma_2 \gamma_3 \frac{\langle \theta_1 \rangle}{M_{10,3}} ,
\end{aligned} \tag{C.2}$$

and y_{33} is a renormalizable Yukawa coupling coefficient.

Removing the messenger pair Z_2, \bar{Z}_2 from the spectrum eliminates two supergraphs and thus the first two terms contributing to y_{12} .¹⁴ However, without $Z_2 \bar{Z}_2$, evaluating y_{12} yields

$$\begin{aligned}
y_{12} &= \gamma_1 \lambda_{10} \eta'_2 \gamma_\theta \frac{\langle \theta_1 \rangle^2 \langle \theta_2 \rangle}{M_1 M_{10,3} M_{10,4}} = y_{13} \frac{\langle \theta_1 \rangle}{M_1} \frac{\eta'_2 \gamma_\theta}{\gamma_3} \\
&= y_{13} y_{22} \frac{M_{10,3}}{\langle \theta_1 \rangle} \frac{1}{\gamma_2 \gamma_3} = \frac{y_{13} y_{22}}{y_{23}} .
\end{aligned} \tag{C.3}$$

This relation is not phenomenologically viable as it would imply $\theta_C = \theta_{13}/\theta_{23}$. To fit Y_u to the observed data an additional degree of freedom is needed. In our model this is realised through $Z_2 \bar{Z}_2$ enabling additional diagrams contributing to y_{12} .

D Simultaneous R-symmetry and GUT breaking

It was claimed in the literature that R-symmetries and four-dimensional (SUSY) GUTs are not compatible, as in the presence of the R-symmetry not all the components of the the GUT breaking Higgs representation(s) can be made heavy enough (cf., e.g., [26]). On the other hand, in flavour models, R-symmetries are often used for generating the required VEVs for the family symmetry breaking Higgs fields (i.e. the flavons) as well as for spontaneous breaking of CP. In the following we present some simple examples that show that it is possible - in particular with discrete R-symmetries - to simultaneously break the R-symmetry and the GUT and hence to avoid this apparent conflict. We will also illustrate that GUT flavour models can rely on a discrete R-symmetry for the flavon VEV alignment, such that our setup can be used to construct flavour models with non-Abelian family symmetries and spontaneous CP violation.

¹⁴A different pair of fields Z'_2, \bar{Z}'_2 would need to be introduced anyway (using different charge assignment than Z_2, \bar{Z}_2) in order to generate y_{11} . The charges can be assigned such that no extra contributions to y_{12} appear.

Simple Example Consider as example a discrete \mathbb{Z}_4^R R-symmetry under which the superfield S is charged and the superfield H is uncharged (for the beginning we assume them to be SU(5) singlets). The fields are also charged under an additional conventional \mathbb{Z}_4 symmetry with charges 1 and 3 respectively. We consider only the two lowest order superpotential terms, neglecting the contributions with S^5 and higher:

$$W_R = aSH + cS^3H^3/M_{\text{Pl}}^3. \quad (\text{D.1})$$

The F-terms for this simple example lead to the conditions

$$aH + 3cS^2H^3/M_{\text{Pl}}^3 = 0 \quad (\text{D.2})$$

$$aS + 3cS^3H^2/M_{\text{Pl}}^3 = 0. \quad (\text{D.3})$$

The allowed solutions are the trivial VEVs $\langle S \rangle = \langle H \rangle = 0$ and a non-trivial solution $\langle S \rangle^2 = -aM_{\text{Pl}}^3/3c\langle H \rangle^2$. If we assign S and H under SU(5) as adjoints (similarly to superpotential (c) where we had the two fields H_{24} and H'_{24}) W_R is SU(5) invariant (by taking the appropriate contractions) and importantly, the R-symmetry and SU(5) are simultaneously broken by the non-trivial VEV configuration. We can achieve a phenomenological viable model by breaking a discrete R-symmetry at the GUT scale.

In general, leaving additional symmetry (or symmetries) unspecified, S may be a singlet of the GUT symmetry group and we denote the (polynomial) functions of the superfield H that make the respective terms invariant as $A(H)$, $C(H)$ (which include the associated couplings and M_{Pl} suppressions). We write

$$W_R = SA(H) + S^3C(H), \quad (\text{D.4})$$

leading to the F-term equations

$$A(H) + 3C(H)S^2 = 0, \quad (\text{D.5})$$

$$\frac{dA(H)}{dH}S + \frac{dC(H)}{dH}S^3 = 0. \quad (\text{D.6})$$

The generalised solution breaking the R-symmetry and the GUT gauge group is then

$$\langle S \rangle^2 = - \left\langle \frac{A(H)}{3C(H)} \right\rangle, \quad (\text{D.7})$$

$$\left\langle \frac{dA(H)}{dH} \right\rangle = \left\langle \frac{A(H)}{3C(H)} \frac{dC(H)}{dH} \right\rangle. \quad (\text{D.8})$$

Note the second equation, $\frac{dA}{A} = \frac{dC}{3C}$, constrains the allowed functions $A(H)$ and $C(H)$ which indirectly imposes conditions on the unspecified additional symmetries.

Generalisations While the R-symmetry must be discrete for the crucial interplay between two terms, it needs not be a \mathbb{Z}_4^R and generalising to other \mathbb{Z}_N^R symmetries is straightforward

$$W_R^N = SA(H) + S^{N-1}C(H). \quad (\text{D.9})$$

This can be further generalised to multiple fields. Consider, for instance, S , R and T charged under a \mathbb{Z}_4^R , some unspecified symmetries with fields H , I , J , and generalized functions $A_{S,R,T}(H, I, J)$ as well as $C(H, I, J)$, such that keeping only the necessary lowest order terms in S , R , T we write

$$W_R = S A_S(H, I, J) + R A_R(H, I, J) + T A_T(H, I, J) + S R T C(H, I, J). \quad (\text{D.10})$$

After some manipulation we find again a non-trivial solution breaking \mathbb{Z}_4^R and the GUT symmetry

$$\frac{A_S}{\langle RT \rangle} = \frac{A_R}{\langle TS \rangle} = \frac{A_T}{\langle SR \rangle} = -C, \quad (\text{D.11})$$

provided the derivatives with respect to each of the GUT superfields H , I , J fulfill

$$\frac{\partial A_S}{A_S} + \frac{\partial A_R}{A_R} + \frac{\partial A_T}{A_T} + \frac{\partial C}{C} = 0. \quad (\text{D.12})$$

The simple examples above illustrate that a simultaneous breaking of R- and GUT symmetry is possible. VEV alignments required by flavour models can therefore still be obtained within GUTs by having additional superfields charged under the R-symmetry.

Flavour Alignment As a very simple example we discuss the ‘‘alignment’’ of a GUT singlet flavon ϕ charged under a non-Abelian family symmetry via a driving field P . As additional symmetries we impose \mathbb{Z}_4^R and a conventional \mathbb{Z}_n . The allowed renormalizable superpotential is

$$W = P \left(\frac{\phi^n}{\Lambda^{n-2}} + M^2 \right) + \kappa P^3, \quad (\text{D.13})$$

where Λ is a generic messenger scale and M a mass parameter. Minimising the F-term conditions we find two possible solutions:

$$\text{solution A: } \langle P \rangle = 0 \text{ and } \langle \phi \rangle^n = -M^2 \Lambda^{n-2}, \quad (\text{D.14})$$

$$\text{solution B: } \langle P \rangle^2 = -\frac{M^2}{3\kappa} \text{ and } \langle \phi \rangle = 0. \quad (\text{D.15})$$

In a flavour model we want the flavon to get a non-vanishing VEV so that we would adopt solution A there. Furthermore solution A shows how the discrete vacuum alignment method [38] can be generalised to models with discrete R-symmetries. This method was invented in the context of spontaneous CP violation. If CP is promoted to be fundamental all the phases of the parameters in eq. (D.13) are fixed and hence the phase of the flavon VEV is fixed as well (up to a discrete choice).

References

- [1] Z. Berezhiani and Z. Tavartkiladze, Phys. Lett. B **396** (1997) 150 [hep-ph/9611277].

- [2] G. Altarelli, F. Feruglio and I. Masina, JHEP **0011** (2000) 040 [hep-ph/0007254].
- [3] H. Georgi and C. Jarlskog, Phys. Lett. B **86** (1979) 297.
- [4] H. Murayama, Y. Okada and T. Yanagida, Prog. Theor. Phys. **88** (1992) 791.
- [5] D. -X. Zhang and J. -h. Zheng, JHEP **1212** (2012) 087 [arXiv:1212.5852 [hep-ph]].
- [6] H. Murayama and A. Pierce, Phys. Rev. D **65** (2002) 055009 [hep-ph/0108104].
- [7] A. Masiero, D. V. Nanopoulos, K. Tamvakis and T. Yanagida, Phys. Lett. B **115** (1982) 380.
- [8] B. Grinstein, Nucl. Phys. B **206** (1982) 387.
- [9] J. Hisano, T. Moroi, K. Tobe and T. Yanagida, Phys. Lett. B **342** (1995) 138 [hep-ph/9406417].
- [10] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86** (2012) 010001.
- [11] S. Antusch and M. Spinrath, Phys. Rev. D **79** (2009) 095004 [arXiv:0902.4644 [hep-ph]].
- [12] S. Antusch and V. Maurer, Phys. Rev. D **84** (2011) 117301 [arXiv:1107.3728 [hep-ph]].
- [13] D. Marzocca, S. T. Petcov, A. Romanino and M. Spinrath, JHEP **1111** (2011) 009 [arXiv:1108.0614 [hep-ph]].
- [14] S. Antusch, C. Gross, V. Maurer and C. Sluka, Nucl. Phys. B **866** (2013) 255 [arXiv:1205.1051 [hep-ph]].
- [15] A. Meroni, S. T. Petcov and M. Spinrath, Phys. Rev. D **86** (2012) 113003 [arXiv:1205.5241 [hep-ph]].
- [16] S. Antusch, C. Gross, V. Maurer and C. Sluka, Nucl. Phys. B **877** (2013) 772 [arXiv:1305.6612 [hep-ph]].
- [17] S. Antusch, C. Gross, V. Maurer and C. Sluka, Nucl. Phys. B **879** (2014) 19 [arXiv:1306.3984 [hep-ph]].
- [18] T. Goto and T. Nihei, Phys. Rev. D **59** (1999) 115009
- [19] I. de Medeiros Varzielas and L. Merlo, JHEP **1102** (2011) 062 [arXiv:1011.6662 [hep-ph]].
- [20] I. de Medeiros Varzielas and D. Pidt, JHEP **1303** (2013) 065 [arXiv:1211.5370 [hep-ph]].
- [21] S. Antusch, S. F. King and M. Spinrath, arXiv:1311.0877 [hep-ph].

- [22] S. Dimopoulos and H. Georgi, Nucl. Phys. B **193** (1981) 150.
- [23] B. Bajc, P. Fileviez Perez and G. Senjanovic, hep-ph/0210374.
- [24] S. Antusch and V. Maurer, JHEP **1311** (2013) 115 [arXiv:1306.6879 [hep-ph]].
- [25] L. J. Hall, Nucl. Phys. B **178** (1981) 75.
- [26] M. Fallbacher, M. Ratz and P. K. S. Vaudrevange, Phys. Lett. B **705** (2011) 503 [arXiv:1109.4797 [hep-ph]].
- [27] L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D **50** (1994) 7048 [arXiv:hep-ph/9306309]; M. S. Carena, M. Olechowski, S. Pokorski and C. E. M. Wagner, Nucl. Phys. B **426** (1994) 269 [arXiv:hep-ph/9402253]; R. Hempfling, Phys. Rev. D **49** (1994) 6168; T. Blazek, S. Raby and S. Pokorski, Phys. Rev. D **52** (1995) 4151 [arXiv:hep-ph/9504364].
- [28] S. F. King, JHEP **1307** (2013) 137 [arXiv:1304.6264 [hep-ph]]; S. F. King, JHEP **1401** (2014) 119 [arXiv:1311.3295 [hep-ph]].
- [29] S. Antusch, S. F. King, M. Malinsky and M. Spinrath, Phys. Rev. D **81** (2010) 033008 [arXiv:0910.5127 [hep-ph]].
- [30] P. Z. Skands, B. C. Allanach, H. Baer, C. Balazs, G. Belanger, F. Boudjema, A. Djouadi and R. Godbole *et al.*, JHEP **0407** (2004) 036 [hep-ph/0311123].
- [31] D. Emmanuel-Costa and S. Wiesenfeldt, Nucl. Phys. B **661** (2003) 62 [hep-ph/0302272].
- [32] S. Wiesenfeldt, Mod. Phys. Lett. A **19** (2004) 2155 [hep-ph/0407173].
- [33] P. Nath, Phys. Rev. Lett. **76** (1996) 2218 [hep-ph/9512415].
- [34] P. Nath and P. Fileviez Perez, Phys. Rept. **441** (2007) 191 [hep-ph/0601023].
- [35] V. Lucas and S. Raby, Phys. Rev. D **55** (1997) 6986 [hep-ph/9610293].
- [36] M. Spinrath, arXiv:1009.2511 [hep-ph].
- [37] D. R. T. Jones, Nucl. Phys. B **87** (1975) 127; D. R. T. Jones and L. Mezincescu, Phys. Lett. B **136** (1984) 242.
- [38] S. Antusch, S. F. King, C. Luhn and M. Spinrath, Nucl. Phys. B **850** (2011) 477 [arXiv:1103.5930 [hep-ph]].