LHC Signatures of τ -Flavoured Vector Leptoquarks

Jordan Bernigaud, ^{a, b} Monika Blanke, ^{a, b} Ivo de Medeiros Varzielas, ^c Jim Talbert, ^{d, e} and José Zurita^f

^a Institute for Astroparticle Physics (IAP), Karlsruhe Institute of Technology, Hermann-von-Helmholtz-Platz 1, D-76344 Eggenstein-Leopoldshafen, Germany,

^b Institute for Theoretical Particle Physics (TTP), Karlsruhe Institute of Technology, Engesserstrasse 7, D-76128 Karlsruhe, Germany

^c CFTP, Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Avenida Rovisco Pais 1, 1049 Lisboa, Portugal

^d DAMTP, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WA, United Kingdom

^e Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark

^f Instituto de Física Corpuscular, CSIC-Universitat de Valéncia, Valencia, Spain

E-mail: jordan.bernigaud@kit.edu, monika.blanke@kit.edu, ivo.de@udo.edu, rjt89@cam.ac.uk, jzurita@ific.uv.es

Abstract

We consider the phenomenological signatures of Simplified Models of Flavourful Leptoquarks, whose Beyond-the-Standard Model (SM) couplings to fermion generations occur via textures that are well motivated from a broad class of ultraviolet flavour models (which we briefly review). We place particular emphasis on the study of the vector leptoquark Δ_{μ} with assignments (3, 1, 2/3) under the SM's gauge symmetry, $SU(3)_C \times SU(2)_L \times U(1)_Y$, which has the tantalising possibility of explaining both $\mathcal{R}_{K^{(\star)}}$ and $\mathcal{R}_{D^{(\star)}}$ anomalies. Upon performing global likelihood scans of the leptoquark's coupling parameter space, focusing in particular on models with tree-level couplings to a single charged lepton species, we then provide confidence intervals and benchmark points preferred by low(er)-energy flavour data. Finally, we use these constraints to further evaluate the (promising) Large Hadron Collider (LHC) detection prospects of pairs of τ -flavoured Δ_{μ} , through their distinct (a)symmetric decay channels. Namely, we consider direct third-generation leptoquark and jets plus missing-energy searches at the LHC, which we find to be complementary. Depending on the simplified model under consideration, the direct searches constrain the Δ_{μ} mass up to 1400-1750 GeV when the branching fraction of Δ_{μ} is entirely to third-generation quarks (but are significantly reduced with decreased branching ratios to the third generation), whereas the missingenergy searches constrain the mass up to 1150-1500 GeV while being largely insensitive to the third-generation branching fraction.

Contents

1	Introduction	1
2	Theoretical Framework2.1Simplified Model Lagrangian2.2Simplified Models of Flavourful Leptoquarks2.3On $\mathcal{R}_{D^{(*)}}$ and Collider Complementarity	4
3	Precision Constraints and Global Likelihoods 3.1 Matching the Vector Singlet 3.2 Scanning the Vector Leptoquark Singlet	
4	LHC Recasting Exclusion Limits4.1 Reinterpretation of the Mixed $b\tau t\nu$ Search4.2 CheckMATE Exclusion Limits	
5	Summary and Outlook	17

1 Introduction

Despite the large amount of data collected and analysed at the Large Hadron Collider (LHC), no new Beyond-the-Standard Model (BSM) particles have been discovered as of yet. Nonetheless, compelling motivations for the existence of BSM physics exist, including the unsolved electroweak (EW) hierarchy, neutrino mass, and strong CP problems, the unexplained presence of a baryon asymmetry in the Universe, the lack of a confirmed dark matter candidate, and of course the flavour puzzle.

Besides these theoretical problems, perhaps the strongest experimental hints for BSM physics are the deviations observed in lepton flavour universality (LFU) tests in B meson decays, the so-called 'flavour anomalies'. The first indications for LFU-violating BSM interactions were found in 2012 by the BaBar collaboration [1,2] in the ratio

$$\mathcal{R}_{D^{(\star)}} = \frac{\mathrm{BR}\left(\overline{B} \to D^{(\star)} \tau^{-} \overline{\nu}_{\tau}\right)}{\mathrm{BR}\left(\overline{B} \to D^{(\star)} l^{-} \overline{\nu}_{l}\right)},\tag{1}$$

with $l \in \{e, \mu\}$. Over the past years, measurements of the same ratios by Belle [3–7] and LHCb [8–10] have confirmed the tension with the SM prediction, with the latest HFLAV average exhibiting a 3.4 σ deviation from the SM [11].

Due to the size of the BSM contribution required to resolve the anomaly – an $\mathcal{O}(10\%)$ enhancement at the matrix element level – new physics plausibly has to enter the relevant $b \to c\tau\nu$ transition at tree level. Possible BSM scenarios then include the exchange of a new colour-singlet charged scalar ("charged Higgs") [12–16] or vector (W') boson [17–20], or of a colour-triplet scalar or vector leptoquark (LQ) [21–33]. The latter have the advantage of being less stringently constrained by precision EW constraints and direct LHC searches. Of particular interest is the isospin-singlet vector LQ¹

$$\Delta^{\mu} \sim (\mathbf{3}, \mathbf{1}, 2/3) , \qquad (2)$$

whose respective charge assignments under the SM gauge group, $\mathcal{G}_{SM} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y$, are given on the right hand side of (2). In contrast to scalar LQ solutions, the coupling structure of Δ^{μ} is not constrained by proton decay. Besides that, Δ^{μ} is contained in the Pati-Salam gauge group $SU(4)_C \times SU(2)_L \times SU(2)_R$ unifying quarks and leptons [38], thereby providing an appealing ansatz for the construction of an ultraviolet (UV)-complete model [39–43].

From the phenomenological perspective, the vector LQ singlet gains additional appeal from the fact that it is the only single-particle solution to anomalies in both $\mathcal{R}_{D^{(\star)}}$ and $\mathcal{R}_{K^{(\star)}}$. The latter ratios, defined as

$$\mathcal{R}_{K^{(\star)}} = \frac{\mathrm{BR}\left(B \to K^{(\star)} \,\mu^+ \mu^-\right)}{\mathrm{BR}\left(B \to K^{(\star)} \,e^+ e^-\right)}\,,\tag{3}$$

exhibit a combined ~ 4σ tension [44,45] with the SM in LHCb [46–48] and Belle [49,50] data. The latter anomaly is further supported by the fact that deviations from the SM predictions are also showing in other observables sensitive to the quark-level $b \to s\mu^+\mu^-$ transition, such as P'_5 , BR($B_s \to \phi\mu^+\mu^-$) and BR($B_s \to \mu^+\mu^-$).

The new physics scale required to address the anomaly in $\mathcal{R}_{D^{(\star)}}$ is as low as a few TeV, and therefore any underlying BSM particle(s) responsible for the (potentially) new physics may be within the expected mass reach of the LHC or, eventually, future high(er)-energy colliders — see e.g. [51–54]. Numerous studies of the LHC phenomenology of LQs responsible for the $\mathcal{R}_{D^{(\star)}}$ anomaly exist, ranging from resonant LQ pair- and single-production, *t*-channel LQ exchange, and other non-resonant processes, see e.g. [37,55–65].

Of these, resonant LQ production processes yield the most direct access to the parameters of the LQ model. LQ pair production is driven by QCD interactions, and hence for a given LQ representation its cross-section yields a direct determination of the LQ mass. The branching ratios of the LQ into different final states then determine the relative coupling strengths to fermions. In combination with the observed values for $\mathcal{R}_{D^{(*)}}$, which are driven by the product of the relevant LQ coupling parameters, the measurement of LQ branching ratios into the various final states then allows for a complete determination of the parameters of the simplified model. We will elaborate more on this point in Section 2.3. In this paper, after reviewing the simplified models we consider, we take specific textures of the LQ couplings to fermions that arise from specific flavour hypotheses and analyze the impact of LHC searches on the respective models.

The remainder of the paper develops as follows: in Section 2 we review the theory formalism embedded in our simplified models, including the symmetry motivation for studying particular LQ flavour patterns. In Section 3 we use smelli [66] to perform a global likelihood scan of said couplings, and ultimately isolate a preferred parameter space at the 1σ and 2σ confidence level. This then provides benchmark points to study collider phenomenology in Section 4,

¹In what follows we will use the notation and nomenclature of [34–36] when considering flavour structures, and [37] for other Lagrangian parameters. We will refer to Δ^{μ} as the 'vector LQ singlet', which is sometimes denoted U_1 in the literature.

where we perform a reinterpretation of several LHC searches for our isospin-singlet vector LQ. Finally, we provide a summary and outlook in Section 5.

2 Theoretical Framework

In this Section we review the simplified model describing the dynamics of the vector LQ singlet Δ^{μ} in (2). We start by introducing the underlying Lagrangian in Section 2.1. Subsequently in Section 2.2 we turn to the discussion of simple LQ coupling structures motivated by flavour symmetries. Turning our attention to one specific case, the τ -isolation pattern, we then outline how the measurements of both the LFU ratios $\mathcal{R}_{D^{(*)}}$ and the LQ pair-production and decay rates at the LHC collude in the determination of the simplified model parameters.

2.1 Simplified Model Lagrangian

When added to the SM field content, the vector LQ singlet Δ^{μ} introduced in (2) sources the following kinetic term and gauge interactions

$$\mathcal{L} \supset -\frac{1}{2}\Delta^{\dagger}_{\mu\nu}\Delta^{\mu\nu} + ig_s(1-k_s)\Delta^{\dagger}_{\mu}T^A\Delta_{\nu}G^{A,\mu\nu} + i\frac{2g'(1-k_Y)}{3}\Delta_{\mu}\Delta_{\nu}B^{\mu\nu}, \qquad (4)$$

where the leptoquark field strength is given by

$$\Delta_{\mu\nu} = D_{\mu}\Delta_{\nu} - D_{\nu}\Delta_{\mu}\,,\tag{5}$$

in terms of the gauge-covariant derivative

$$D_{\mu} = \partial_{\mu} + ig_{s}T^{A}G_{\mu}^{A} + i\frac{2g'}{3}B_{\mu}.$$
 (6)

In the above equations T^A are colour generators, g_s and g' are the standard QCD and hypercharge gauge couplings, and we study two scenarios for the $k_{s,Y}$ parameters: (**A**) $k_Y = k_s = 0$, which tames divergences in leptoquark-gauge boson scattering and dipole processes — see [24,67] for details — and is also motivated by UV-completing the vector LQ singlet as a gauge boson of an extended gauge symmetry [25], and (**B**) $k_Y = k_s = 1$, which corresponds to the so-called minimal coupling scenario as may appear in strongly coupled UV models [58]. For simplicity we will also refer to Scenarios (**A**) and (**B**) as k = 0 and k = 1, respectively.

In addition to the above interactions, gauge-invariant coupling terms of the form²

$$\mathcal{L} \supset x_{1,ij}^{LL} \bar{Q}_L^{i,a} \gamma^\mu \Delta_\mu L_L^{j,a} + x_{1,ij}^{RR} \bar{d}_R^i \gamma^\mu \Delta_\mu e_R^j + x_{1,ij}^{\overline{RR}} \bar{u}_R^i \gamma^\mu \Delta_\mu \nu_R^j + \text{h.c.} , \qquad (7)$$

also appear, where $\{i, j\}$ denote flavour indices, $\{a, b\}$ denote SU(2) indices, and k = 1, 2, 3 indicate the Pauli matrices. For a thorough review of the physics of leptoquarks, see e.g. [68]. Moving to the SM fermion mass basis via

$$u_L \to U_u u_L$$
, $d_L \to U_d d_L$, $l_L \to U_l l_L$, $\nu_L \to U_\nu \nu_L$,

²While right-handed neutrinos play no role throughout our phenomenological analysis, we introduce them here in order to allow for a complete discussion of flavour symmetries.

$$u_R \to U_U u_R$$
, $d_R \to U_D d_R$, $E_R \to U_E E_R$, $\nu_R \to U_R \nu_R$, (8)

and further decomposing $SU(2)_L$ indices, one then finds that (7) expands to

$$\mathcal{L} \supset (U_u^{\dagger} x_1^{LL} U_{\nu})_{ij} \bar{u}_L^i \gamma^{\mu} \Delta_{\mu} \nu_L^j + (U_d^{\dagger} x_1^{LL} U_l)_{ij} \bar{d}_L^i \gamma^{\mu} \Delta_{\mu} l_L^j$$

$$+ (U_D^{\dagger} x_1^{RR} U_E)_{ij} \bar{d}_R^i \gamma^{\mu} \Delta_{\mu} E_R^j + (U_U^{\dagger} x_1^{RR} U_R)_{ij} \bar{u}_R^i \gamma^{\mu} \Delta_{\mu} \nu_R^j$$

$$+ \text{h.c.}, \qquad (9)$$

such that the novel BSM interactions between generations of quarks and leptons are manifest, including terms with left-left (LL) and right-right (RR) chiral structure (although we do not consider the terms with RR chiral structure in our analysis below). As is clear, these couplings are all 3×3 matrices in flavour space, such that (e.g.)³

$$\left(U_d^{\dagger} x_1^{LL} U_l\right) \equiv \lambda_{dl} = \begin{pmatrix} \lambda_{de} \ \lambda_{d\mu} \ \lambda_{d\tau} \\ \lambda_{se} \ \lambda_{s\mu} \ \lambda_{s\tau} \\ \lambda_{be} \ \lambda_{b\mu} \ \lambda_{b\tau} \end{pmatrix},$$
(10)

so implying that the LL $u-\nu$ coupling is related to λ_{dl} via SU(2)_L rotations, $\lambda_{u\nu} = U_{\text{CKM}} \lambda_{dl} U_{\text{PMNS}}$. Here U_{CKM} and U_{PMNS} are the standard quark and lepton mixing matrices of the SM, defined by

$$U_{\rm CKM} \equiv U_u^{\dagger} U_d \,, \qquad U_{\rm PMNS} \equiv U_l^{\dagger} U_{\nu} \,, \tag{11}$$

and whose matrix elements are constrained by a host of low-energy precision flavour data — see e.g. [69,70].

Finally, the RR terms of (9) are, a priori, fully independent. In the remainder of our analysis we will set these couplings to zero, which we motivate below.

2.2 Simplified Models of Flavourful Leptoquarks

In general, one can study the phenomenology of (10) with arbitrary values/shapes for the couplings λ_{dl} . However, it is appealing to instead examine textures that are motivated by both experimental and theoretical considerations. To that end, we will study lepton isolation patterns of the form

$$\lambda_{dl}^{[e]} = \begin{pmatrix} \lambda_{de} \ 0 \ 0 \\ \lambda_{se} \ 0 \ 0 \\ \lambda_{be} \ 0 \ 0 \end{pmatrix}, \quad \lambda_{dl}^{[\mu]} = \begin{pmatrix} 0 \ \lambda_{d\mu} \ 0 \\ 0 \ \lambda_{s\mu} \ 0 \\ 0 \ \lambda_{b\mu} \ 0 \end{pmatrix}, \quad \lambda_{dl}^{[\tau]} = \begin{pmatrix} 0 \ 0 \ \lambda_{d\tau} \\ 0 \ 0 \ \lambda_{s\tau} \\ 0 \ 0 \ \lambda_{b\tau} \end{pmatrix}, \quad (12)$$

where the meaning of the red entries in the first row will be explained below. Such matrices are some of the minimal patterns motivated by the flavour-symmetry breaking embedded in the Simplified Models of Flavourful Leptoquarks (SMFL) developed in [34–36]. The principle

³Observe that in [34,35] the vector LQ singlet d-l coupling features an additional superscript, $\lambda_{dl}^{V_1}$. Since we will not study the scalar triplet or vector triplet states in this work, we remove this superscript for simplicity.

assumption of SMFL is that the LQ couplings to fermions of (e.g.) (10) are invariant under Abelian residual family symmetries (RFS),

$$\exists \{Q, L\}, \quad T_Q^{\dagger} \lambda_{QL} T_L \stackrel{!}{=} \lambda_{QL} \,. \tag{13}$$

Here $T_{Q,L}$ are (reducible) generator representations of said RFS in arbitrary quark (Q) or lepton (L) family sectors, which simultaneously act on the SM Yukawa sector, where it is well known that each family's mass sector is invariant under U(1)³ RFS (in the broken phase) and that, if present, a Majorana neutrino mass term is instead invariant under a Klein $\mathbb{Z}_2 \times \mathbb{Z}_2$ [71,72]. When RFS are interpreted as remnants of the breakdown of a UV parent symmetry, e.g. through a breaking chain

$$\mathcal{G}_{\mathcal{F}} \to \begin{cases} \mathcal{G}_{\mathcal{L}} \to \begin{cases} \mathcal{G}_{\nu} \\ \mathcal{G}_{l} \\ \\ \mathcal{G}_{\mathcal{Q}} \to \begin{cases} \mathcal{G}_{u} \\ \mathcal{G}_{d} \end{cases} \end{cases}$$
(14)

where $\mathcal{G}_{u,d,\nu,l}$ denote the RFS controlling infrared (IR) flavour structures and $\mathcal{G}_{F,\mathcal{L},\mathcal{Q}}$ are larger parent flavour groups,⁴ they can be used to algorithmically study the origins of CKM and PMNS mixing matrices [72–84], control FCNC in multi-Higgs-doublet models [85], and of course structure the LQ couplings of interest here. Critically, this analysis can be done without reference to the details of the UV flavour model's dynamics, and is therefore a largely modelindependent formalism for studying (B)SM flavour.

We leave the details of the RFS mechanism embedded in SMFL to [34–36], and proceed by considering (13) with respect to the d-l operator only,⁵ where the 3×3 coupling is constrained entry-by-entry through the RFS relation

$$\begin{pmatrix} e^{i(-\alpha_d+\alpha_l)} \lambda_{de} & e^{i(-\alpha_d+\beta_l)} \lambda_{d\mu} & e^{i(-\alpha_d+\gamma_l)} \lambda_{d\tau} \\ e^{i(-\beta_d+\alpha_l)} \lambda_{se} & e^{i(-\beta_d+\beta_l)} \lambda_{s\mu} & e^{i(-\beta_d+\gamma_l)} \lambda_{s\tau} \\ e^{i(-\gamma_d+\alpha_l)} \lambda_{be} & e^{i(-\gamma_d+\beta_l)} \lambda_{b\mu} & e^{i(-\gamma_d+\gamma_l)} \lambda_{b\tau} \end{pmatrix} \stackrel{!}{=} \begin{pmatrix} \lambda_{de} & \lambda_{d\mu} & \lambda_{d\tau} \\ \lambda_{se} & \lambda_{s\mu} & \lambda_{s\tau} \\ \lambda_{be} & \lambda_{b\mu} & \lambda_{b\tau} \end{pmatrix} .$$
(15)

This invariance is clearly not realized in the absence of special relationships amongst the phases of the RFS generator, which are themselves IR realizations of UV flavour-symmetry breaking in specific directions of flavour space. Indeed, the patterns of (12) appear when one of the T_l phases is also equal to $\beta_d = \gamma_d$, e.g. $\alpha_l = \beta_d = \gamma_d$ (which gives the first texture, etc.). These matrices are given in the fermion mass basis, and the red entries in the top row of (12) highlight that it is perhaps more interesting to consider RFS which distinguish at least two fermion generations, thereby forcing $\alpha_d \neq \beta_d = \gamma_d$, which forbids these *d*-quark entries. It was also noted in [34] that zero entries in the first row are consistent with scenarios where the RFS successfully controls the dominant Cabibbo mixing observed in the CKM matrix, thereby connecting potentially anomalous signals of new physics with partial solutions to the SM's longstanding flavour puzzle. Furthermore, it was generically shown that (12) can

⁴The parent group can be continuous or discrete, Abelian or non-Abelian.

 $^{{}^{5}}$ In [34] the consequences of applying (13) to all relevant family sectors were explored. While the presence of more symmetry removes parametric degrees of freedom in a simplified model setup, it is also more challenging to accommodate in UV flavour models – cf. [35, 36] (e.g.) for some discussion on this point.

arise from the breakdown of non-Abelian family symmetries [35], which can be described by an effective Lagrangian composed of non-renormalizable interactions between scalar flavons and SM fermion multiplets [36,86]. In short, evidence of new SMFL physics can be directly connected to more complete models of (B)SM flavour physics.

2.3 On $\mathcal{R}_{D^{(*)}}$ and Collider Complementarity

Precision flavour constraints from *B*-meson decays and other low-energy processes give information that is potentially complementary to direct searches at the LHC. Consider the LFU ratio $\mathcal{R}_{D^{(\star)}}$, which for the flavoured LQ Δ_{μ} we consider is approximately given by [26]

$$\mathcal{R}_{D^{(\star)}} \simeq \mathcal{R}_{D^{(\star)}}^{\mathrm{SM}} \cdot \left[1 + \frac{1}{\sqrt{2}G_F V_{cb}} \operatorname{Re}\left(\lambda \lambda^{\star}|_{\tau} - \lambda \lambda^{\star}|_l\right) \left(\frac{\mathrm{TeV}}{M_{\Delta}}\right)^2 \right] , \qquad (16)$$

including only linear matching effects to the dimension-six $(V - A) \otimes (V - A)$ operator $[\bar{c}\gamma^{\mu}(1-\gamma_5)b] [\bar{l}\gamma_{\mu}(1-\gamma_5)\nu]$ in the weak effective theory (WET) (which holds up to $\mathcal{O}(10\%)$ corrections to the BSM contribution).⁶ On the other hand, for a two-body decay into a given quark-lepton pair, the branching ratio (BR) for a particular vector LQ decay channel is given by (see e.g. [68])

$$BR (\Delta \to QL) \simeq \frac{|\lambda_{QL}|^2}{\sum_{\{QL\}} |\lambda_{QL}|^2} \quad \text{where} \quad m_{Q,L} \to 0.$$
(17)

Neglecting quark and lepton masses is an excellent approximation for the LQ mass scales we consider.

Comparing (16) to (17), one sees that the former constrains a product of LQ couplings, whereas the latter constrains a ratio. Hence, in the event of a discovery being made at the LHC, we note that combining this information would allow for a direct experimental probe at the level of individual couplings in the overall flavour matrices $\lambda_{dl,u\nu}$. We show this qualitatively in Fig. 1, given the approximations in (16)-(17), where we have presented experimentally relevant values of $\mathcal{R}_{D^{(\star)}}$ and various values of BR ($\Delta \rightarrow s\tau(b\tau)$), in the context of the τ -isolation model $\lambda_{dl}^{[\tau]}$. Note that in this model SU(2) rotations lead to non-negligible contributions from the $u - \nu$ sector, cf. Fig. 3.

In order to fully determine the parameters of the SMFL, also the LQ mass M_{Δ} and the LQ coupling parameters k_s, k_Y need to be measured. k_s and M_{Δ} can be accessed through the LQ pair-production cross-section in pp collisions and the invariant mass of the LQ decay products. The parameter k_Y is more difficult to access, as it is responsible for the coupling strength of the LQ to the photon and, to a lesser extent, to the Z boson.

3 Precision Constraints and Global Likelihoods

Our goal is to now provide a robust examination of the SMFL parameter space favored by present-day experiment, and our primary tool in this effort will be **smelli** [66], an open-source

⁶Note of course that the analysis in Section 3.2 accounts for running effects, etc. Here we are simply making a qualitative (and motivating) point.




Figure 1: A comparison of contours of $\mathcal{R}_{D^{(\star)}}$ and BR $(\Delta \to s\tau(b\tau))$ in black and red (blue), respectively.

python package which builds upon the flavio [87] and Wilson [88] programs. The goal of flavio is to enable the automated calculation of flavoured processes in terms of dimension-six Wilson coefficients C of the SM effective field theory (SMEFT, valid above the EW scale) or the Weak Effective Theory (WET, valid below the EW scale). This package includes a large library of experimental measurements in the flavour and EW sectors. In addition., Wilson computes the renormalization group evolution (RGE) of SMEFT operators, their matching onto the WET at relevant scales, and the further RGE running of WET operators to hadronic scales that are (often) of interest in flavour physics. Augmenting these capabilities, smelli computes a global likelihood function in the space of SMEFT coefficients, i.e. the object

$$L_{SMEFT}(\vec{C}) = \prod_{i} L_{exp} \left(\vec{O}_{i}^{exp}, \vec{O}_{i}^{th} \left(\vec{C}, \vec{\theta} \right) \right) \times L_{\theta} \left(\vec{\theta} \right) \,. \tag{18}$$

Here L_{exp} are likelihood distribution functions which depend on independent experimental measurements \vec{O}_i^{exp} with associated theory predictions \vec{O}_i^{th} , which are themselves functions of the Wilson coefficients \vec{C} and additional model-independent phenomenological parameters θ (e.g. form factors and decay constants). Then L_{θ} accounts for any experimental or theoretical constraints on these nuisance parameters. Note however that the actual **smelli** implementation of (18) relies on a 'nuisance-free' approximation to the total global likelihood function, which effectively 'integrates out' the theoretical errors associated to θ , treating them as additional experimental uncertainties. This is achieved by a factorization of L_{SMEFT} into likelihoods for observables that a) have negligible theoretical vs. experimental uncertainty or b) that have reliably Gaussian theory and experimental uncertainties, and where the former only weakly depends on \vec{C} and $\vec{\theta}$. While care must be taken for certain observables (e.g. CKM angles) that do not necessarily respect these assumptions, they are reliable and frequently employed (see e.g. [89–92]) for the analyses we are attempting here. Finally, smelli also includes the one-loop RGE that mix flavour structures in the SMEFT, and which are required for a consistent matching to the WET, where QCD and QED renormalization is flavourblind [93–99]. This matching and associated RGE is performed automatically in smelli, allowing coherent comparisons to low-energy experimental data given a UV new physics scale Λ . For a more complete description of smelli functionality, its built-in assumptions, and an exhaustive list of observables included in its likelihoods, see [66] and the documentation at https://github.com/smelli/smelli.

3.1 Matching the Vector Singlet

In order to perform a smelli analysis for our SMFL, we first recall the tree-level SMEFT matching onto the LL operators of the Δ^{μ} vector singlet given in (7), which when computed at the new physics scale $\Lambda = M_{\Delta}$ yields (see e.g. [100]):

$$\left[C_{LQ}^{(1)}\right]_{ijkl} = \left[C_{LQ}^{(3)}\right]_{ijkl} = -\frac{\lambda_{LQ}^{kj} \lambda_{LQ}^{li\star}}{2M_{\Delta}^2},\tag{19}$$

which are the Wilson coefficients of the dim-6 four-fermion SMEFT operators given by

$$\left[\mathcal{O}_{LQ}^{(1)}\right]_{ijkl} = \left(\overline{L}_i\gamma_{\mu}L_j\right)\left(\overline{Q}_k\gamma_{\mu}Q_l\right), \qquad \left[\mathcal{O}_{LQ}^{(3)}\right]_{ijkl} = \left(\overline{L}_i\gamma_{\mu}\tau^I L_j\right)\left(\overline{Q}_k\gamma_{\mu}\tau^I Q_l\right). \tag{20}$$

In addition to this tree-level matching, we follow [37] and include additional one-loop matching contributions to quark dipole operators in the SMEFT, which can generate the known [101] vector singlet matching contributions to electric and chromomagnetic dipole operators in the low-energy WET. The relevant dimension-six SMEFT operators are

$$[\mathcal{O}_{dB}]_{ij} = \left(\overline{Q}_i \sigma^{\mu\nu} d_j\right) \phi B_{\mu\nu}, \quad [\mathcal{O}_{dW}]_{ij} = \left(\overline{Q}_i \sigma^{\mu\nu} d_j\right) \tau^I \phi W^I_{\mu\nu}, \quad [\mathcal{O}_{dG}]_{ij} = \left(\overline{Q}_i \sigma^{\mu\nu} T^A d_j\right) \phi G^A_{\mu\nu},$$

which are catalogued alongside all remaining independent dimension-six operators in [102]. Note that in **smelli** the Warsaw basis of [102] (the **flavio** basis of [87]) is the default basis when obtaining likelihoods in the SMEFT (WET). Again computing the relevant matching at $\Lambda = M_{\Delta}$, one finds [37]

$$[C_{dW}]_{23} = \frac{Y_b}{6} \frac{g}{16\pi^2} \frac{\lambda_{LQ}^{2i} \lambda_{LQ}^{3i\star}}{M_{\Delta}^2}, \quad [C_{dB}]_{23} = -\frac{4Y_b}{9} \frac{g'}{16\pi^2} \frac{\lambda_{LQ}^{2i} \lambda_{LQ}^{3i\star}}{M_{\Delta}^2}, \quad [C_{dG}]_{23} = -\frac{5Y_b}{12} \frac{g_s}{16\pi^2} \frac{\lambda_{LQ}^{2i} \lambda_{LQ}^{3i\star}}{M_{\Delta}^2}$$

$$[C_{dW}]_{32} = \frac{Y_s}{6} \frac{g}{16\pi^2} \frac{\lambda_{LQ}^{3i} \lambda_{LQ}^{2i\star}}{M_{\Delta}^2}, \quad [C_{dB}]_{32} = -\frac{4Y_s}{9} \frac{g'}{16\pi^2} \frac{\lambda_{LQ}^{3i} \lambda_{LQ}^{2i\star}}{M_{\Delta}^2}, \quad [C_{dG}]_{32} = -\frac{5Y_s}{12} \frac{g_s}{16\pi^2} \frac{\lambda_{LQ}^{3i} \lambda_{LQ}^{2i\star}}{M_{\Delta}^2}$$

$$(21)$$

Here the lepton index i is summed over. In addition to the Lagrangian conventions discussed above Section 2.2, these were also computed in the limit of a diagonal down-quark Yukawa matrix, with $Y_{s,b}$ the respective Yukawa couplings to strange and bottom quarks. Note that,

SMFL	M_{Δ}	Best Fit $(\lambda_{sl}, \lambda_{bl})$	$\ln \Delta L _{\mathcal{R}_{D^{(\star)}}}$	$\ln \Delta L _{\mathcal{R}_{K^{(\star)}}}$	$\ln \Delta L _{Global}$
$\lambda_{dl}^{[au]}$	2 TeV	(0.68, 0.72)	12.694	N.A.	20.771
\wedge_{dl}	1 TeV	(0.32, 0.38)	12.692	IN.A.	20.075
$\lambda_{dl}^{[\mu]}$	2 TeV	(0.15, -0.024)	0.125	3.664	21.643
λ_{dl}	1 TeV	(0.06, -0.015)	0.124	3.629	21.604
$\lambda_{dl}^{[e]}$	2 TeV	(0.016, 0.184)	$-\mathcal{O}(10^{-2})$	4.866	5.827
\wedge_{dl}	1 TeV	(0.006, 0.12)	$-\mathcal{O}(10^{-2})$	4.872	5.814

Table 1: Results from the $M_{\Delta} = \{1, 2\}$ TeV two-parameter likelihood (ln $\Delta L = -\Delta \chi^2/2$) scans of Section 3.2, including the individual contributions of the (potentially) anomalous observables $\mathcal{R}_{K^{(\star)},D^{(\star)}}$. Column 3 gives the best-fit values of $(\lambda_{sl}, \lambda_{bl})$, corresponding to the global minimum of the scans, found in column 6, which consider all available data in smelli. Columns 4-5 then give the individual contributions of $\mathcal{R}_{K^{(\star)},D^{(\star)}}$ to this likelihood (again at the best-fit coordinates). See the text and Figure 2 for more details.

as mentioned above, one expects logarithmic divergences to appear in dipole processes in the 'minimal' coupling scenario where $k_s = k_Y = 1$ in (4) [24,67], and therefore the operators in (21) may not be induced in a sensible UV matching with this parameter choice. It is also clear that, when $k_s = k_Y = 1$, triple-vector couplings between $\Delta - \Delta - B/G$ are not present at tree level, and therefore the one-loop diagram leading to non-zero C_{dG} (e.g.) is not present. Regardless, we note that our analysis in Section 4 is largely insensitive to these UV details, as we have found that turning off all of the dipole operators in (21) only results in a roughly 1% correction to the best-fit ratio of $\lambda_{s\tau}/\lambda_{b\tau}$ when $M_{\Delta} = 1$ TeV (e.g.). In what follows we therefore show fit results in the 'full' $k_s = k_Y = 0$ scenario.

Finally, we recall that as with other similar studies, we have chosen to set the RR couplings x^{RR} and $x^{\overline{RR}}$ in (7) to zero. As can be deduced from model-independent EFT fits (see e.g. [37, 103–106]) and as will be seen explicitly below, these RH couplings are not necessary in minimal explanations of the observed LFU anomalies, and it has been further shown [58] that exclusion limits on M_{Δ} strengthen when $x^{RR} \neq 0$. Hence (19)-(21) represent the complete set of relevant SMEFT Wilson coefficients implemented in our smelli analysis.

3.2 Scanning the Vector Leptoquark Singlet

Given (19)-(21), one is in a position to scan over the SMFL couplings λ_{dl} that compose them and allow **smelli** to compute likelihoods at each phase-space point. One can collect this information as a function of λ_{dl} and determine the experimentally favoured space of couplings for a given SMFL pattern/model, as well as the observables which contribute the most significant pulls.

Specifically, in performing our scans we

- 1. take the leptoquark couplings λ_{dl} to be positive and real. As will be seen, there is ample parameter space of interest even without additional complex degrees of freedom.
- 2. build an array of $\{\lambda_{sl}, \lambda_{bl}\}$ by dividing the phase space in either dimension by a predefined set of intervals. We then calculate the Wilson coefficients in (19)-(21) for all points on this grid.
- 3. perform a global likelihood analysis using smelli v2.0.0 at each parameter point on the λ_{dl} grid. This is performed by calling smelli.GlobalLikelihood(), which we also modify using the custom_likelihoods attribute. This latter functionality allows us to define custom sets of observables contributing to a likelihood computation.
- 4. collect all likelihoods computed and determine the values associated to the minimum χ^2 for a given pattern (and a given set of observables). We do so by calling the log_likelihood_global() method, which returns $\ln \Delta L = -\Delta \chi^2/2$, the BSM χ^2 minus its SM value. We use this to then compute 1σ and 2σ likelihood contours about the minimum.
- 5. in order to determine the relative pull of any given set of observables, we also use the $log_likelihood_dict()$ method, which returns the dictionary of all contributions to $ln \Delta L$ from the individual products in (the smelli implementation of) (18). Note that in addition to the classes of observables already segregated in smelli through the inclusion of separate internal YAML files, any of the custom_likelihoods we defined ourselves will also be given as independent contributions.

We now report the results of these scans for SMFL patterns of distinct phenomenological interest: the lepton isolation patterns $\lambda_{dl}^{[e,\mu,\tau]}$. We also report the relative pull of $\mathcal{R}_{K^{(*)},D^{(*)}}$ (at the global minimum), which are especially interesting due to their present deviations from SM predictions.

Electron and Muon Isolation Patterns

We first investigate $\lambda_{dl}^{[e,\mu]}$ from (12), the e- and μ -isolation patterns, where the results of the **smelli** scans we performed as described above are given in the top (middle) panels of Figure 2 for the electron (muon) patterns, as well as numerically in Table 1. One observes in Figure 2 that a broad range of couplings is allowed at the 1σ and 2σ confidence level, considering all available data, and that the potentially anomalous measurement of $\mathcal{R}_{K^{(\star)}}$ is well-described in these models. Indeed, the overall shape of the global parameter space preferred largely follows that preferred by $\mathcal{R}_{K^{(\star)}}$ alone. The stronger pull to non-zero couplings in the moun-isolation pattern originates in the so-called $b \to s\mu^+\mu^-$ anomalies observed in the decays $B \to K^*\mu^+\mu^-$, $B_s \to \phi\mu^+\mu^-$ etc. that can not be addressed in the electron-isolation pattern. Additionally, as seen in Table 1, the global best-fit value is also favored over SM couplings alone as it slightly softens the charged-current LFU anomaly $\mathcal{R}_{D^{(\star)}}$, at least for the μ -isolation pattern. When considering only $\mathcal{R}_{D^{(\star)}}$ data, better likelihoods can be obtained in the parameter space scanned. $\lambda_{dl}^{[\mu]}$'s ability to resolve LFU anomalies whilst generating a distinct collider phenomenology has been known for some time (see e.g. [107, 108]).




Figure 2: **Top Left:** Likelihood contours for the two-parameter *e*-isolation pattern $\lambda_{dl}^{[e]}$ at $M_{\Delta} = 1$ TeV, including 2σ contours from $\mathcal{R}_{K^{(\star)}}$ constraints alone (hatched region), as well as global 1σ (orange region) and 2σ (blue region) preferred contours considering all data available to **smelli**. **Top Right:** The same, but for $M_{\Delta} = 2$ TeV. **Middle Row:** The same, but for the μ -isolation pattern $\lambda_{dl}^{[\mu]}$. **Bottom Row:** The same, but for the τ -isolation pattern $\lambda_{dl}^{[\tau]}$, and including 2σ contours from $\mathcal{R}_{D^{(\star)}}$ constraints alone (hatched region), and $\mathcal{R}_{D^{(\star)}} + \tau$ decay modes (gray region). Here the global best-fit (black, triangular) and benchmark (red, circular) points in $\{\lambda_{s\tau}, \lambda_{b\tau}\}$ we obtained from our scan are also shown.

As a final note, we have observed that the overall $-\Delta\chi^2/2$ likelihood distributions for both $\lambda_{dl}^{[e,\mu]}$ are somewhat flat, in that multiple points spanning a broad domain in the twodimensional contours presented fall very close to likelihood found for the global best-fit point. For example, we can identify a number of parameter space points whose likelihoods are within a percent (or less) of the global minimum, but whose coordinates are a factor of 2 (or more) away from those presented in Table 1. We found a similar behavior for $\lambda_{dl}^{[e]}$ at $M_{\Delta} = 1$ TeV, and for both $\lambda_{dl}^{[e,\mu]}$ at $M_{\Delta} = 2$ TeV. It is for this reason that we do not find it instructive to plot a 'global' best-fit point in Figure 2, but instead give this information in Table 1, to illustrate the overall quality of the fits.

Tau Isolation Pattern

We next investigate $\lambda_{dl}^{[\tau]}$ from (12), the τ -isolation pattern. This texture has been studied before in the context of a toy vector singlet LQ model [37] (although without the SMFL symmetry-based motivation for its flavour structure), and here we update those results given improved experimental and theoretical developments over the last years.⁷

Using the algorithm described in Section 3, we produce the bottom panels in Figure 2. The graphics illustrate the contours contributing to the global likelihood coming from the ratio observables $\mathcal{R}_{D^{(\star)}}$ (hatched region), those from a custom fit combining both $\mathcal{R}_{D^{(\star)}}$ and the leptonic τ -decay modes BR ($\tau^- \rightarrow l^- \nu \bar{\nu}$) with $l = e, \mu$, BR ($\tau^+ \rightarrow K^+ \bar{\nu}$), and BR ($\tau^+ \rightarrow \pi^+ \bar{\nu}$) (gray region), and finally the global 1 σ (orange region) and 2 σ (blue region) preferred contours upon considering all experimental datasets in smelli. Note that leptonic τ decays and $\mathcal{R}_{D^{(\star)}}$ were identified as dominant contributors to the overall likelihoods for $\lambda_{dl}^{[\tau]}$ in [37], and we confirm that observation in our analysis. We give these at $M_{\Delta} = \{1,2\}$ TeV (left and right panels, respectively), and the best-fit ($\lambda_{s\tau}, \lambda_{b\tau}$) values are shown in black, where the global minimum likelihood of our scans are realized. The numerical value of this coordinate as well as the minimum $-\Delta\chi^2/2$ is again given in Table 1. For the 2 TeV contours we also plot 'benchmark points' in red that we will use in the upcoming collider analysis of Section 4.

Summary

In conclusion, it is clear that the μ - and the τ -isolation patterns both provide excellent fits to the available data across a broad range of parameter space, and are able to solve the LFU anomalies $\mathcal{R}_{K^{(*)}}$ and $\mathcal{R}_{D^{(*)}}$, respectively. While the electron isolation pattern can also successfully address $\mathcal{R}_{K^{(*)}}$, it falls short of explaining the related anomalies in $b \to s\mu^+\mu^$ transitions, and is hence less motivated from a phenomenological perspective.

In what follows we will further pursue the analysis of the τ -isolation pattern and study its LHC signatures. A high- p_T study of this scenario is particularly motivated, since we have seen large LQ couplings are required by the global fit. In turn this precludes the possibility of evading direct searches by simply raising the LQ mass M_{Δ} beyond the reach of the LHC.

⁷We thank Peter Stangl for pointing out (e.g.) [109], whose updated $B \to D^{(\star)}$ form factors impact predictions (and uncertainty estimates) for $\mathcal{R}_{D^{(\star)}}$. Note also that the gray-shaded contour of the bottom-right panel of Figure 2 can be compared to the corresponding contours in Figure 6 of [37], where we find good qualitative agreement, given the updates in the code and both experimental and theoretical inputs.

$\lambda_{dl}^{[\tau]}$ Benchmark Points								
BP_i	$(\lambda_{s au},\lambda_{b au})$							
BP_1	(1.29, 0.4)							
BP_2	(0.98, 0.49)							
BP_3	(0.68, 0.68)							
BP_4	(0.48, 0.96)							
BP_5	(0.32, 1.43)							

$BR(\Delta \rightarrow QL) \ [\%]$										
_	$\mathrm{b} au$	tν	s $ au$	cv						
BP_1	4.4	3.3	45.6	44.3		40				
BP_2	10.0	8.4	40.0	39.5		-30				
BP_3	25.1	22.9	25.1	25.7						
BP_4	40.2	38.1	10.0	11.1		20				
BP ₅	47.8	46.4	2.4	3.2		10				

 $\text{BR}(\Delta \rightarrow \text{QLQL}) \ [\%]$

	bτbτ	tvtv	bτtv	bτsτ	bτcv	tνsτ	tvcv	s $ au$ s $ au$ s $ au$	sτcν	сиси		
BP_1	0.2	0.1	0.3	4.0	3.9	3.0	3.0	20.8	40.5	19.7	-	40
BP_2	1.0	0.7	1.7	8.0	7.9	6.7	6.6	16.0	31.6	15.6		30
BP_3	6.3	5.2	11.5	12.6	12.9	11.5	11.8	6.3	12.9	6.6	-	20
BP_4	16.1	14.5	30.6	8.1	8.9	7.7	8.5	1.0	2.2	1.2		10
BP_5	22.9	21.5	44.4	2.3	3.1	2.3	3.0	0.1	0.2	0.1		-

Table 2: $BR(\Delta(\Delta) \to QL(QL))$ for 2 TeV LQ Benchmark Points from the $\lambda_{dl}^{[\tau]}$ scan of Section 3. All asymmetric decay BR have an additional $\times 2$ factor because of permutation.

4 LHC Recasting Exclusion Limits

In this Section we consider existing direct searches published by ATLAS [110] and reinterpret them in the context of our well motivated τ —isolation scenario. In particular, both second- and third-generation quarks can be involved in the LQ decay. Assuming only left-handed couplings, and neglecting the fermion masses as well as CKM rotations, the different branching ratios exhibit the following structure

$$BR_{\Delta}^{3^{rd}} = BR(\Delta \to b\tau) \simeq BR(\Delta \to t\nu), \qquad (22)$$




Figure 3: 2 TeV LQ branching ratios into quarks and leptons for the τ - isolation pattern. The channel magnitudes only depend on the ratio of the couplings. The non-vanishing $u - \nu$ contribution appears due to CKM rotations from $\lambda_{d\ell} \rightarrow \lambda_{u\nu}$. The other channels are (almost) exactly vanishing in the context of the τ -isolation pattern. We have highlighted in red the position of the different benchmark points shown in Figure 2. Note that P_2^{τ} , P_3^{τ} and P_4^{τ} have $\lambda_{b\tau}/\lambda_{s\tau}$ values of 0.5, 1 and 2, respectively.

$$BR_{\Delta}^{2^{nd}} = BR(\Delta \to s\tau) \simeq BR(\Delta \to c\nu) \simeq 0.5 - 2 \cdot BR_{\Delta}^{3^{rd}}, \qquad (23)$$

where $BR_{\Delta}^{2^{nd}}$ ($BR_{\Delta}^{3^{rd}}$) corresponds to the decay into the second (third) down-type quark. This structure holds very well as shown in Figure 3 (note that for readability we fixed the only coupling free parameters $\lambda_{dl} \doteq \lambda$) where the CKM and fermion masses effects have been considered with a 2 TeV mass for the LQ. Using (17) we obtain the following relation for the branching ratio as a function of the couplings:

$$BR_{\Delta}^{3^{rd}} \simeq \frac{\lambda_{33}^2}{\lambda_{33}^2 + \lambda_{23}^2}.$$
(24)

As an example, we selected five benchmark points labelled P_i^{τ} across the 1σ region fit of Figure 2. These scenarios have distinct decay channel magnitudes and are therefore rather illustrative for collider considerations. All information regarding these benchmark points is gathered in Table 2.

The procedure regarding our collider analysis is as follows: we first consider the mixed search $\Delta \rightarrow b\tau/t\nu$ investigated in [110] where we adapt to match the extra opened secondgeneration quark channel. We then complete the analysis by confronting our model with the implemented LHC searches in CheckMATE [111,112]. Of particular interest would be the jets and missing energy searches, since the channel involving jets and neutrino remains stable around 50% as discussed in (23). We present results for the two extreme cases for pair production: k = 0 and k = 1.




Figure 4: Pair production cross-section for different LQ models. For the vector LQ Δ we present results for both the k = 0 and k = 1 scenarios studied in the text. The scalar model (S_1) is presented as an illustration, being the benchmark model employed in most of the ATLAS and CMS studies.

4.1 Reinterpretation of the Mixed $b\tau t\nu$ Search

Several searches at the LHC by the ATLAS [110, 113] and CMS [114–116] collaborations explicitly target up-type vector leptoquarks with $b\tau$ and $t\nu$ final states. Among those, ATLAS in [110] (CMS in [116]) considers for the first time the possibility of a $b\tau t\nu$ final state (originally pointed out in [117] and scrutinized in [118,119]), where each leptoquark decayed in a different channel. We informally refer to this as the "mixed" channel. Moreover, for the specific case where BR($\Delta \rightarrow b\tau$) and BR($\Delta \rightarrow t\nu$) are approximately equal, the study of reference [110] provides the most stringent constraints, and hence, among the whole suite of leptoquark studies, we focus on this particular channel.

The ATLAS study [110] considers LQ decays exclusively to third-generation quarks. In other words, the branching ratios into $t\nu$ and $b\tau$ add up to unity. However, in more generic setups additional decay channels can be opened. Following the strategy proposed in [119] we reinterpret the search in this more generic framework, for our vector LQ model.

As discussed above, within our assumptions, we can use (23). Therefore, the excluded fiducial cross-section given by ATLAS is simply re-expressed as

$$\sigma(pp \to \Delta\Delta^*) \times BR(\Delta \to b\tau) \times BR(\Delta \to t\nu) \Longrightarrow \sigma(pp \to \Delta\Delta^*) \times \left(BR_{\Delta}^{3^{rd}}\right)^2, \qquad (25)$$

which corresponds to $BR(\Delta \rightarrow b\tau) = 0.5$ in the ATLAS paper.

Using our results for the production cross-section as a function of the mass, given in Figure 4, it is possible to reproduce the exclusion given by ATLAS while including the effect of the





Figure 5: Reinterpretation from [110]. The main assumptions are $BR(\Delta \to b\tau) = BR(\Delta \to t\nu)$ and that the opened second generation decay is invisible to the analysis, and hence only affects the fiducial cross-section $\sigma(pp \to \Delta\Delta^*) \times (BR_{\Delta}^{3rd})^2$

additional opened channels by varying $BR_{\Delta}^{3^{rd}}$ instead, which can be matched to the ratio $\lambda_{33}/\lambda_{23}$. The result is presented in Figure 5. We observe that, as expected, the excluded mass drops significantly as the branching ratio into the third generation quarks decreases. The two extreme cases from the best fits presented in Section 3.2 are reached for BP1 and BP5, for which we exclude masses close to 600 GeV (900 GeV) and 1500 GeV (1750 GeV) for k = 1 (k = 0), respectively.

4.2 CheckMATE Exclusion Limits

As discussed previously, we expect searches involving multi-jets and missing energy to complete the exclusion limits provided by the reinterpretation of the mixed channel. Indeed, at parton level, the channel involving neutrino decay of the leptoquark $(t\nu + c\nu)$ remains stable around 50% across the coupling parameter space as only the left-handed coupling is present. Decay channels involving either one or two neutrinos will therefore be probed by these searches.

We perform a parameter scan in the plane $(M, \lambda_{33}/\lambda_{23})$ as only the coupling ratio has an impact on the collider phenomenology. Using **CheckMATE** and previously implemented analysis we derive the complementary exclusion limits. The results are shown in Fig. 6 together with the mixed search, adapted to the $(M, \lambda_{33}/\lambda_{23})$ plane using the valid simplified expression for the branching ratios (24). We note that the limits seem rather independent of the coupling ratio with the excluded values for k = 0 (k = 1) below ~ 1500 GeV (~ 1100 GeV), reproducing our expectations from the parton-level consideration where the neutrino channels sum up to 50% independently of the coupling ratio. This further suggests that at the reconstructed level, where in principle the nature of the up-type quark (c or t) might affect the jet multiplicity,



the different signal regions have similar sensitivity. Most of the exclusion limits come from the ATLAS analysis [120] where they originally considered squark and gluino multi-jets and missing energy with 36 fb⁻¹. Note that a new analysis by ATLAS of the same channels was released in [121] where they performed a more sophisticated search with more luminosity. However, recasting this analysis is beyond the scope of this work as this search has not been implemented and validated in **CheckMATE**. Being conservative by assuming that the efficiency of future searches will remain stable and therefore simply rescale by the luminosity, one can estimate prospects for HL-LHC for 3000 fb⁻¹, which should exclude masses of order 1400 GeV (1850 GeV) for k = 1 (k = 0). However, due to the improvement of the analysis techniques (for example in [121] compared to [120]), one can be more optimistic and expect even higher masses to be reached. We then see that both classes of studies (searches for leptoquarks and missing energy searches) complement each other and that enhancing the sensitivity of these studies is of crucial importance towards shedding light on the mechanism(s) behind the flavour anomalies.

5 Summary and Outlook

We have studied the phenomenological consequences of a subset of Simplified Models of Flavourful Leptoquarks [34–36] that only couple to a single generation of charged leptons, with a central focus on the $(\mathbf{3}, \mathbf{1}, 2/3)$ lepqtoquark Δ_{μ} coupling quarks to τ leptons. Lepton isolation patterns/models are readily motivated by ultraviolet flavour-symmetry breaking, and are amongst the most minimal scenarios that can explain experimental anomalies in the leptonflavour-universality observables $\mathcal{R}_{D^{(\star)},K^{(\star)}}$, depending on the actual tree-level lepton couplings allowed. We reviewed their theoretical motivation and used smelli [66] to scan over the global parameter space favoured by low-energy flavour data when couplings to electrons, muons, or taus are permitted. This analysis revealed that all three patterns are favored over SM physics alone, and we presented 1σ and 2σ global likelihood contours in the two-parameter spaces of (12), along with likelihood contours in the same space, but coming from $\mathcal{R}_{D^{(\star)},K^{(\star)}}$ data alone. Upon focusing on the τ -isolation pattern, we then studied the decay signatures of Δ_{μ} at the LHC.

Based on our parameter-space scans, we have defined a series of benchmark points to illustrate the main phenomenological features, where the branching fraction into the third generation ranges from a few percent up to almost 100 %. We have confronted the parameter space of our model with two different classes of LHC searches, namely i) direct searches for leptoquarks and ii) missing energy searches. Given that our setup imposes approximately equal branching ratios into $t\nu$ and $b\tau$, it turns out that the mixed ATLAS search looking for $b\tau t\nu$ provides the strongest constraints among the set of direct leptoquark searches. We carried out the reinterpretation of this search for our vector leptoquark models, where the sensitivity would obviously depend on the branching ratio into the third generation. If no additional channels are open, these searches can constrain a leptoquark mass up to 1400 (1750) GeV for the k = 1 (k = 0) scenarios, while if instead the branching fraction into the third generation were to be of 5 %, these upper limits would be reduced to 800 (1100) GeV. Regarding the missing energy searches, since our setup also imposes that our leptoquark decays 50 % of the time in a channel with one neutrino, it is not surprising that the mass reach is pretty insensitive to the third generation branching fraction, being about 1150 (1500) GeV for the k = 1 (k = 0) case. The interplay between both types of searches is interesting.

Last but not least, we would like to point out that it would be desirable to expand the set of direct leptoquark searches to specifically target decays into the second generation. While experimentally challenging, it is clear that since the flavour anomalies are directly related to the second generation, searches including c quarks, e.g. $c\nu c\nu$ or $b\tau c\nu$ would have an important impact on both the discovery and the characterization of leptoquark models. While outside the scope of this work, it would also be desirable to study in similar fashion how future colliders can fully probe the vector leptoquark scenario.

Acknowledgements

We thank Peter Stangl and Jason Aebischer for very helpful discussions regarding smelli. JB and MB are supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762 – TRR 257. IdMV acknowledges funding from Fundação para a Ciência e a Tecnologia (FCT, Portugal) through the contract IF/00816/2015 and through the projects CFTP-FCT Unit 777 (UIDB/00777/2020 and UIDP/00777/2020), PTDC/FIS-PAR/29436/2017 and CERN/FIS-PAR/0008/2019, which are partially funded through POCTI (FEDER), COMPETE, QREN and EU. JT gratefully acknowledges funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 101022203, as well as prior support from the Villum Fund (project No. 00010102). JZ is supported by the Generalitat Valenciana (Spain) through

the *plan GenT* program (CIDEGENT/2019/068), by the Spanish Government (Agencia Estatal de Investigación) and ERDF funds from European Commission (MCIN/AEI/10.13039/501100011033, Grant No. PID2020-114473GB-I00).

References

- [1] J. P. Lees *et al.* [BaBar], Phys. Rev. Lett. **109** (2012), 101802 doi:10.1103/PhysRevLett.109.101802 [arXiv:1205.5442 [hep-ex]].
- [2] J. P. Lees *et al.* [BaBar], Phys. Rev. D 88 (2013) no.7, 072012 doi:10.1103/PhysRevD.88.072012 [arXiv:1303.0571 [hep-ex]].
- [3] M. Huschle *et al.* [Belle], Phys. Rev. D **92** (2015) no.7, 072014 doi:10.1103/PhysRevD.92.072014 [arXiv:1507.03233 [hep-ex]].
- [4] S. Hirose *et al.* [Belle], Phys. Rev. Lett. **118** (2017) no.21, 211801 doi:10.1103/PhysRevLett.118.211801 [arXiv:1612.00529 [hep-ex]].
- [5] Y. Sato *et al.* [Belle], Phys. Rev. D **94** (2016) no.7, 072007 doi:10.1103/PhysRevD.94.072007 [arXiv:1607.07923 [hep-ex]].
- [6] S. Hirose *et al.* [Belle], Phys. Rev. Lett. **118** (2017) no.21, 211801 doi:10.1103/PhysRevLett.118.211801 [arXiv:1612.00529 [hep-ex]].
- [7] A. Abdesselam *et al.* [Belle], [arXiv:1904.08794 [hep-ex]].
- [8] R. Aaij et al. [LHCb], Phys. Rev. Lett. 115 (2015) no.11, 111803 [erratum: Phys. Rev. Lett. 115 (2015) no.15, 159901] doi:10.1103/PhysRevLett.115.111803 [arXiv:1506.08614 [hep-ex]].
- [9] R. Aaij *et al.* [LHCb], Phys. Rev. Lett. **120** (2018) no.17, 171802 doi:10.1103/PhysRevLett.120.171802 [arXiv:1708.08856 [hep-ex]].
- [10] R. Aaij et al. [LHCb], Phys. Rev. D 97 (2018) no.7, 072013 doi:10.1103/PhysRevD.97.072013 [arXiv:1711.02505 [hep-ex]].
- [11] Y. S. Amhis *et al.* [HFLAV], Eur. Phys. J. C 81 (2021) no.3, 226 doi:10.1140/epjc/s10052-020-8156-7 [arXiv:1909.12524 [hep-ex]].
- [12] A. Crivellin, C. Greub and A. Kokulu, Phys. Rev. D 86 (2012), 054014 doi:10.1103/PhysRevD.86.054014 [arXiv:1206.2634 [hep-ph]].
- [13] A. Crivellin, A. Kokulu and C. Greub, Phys. Rev. D 87 (2013) no.9, 094031 doi:10.1103/PhysRevD.87.094031 [arXiv:1303.5877 [hep-ph]].
- [14] A. Celis, M. Jung, X. Q. Li and A. Pich, JHEP **01** (2013), 054 doi:10.1007/JHEP01(2013)054 [arXiv:1210.8443 [hep-ph]].

- [15] P. Ko, Y. Omura and C. Yu, JHEP 03 (2013), 151 doi:10.1007/JHEP03(2013)151 [arXiv:1212.4607 [hep-ph]].
- [16] A. Crivellin, J. Heeck and P. Stoffer, Phys. Rev. Lett. **116** (2016) no.8, 081801 doi:10.1103/PhysRevLett.116.081801 [arXiv:1507.07567 [hep-ph]].
- [17] X. G. He and G. Valencia, Phys. Rev. D 87 (2013) no.1, 014014 doi:10.1103/PhysRevD.87.014014 [arXiv:1211.0348 [hep-ph]].
- [18] A. Greljo, G. Isidori and D. Marzocca, JHEP 07 (2015), 142 doi:10.1007/JHEP07(2015)142 [arXiv:1506.01705 [hep-ph]].
- [19] S. M. Boucenna, A. Celis, J. Fuentes-Martin, A. Vicente and J. Virto, Phys. Lett. B 760 (2016), 214-219 doi:10.1016/j.physletb.2016.06.067 [arXiv:1604.03088 [hep-ph]].
- [20] X. G. He and G. Valencia, Phys. Lett. B 779 (2018), 52-57 doi:10.1016/j.physletb.2018.01.073 [arXiv:1711.09525 [hep-ph]].
- [21] R. Alonso, B. Grinstein and J. Martin Camalich, JHEP 10 (2015), 184 doi:10.1007/JHEP10(2015)184 [arXiv:1505.05164 [hep-ph]].
- [22] L. Calibbi, A. Crivellin and T. Ota, Phys. Rev. Lett. 115 (2015), 181801 doi:10.1103/PhysRevLett.115.181801 [arXiv:1506.02661 [hep-ph]].
- [23] S. Fajfer and N. Košnik, Phys. Lett. B 755 (2016), 270-274 doi:10.1016/j.physletb.2016.02.018 [arXiv:1511.06024 [hep-ph]].
- [24] R. Barbieri, G. Isidori, A. Pattori and F. Senia, Eur. Phys. J. C 76 (2016) no.2, 67 doi:10.1140/epjc/s10052-016-3905-3 [arXiv:1512.01560 [hep-ph]].
- [25] R. Barbieri, C. W. Murphy and F. Senia, Eur. Phys. J. C 77 (2017) no.1, 8 doi:10.1140/epjc/s10052-016-4578-7 [arXiv:1611.04930 [hep-ph]].
- [26] G. Hiller, D. Loose and K. Schönwald, JHEP **12** (2016), 027 doi:10.1007/JHEP12(2016)027 [arXiv:1609.08895 [hep-ph]].
- [27] N. G. Deshpande and A. Menon, JHEP 01 (2013), 025 doi:10.1007/JHEP01(2013)025 [arXiv:1208.4134 [hep-ph]].
- [28] M. Tanaka and R. Watanabe, Phys. Rev. D 87 (2013) no.3, 034028 doi:10.1103/PhysRevD.87.034028 [arXiv:1212.1878 [hep-ph]].
- [29] Y. Sakaki, M. Tanaka, A. Tayduganov and R. Watanabe, Phys. Rev. D 88 (2013) no.9, 094012 doi:10.1103/PhysRevD.88.094012 [arXiv:1309.0301 [hep-ph]].
- [30] M. Freytsis, Z. Ligeti and J. T. Ruderman, Phys. Rev. D 92 (2015) no.5, 054018 doi:10.1103/PhysRevD.92.054018 [arXiv:1506.08896 [hep-ph]].
- [31] M. Bauer and M. Neubert, Phys. Rev. Lett. **116** (2016) no.14, 141802 doi:10.1103/PhysRevLett.116.141802 [arXiv:1511.01900 [hep-ph]].

- [32] D. Bečirević, S. Fajfer, N. Košnik and O. Sumensari, Phys. Rev. D 94 (2016) no.11, 115021 doi:10.1103/PhysRevD.94.115021 [arXiv:1608.08501 [hep-ph]].
- [33] D. Bečirević, I. Doršner, S. Fajfer, N. Košnik, D. A. Faroughy and O. Sumensari, Phys. Rev. D 98 (2018) no.5, 055003 doi:10.1103/PhysRevD.98.055003 [arXiv:1806.05689 [hep-ph]].
- [34] I. de Medeiros Varzielas and J. Talbert, Eur. Phys. J. C 79 (2019) no.6, 536 doi:10.1140/epjc/s10052-019-7047-2 [arXiv:1901.10484 [hep-ph]].
- [35] J. Bernigaud, I. de Medeiros Varzielas and J. Talbert, JHEP 01 (2020), 194 doi:10.1007/JHEP01(2020)194 [arXiv:1906.11270 [hep-ph]].
- [36] J. Bernigaud, I. de Medeiros Varzielas and J. Talbert, Eur. Phys. J. C 81 (2021) no.1, 65 doi:10.1140/epjc/s10052-021-08882-7 [arXiv:2005.12293 [hep-ph]].
- [37] J. Aebischer, W. Altmannshofer, D. Guadagnoli, M. Reboud, P. Stangl and D. M. Straub, Eur. Phys. J. C 80 (2020) no.3, 252 doi:10.1140/epjc/s10052-020-7817-x [arXiv:1903.10434 [hep-ph]].
- [38] J. C. Pati and A. Salam, Phys. Rev. D 10 (1974), 275-289 [erratum: Phys. Rev. D 11 (1975), 703-703] doi:10.1103/PhysRevD.10.275
- [39] L. Di Luzio, A. Greljo and M. Nardecchia, Phys. Rev. D 96 (2017) no.11, 115011 doi:10.1103/PhysRevD.96.115011 [arXiv:1708.08450 [hep-ph]].
- [40] L. Calibbi, A. Crivellin and T. Li, Phys. Rev. D 98 (2018) no.11, 115002 doi:10.1103/PhysRevD.98.115002 [arXiv:1709.00692 [hep-ph]].
- [41] M. Bordone, C. Cornella, J. Fuentes-Martin and G. Isidori, Phys. Lett. B 779 (2018), 317-323 doi:10.1016/j.physletb.2018.02.011 [arXiv:1712.01368 [hep-ph]].
- [42] R. Barbieri and A. Tesi, Eur. Phys. J. C 78 (2018) no.3, 193 doi:10.1140/epjc/s10052-018-5680-9 [arXiv:1712.06844 [hep-ph]].
- [43] M. Blanke and A. Crivellin, Phys. Rev. Lett. **121** (2018) no.1, 011801 doi:10.1103/PhysRevLett.121.011801 [arXiv:1801.07256 [hep-ph]].
- [44] M. Algueró, B. Capdevila, S. Descotes-Genon, J. Matias and M. Novoa-Brunet, [arXiv:2104.08921 [hep-ph]].
- [45] W. Altmannshofer and P. Stangl, Eur. Phys. J. C 81 (2021) no.10, 952 doi:10.1140/epjc/s10052-021-09725-1 [arXiv:2103.13370 [hep-ph]].
- [46] R. Aaij et al. [LHCb], JHEP 08 (2017), 055 doi:10.1007/JHEP08(2017)055 [arXiv:1705.05802 [hep-ex]].
- [47] R. Aaij et al. [LHCb], [arXiv:2103.11769 [hep-ex]].
- [48] R. Aaij et al. [LHCb], [arXiv:2110.09501 [hep-ex]].

- [49] S. Choudhury et al. [BELLE], JHEP 03 (2021), 105 doi:10.1007/JHEP03(2021)105 [arXiv:1908.01848 [hep-ex]].
- [50] A. Abdesselam *et al.* [Belle], Phys. Rev. Lett. **126** (2021) no.16, 161801 doi:10.1103/PhysRevLett.126.161801 [arXiv:1904.02440 [hep-ex]].
- [51] B. C. Allanach, B. Gripaios and T. You, JHEP 03 (2018), 021 doi:10.1007/JHEP03(2018)021 [arXiv:1710.06363 [hep-ph]].
- [52] B. Garland, S. Jäger, C. K. Khosa and S. Kvedaraitė, [arXiv:2112.05127 [hep-ph]].
- [53] P. Asadi, R. Capdevilla, C. Cesarotti and S. Homiller, JHEP 10 (2021), 182 doi:10.1007/JHEP10(2021)182 [arXiv:2104.05720 [hep-ph]].
- [54] B. C. Allanach, T. Corbett and M. Madigan, Eur. Phys. J. C 80 (2020) no.2, 170 doi:10.1140/epjc/s10052-020-7722-3 [arXiv:1911.04455 [hep-ph]].
- [55] T. Mandal, S. Mitra and S. Seth, JHEP 07 (2015), 028 doi:10.1007/JHEP07(2015)028 [arXiv:1503.04689 [hep-ph]].
- [56] A. Bhaskar, T. Mandal, S. Mitra and M. Sharma, Phys. Rev. D 104 (2021) no.7, 075037 doi:10.1103/PhysRevD.104.075037 [arXiv:2106.07605 [hep-ph]].
- [57] U. Aydemir, T. Mandal and S. Mitra, Phys. Rev. D 101 (2020) no.1, 015011 doi:10.1103/PhysRevD.101.015011 [arXiv:1902.08108 [hep-ph]].
- [58] M. J. Baker, J. Fuentes-Martín, G. Isidori and M. König, Eur. Phys. J. C 79 (2019) no.4, 334 doi:10.1140/epjc/s10052-019-6853-x [arXiv:1901.10480 [hep-ph]].
- [59] M. Endo, S. Iguro, T. Kitahara, M. Takeuchi and R. Watanabe, [arXiv:2111.04748 [hep-ph]].
- [60] C. Cornella, D. A. Faroughy, J. Fuentes-Martin, G. Isidori and M. Neubert, JHEP 08 (2021), 050 doi:10.1007/JHEP08(2021)050 [arXiv:2103.16558 [hep-ph]].
- [61] A. Angelescu, D. Bečirević, D. A. Faroughy and O. Sumensari, JHEP 10 (2018), 183 doi:10.1007/JHEP10(2018)183 [arXiv:1808.08179 [hep-ph]].
- [62] F. Feruglio, P. Paradisi and O. Sumensari, JHEP **11** (2018), 191 doi:10.1007/JHEP11(2018)191 [arXiv:1806.10155 [hep-ph]].
- [63] A. Angelescu, D. Bečirević, D. A. Faroughy, F. Jaffredo and O. Sumensari, Phys. Rev. D 104 (2021) no.5, 055017 doi:10.1103/PhysRevD.104.055017 [arXiv:2103.12504 [hep-ph]].
- [64] A. Greljo, J. Martin Camalich and J. D. Ruiz-Álvarez, Phys. Rev. Lett. **122** (2019) no.13, 131803 doi:10.1103/PhysRevLett.122.131803 [arXiv:1811.07920 [hep-ph]].
- [65] T. Husek, K. Monsalvez-Pozo and J. Portoles, [arXiv:2111.06872 [hep-ph]].
- [66] J. Aebischer, J. Kumar, P. Stangl and D. M. Straub, Eur. Phys. J. C 79 (2019) no.6, 509 doi:10.1140/epjc/s10052-019-6977-z [arXiv:1810.07698 [hep-ph]].

- [67] S. Ferrara, M. Porrati and V. L. Telegdi, Phys. Rev. D 46 (1992), 3529-3537 doi:10.1103/PhysRevD.46.3529
- [68] I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik and N. Košnik, Phys. Rept. 641 (2016), 1-68 doi:10.1016/j.physrep.2016.06.001 [arXiv:1603.04993 [hep-ph]].
- [69] P. A. Zyla *et al.* [Particle Data Group], PTEP **2020** (2020) no.8, 083C01 doi:10.1093/ptep/ptaa104
- [70] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, JHEP 09 (2020), 178 doi:10.1007/JHEP09(2020)178 [arXiv:2007.14792 [hep-ph]].
- [71] S. F. King and C. Luhn, Rept. Prog. Phys. 76 (2013), 056201 doi:10.1088/0034-4885/76/5/056201 [arXiv:1301.1340 [hep-ph]].
- [72] C. S. Lam, Phys. Lett. B 656 (2007), 193-198 doi:10.1016/j.physletb.2007.09.032
 [arXiv:0708.3665 [hep-ph]].
- [73] D. Hernandez and A. Y. Smirnov, Phys. Rev. D 86 (2012), 053014 doi:10.1103/PhysRevD.86.053014 [arXiv:1204.0445 [hep-ph]].
- [74] C. S. Lam, Phys. Rev. D 87 (2013) no.1, 013001 doi:10.1103/PhysRevD.87.013001 [arXiv:1208.5527 [hep-ph]].
- [75] M. Holthausen, K. S. Lim and M. Lindner, Phys. Lett. B 721 (2013), 61-67 doi:10.1016/j.physletb.2013.02.047 [arXiv:1212.2411 [hep-ph]].
- [76] S. F. King, T. Neder and A. J. Stuart, Phys. Lett. B 726 (2013), 312-315 doi:10.1016/j.physletb.2013.08.052 [arXiv:1305.3200 [hep-ph]].
- [77] M. Holthausen and K. S. Lim, Phys. Rev. D 88 (2013), 033018 doi:10.1103/PhysRevD.88.033018 [arXiv:1306.4356 [hep-ph]].
- [78] L. Lavoura and P. O. Ludl, Phys. Lett. B 731 (2014), 331-336 doi:10.1016/j.physletb.2014.03.001 [arXiv:1401.5036 [hep-ph]].
- [79] R. M. Fonseca and W. Grimus, JHEP 09 (2014), 033 doi:10.1007/JHEP09(2014)033
 [arXiv:1405.3678 [hep-ph]].
- [80] A. S. Joshipura and K. M. Patel, Phys. Rev. D 90 (2014) no.3, 036005 doi:10.1103/PhysRevD.90.036005 [arXiv:1405.6106 [hep-ph]].
- [81] J. Talbert, JHEP 12 (2014), 058 doi:10.1007/JHEP12(2014)058 [arXiv:1409.7310 [hep-ph]].
- [82] C. Y. Yao and G. J. Ding, Phys. Rev. D 92 (2015) no.9, 096010 doi:10.1103/PhysRevD.92.096010 [arXiv:1505.03798 [hep-ph]].
- [83] J. N. Lu and G. J. Ding, Phys. Rev. D 95 (2017) no.1, 015012 doi:10.1103/PhysRevD.95.015012 [arXiv:1610.05682 [hep-ph]].

- [84] I. de Medeiros Varzielas, R. W. Rasmussen and J. Talbert, Int. J. Mod. Phys. A 32 (2017) no.06n07, 1750047 doi:10.1142/S0217751X17500476 [arXiv:1605.03581 [hep-ph]].
- [85] I. de Medeiros Varzielas and J. Talbert, Phys. Lett. B 800 (2020), 135091 doi:10.1016/j.physletb.2019.135091 [arXiv:1908.10979 [hep-ph]].
- [86] I. de Medeiros Varzielas and G. Hiller, JHEP 06 (2015), 072 doi:10.1007/JHEP06(2015)072 [arXiv:1503.01084 [hep-ph]].
- [87] D. M. Straub, [arXiv:1810.08132 [hep-ph]].
- [88] J. Aebischer, J. Kumar and D. M. Straub, Eur. Phys. J. C 78 (2018) no.12, 1026 doi:10.1140/epjc/s10052-018-6492-7 [arXiv:1804.05033 [hep-ph]].
- [89] A. Efrati, A. Falkowski and Y. Soreq, JHEP 07 (2015), 018 doi:10.1007/JHEP07(2015)018 [arXiv:1503.07872 [hep-ph]].
- [90] A. Falkowski, M. González-Alonso and K. Mimouni, JHEP 08 (2017), 123 doi:10.1007/JHEP08(2017)123 [arXiv:1706.03783 [hep-ph]].
- [91] W. Altmannshofer and D. M. Straub, Eur. Phys. J. C 75 (2015) no.8, 382 doi:10.1140/epjc/s10052-015-3602-7 [arXiv:1411.3161 [hep-ph]].
- [92] S. Descotes-Genon, L. Hofer, J. Matias and J. Virto, JHEP 06 (2016), 092 doi:10.1007/JHEP06(2016)092 [arXiv:1510.04239 [hep-ph]].
- [93] E. E. Jenkins, A. V. Manohar and M. Trott, JHEP 10 (2013), 087 doi:10.1007/JHEP10(2013)087 [arXiv:1308.2627 [hep-ph]].
- [94] E. E. Jenkins, A. V. Manohar and M. Trott, JHEP 01 (2014), 035 doi:10.1007/JHEP01(2014)035 [arXiv:1310.4838 [hep-ph]].
- [95] R. Alonso, E. E. Jenkins, A. V. Manohar and M. Trott, JHEP 04 (2014), 159 doi:10.1007/JHEP04(2014)159 [arXiv:1312.2014 [hep-ph]].
- [96] J. Aebischer, A. Crivellin, M. Fael and C. Greub, JHEP 05 (2016), 037 doi:10.1007/JHEP05(2016)037 [arXiv:1512.02830 [hep-ph]].
- [97] E. E. Jenkins, A. V. Manohar and P. Stoffer, JHEP 03 (2018), 016 doi:10.1007/JHEP03(2018)016 [arXiv:1709.04486 [hep-ph]].
- [98] J. Aebischer, M. Fael, C. Greub and J. Virto, JHEP 09 (2017), 158 doi:10.1007/JHEP09(2017)158 [arXiv:1704.06639 [hep-ph]].
- [99] E. E. Jenkins, A. V. Manohar and P. Stoffer, JHEP 01 (2018), 084 doi:10.1007/JHEP01(2018)084 [arXiv:1711.05270 [hep-ph]].
- [100] J. de Blas, J. C. Criado, M. Perez-Victoria and J. Santiago, JHEP 03 (2018), 109 doi:10.1007/JHEP03(2018)109 [arXiv:1711.10391 [hep-ph]].

- [101] A. Crivellin, C. Greub, D. Müller and F. Saturnino, Phys. Rev. Lett. **122** (2019) no.1, 011805 doi:10.1103/PhysRevLett.122.011805 [arXiv:1807.02068 [hep-ph]].
- [102] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, JHEP 10 (2010), 085 doi:10.1007/JHEP10(2010)085 [arXiv:1008.4884 [hep-ph]].
- [103] C. Murgui, A. Peñuelas, M. Jung and A. Pich, JHEP 09 (2019), 103 doi:10.1007/JHEP09(2019)103 [arXiv:1904.09311 [hep-ph]].
- [104] R. X. Shi, L. S. Geng, B. Grinstein, S. Jäger and J. Martin Camalich, JHEP 12 (2019), 065 doi:10.1007/JHEP12(2019)065 [arXiv:1905.08498 [hep-ph]].
- [105] M. Blanke, A. Crivellin, S. de Boer, T. Kitahara, M. Moscati, U. Nierste and I. Nišandžić, Phys. Rev. D 99 (2019) no.7, 075006 doi:10.1103/PhysRevD.99.075006 [arXiv:1811.09603 [hep-ph]].
- [106] M. Blanke, A. Crivellin, T. Kitahara, M. Moscati, U. Nierste and I. Nišandžić, doi:10.1103/PhysRevD.100.035035 [arXiv:1905.08253 [hep-ph]].
- [107] G. Hiller, D. Loose and I. Nišandžić, Phys. Rev. D 97 (2018) no.7, 075004 doi:10.1103/PhysRevD.97.075004 [arXiv:1801.09399 [hep-ph]].
- [108] G. Hiller, D. Loose and I. Nišandžić, JHEP 06 (2021), 080 doi:10.1007/JHEP06(2021)080 [arXiv:2103.12724 [hep-ph]].
- [109] M. Bordone, M. Jung and D. van Dyk, Eur. Phys. J. C 80 (2020) no.2, 74 doi:10.1140/epjc/s10052-020-7616-4 [arXiv:1908.09398 [hep-ph]].
- [110] G. Aad *et al.* [ATLAS], [arXiv:2108.07665 [hep-ex]].
- [111] M. Drees, H. Dreiner, D. Schmeier, J. Tattersall and J. S. Kim, Comput. Phys. Commun. 187 (2015), 227-265 doi:10.1016/j.cpc.2014.10.018 [arXiv:1312.2591 [hep-ph]].
- [112] D. Dercks, N. Desai, J. S. Kim, K. Rolbiecki, J. Tattersall and T. Weber, Comput. Phys. Commun. 221 (2017), 383-418 doi:10.1016/j.cpc.2017.08.021 [arXiv:1611.09856 [hep-ph]].
- [113] M. Aaboud *et al.* [ATLAS], JHEP **06** (2019), 144 doi:10.1007/JHEP06(2019)144 [arXiv:1902.08103 [hep-ex]].
- [114] A. M. Sirunyan *et al.* [CMS], Phys. Rev. D **98** (2018) no.3, 032005 doi:10.1103/PhysRevD.98.032005 [arXiv:1805.10228 [hep-ex]].
- [115] A. M. Sirunyan et al. [CMS], JHEP 07 (2018), 115 doi:10.1007/JHEP07(2018)115 [arXiv:1806.03472 [hep-ex]].
- [116] A. M. Sirunyan *et al.* [CMS], Phys. Lett. B **819** (2021), 136446 doi:10.1016/j.physletb.2021.136446 [arXiv:2012.04178 [hep-ex]].
- [117] B. Gripaios, A. Papaefstathiou, K. Sakurai and B. Webber, JHEP 01 (2011), 156 doi:10.1007/JHEP01(2011)156 [arXiv:1010.3962 [hep-ph]].

- [118] G. Brooijmans, A. Buckley, S. Caron, A. Falkowski, B. Fuks, A. Gilbert, W. J. Murray, M. Nardecchia, J. M. No and R. Torre, *et al.* [arXiv:2002.12220 [hep-ph]].
- [119] G. Bélanger, A. Bharucha, B. Fuks, A. Goudelis, J. Heisig, A. Jueid, A. Lessa, K. A. Mohan, G. Polesello and P. Pani, et al. [arXiv:2111.08027 [hep-ph]].
- [120] M. Aaboud *et al.* [ATLAS], Phys. Rev. D **97** (2018) no.11, 112001 doi:10.1103/PhysRevD.97.112001 [arXiv:1712.02332 [hep-ex]].
- [121] G. Aad *et al.* [ATLAS], JHEP **02** (2021), 143 doi:10.1007/JHEP02(2021)143 [arXiv:2010.14293 [hep-ex]].