

Current Status of the Muon $g - 2$ Interpretations within Two-Higgs-Doublet Models

Syuhei Iguro,^(a,b) Teppei Kitahara,^(c,d) Martin S. Lang,^(a)
and Michihisa Takeuchi^(e,f)

- (a) *Institute for Theoretical Particle Physics (TTP), Karlsruhe Institute of Technology (KIT), Wolfgang-Gaede-Str. 1, 76131 Karlsruhe, Germany*
- (b) *Institute for Astroparticle Physics (IAP), KIT, Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany*
- (c) *Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Nagoya 464–8602, Japan*
- (d) *CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China*
- (e) *School of Physics and Astronomy, Sun Yat-sen University, 519082 Zhuhai, China*
- (f) *Graduate School of Information Science and Technology, Osaka University, Suita, Osaka 565-0871, Japan*

igurosyuhei@gmail.com, teppeik@kmi.nagoya-u.ac.jp, m.lang@kit.edu,
takeuchi@mail.sysu.edu.cn

Abstract

In this review article, we outline implications of the muon anomalous magnetic moment (muon $g - 2$) anomaly for two-Higgs-doublet models (2HDMs), which are classified according to their Yukawa sector and imposed symmetries. In the minimal setup, the muon $g - 2$ anomaly can be accommodated by the type-X (lepto-philic) 2HDM, flavor-aligned 2HDM (FA2HDM), muon-specific 2HDM (μ 2HDM), and $\mu\tau$ -flavor violating 2HDM. We summarize all relevant experimental constraints from high-energy collider experiments and flavor experiments, as well as the theoretical constraints from the perturbative unitarity and vacuum stability bounds, to these 2HDMs in light of the muon $g - 2$ anomaly. We clarify the available parameter spaces of these 2HDMs and investigate how to probe the remaining parameter regions in future experiments. In particular, we find that, due to the updated $B_s \rightarrow \mu^+\mu^-$ measurement, the remaining parameter region of the FA2HDM is almost equivalent to the one of the type-X 2HDM. Furthermore, based on collider simulations, we find that the type-X 2HDM and μ 2HDM scenarios will be covered with the LHC Run 2 full data and the Run 3 full data, respectively.

KEYWORDS: MUON $g - 2$, 2HDM, LARGE HADRON COLLIDER, FLAVOR PHYSICS

Contents

1	Introduction	1
2	Two-Higgs-doublet models	4
2.1	Type-X 2HDM	7
2.2	Flavor-aligned 2HDM	8
2.3	Muon-specific 2HDM	9
2.4	$\mu\tau$ -flavor violating 2HDM	9
3	Current status and prospects for muon $g - 2$ interpretations	10
3.1	Type-X 2HDM	10
3.2	Flavor-aligned 2HDM	16
3.3	Muon-specific 2HDM	18
3.4	$\mu\tau$ -flavor violating 2HDM	20
4	Summary and discussion	22
A	Explicit formulae	24
A.1	Muon $g - 2$	24
A.2	Tau decays	24
A.3	Z decays	26
A.4	$B_s \rightarrow \mu^+ \mu^-$	27
A.5	Perturbative unitarity and vacuum stability	27
A.6	Renormalization group equations	28

1 Introduction

The muon anomalous magnetic moment (the muon $g-2$), defined by $a_\mu = (g_\mu - 2)/2$, is known as a long-standing anomaly in the Standard Model (SM). Based on the white paper recommended values [1], the SM prediction deviates from the experimental values measured at Brookhaven [2–4] and at Fermilab [5],

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (25.1 \pm 5.9) \times 10^{-10}, \quad (1.1)$$

at a significance of 4.2σ . The measured value will be more accurately (and independently) checked by further runs of the Fermilab experiment and by the upcoming J-PARC experiment, which is based on a different measurement technique [6, 7].

One should note that the recent evaluation of the leading-order hadronic vacuum polarization (HVP) contribution to the muon $g - 2$, based on a lattice QCD+QED

simulation [8], casts doubt on the SM prediction of the white paper in Eq. (1.1), giving $\Delta a_\mu = (10.7 \pm 7.0) \times 10^{-10}$, which is consistent with the measured data a_μ^{exp} . This lattice evaluation has been partially confirmed by different lattice collaborations [9–12], using window observables for the HVP [13]. The lattice average based on these window observables provides $\Delta a_\mu = (18.3 \pm 5.9) \times 10^{-10}$, corresponding to a significance of 3.1σ [14]. However, these lattice results are incompatible with $e^+e^- \rightarrow \pi^+\pi^-$ data [15–17] at the 3.8σ level [14].^{#1}

In this article, we assume that the muon $g - 2$ anomaly in Eq. (1.1) is a hint of new physics (NP) beyond the SM and suppose $a_\mu^{\text{NP}} = (25.1 \pm 5.9) \times 10^{-10}$. It is known that many kinds of NP models can accommodate the muon $g - 2$ anomaly, see the recent review paper [22]. We will discuss NP interpretations in the context of two-Higgs-doublet models (2HDMS) in this review article.

In the last decade, our understanding of the Higgs sector has significantly been renewed after the discovery of a 125 GeV scalar boson, with null results in searches for additional Higgs bosons at the large hadron collider (LHC). In the meantime, precision measurements have been improved greatly. For instance, the Yukawa couplings to third-generation fermions have been determined at the 10% level, and furthermore the muon and charm Yukawa couplings were started to be measured. With the larger statistics in the high-luminosity phase of the LHC (HL-LHC), the LHC will gain access to the triple-Higgs coupling and thus will determine the shape of the Higgs potential.

However, the fundamental question of whether the discovered 125 GeV Higgs is the only scalar boson or just one of several scalars remains unanswered. For example, one of the minimal extensions of the SM features an additional Higgs doublet, the so-called 2HDM [23], which naturally accommodates electroweak (EW) precision tests and has rich phenomenology. The 2HDM appears as a low-energy effective scalar sector of many UV-complete theories, *e.g.*, the minimal supersymmetric standard model (MSSM) [24,25], left-right model [26], Pati-Salam model [27], little Higgs model [28,29], and so on.

It is known that the muon $g - 2$ anomaly in Eq. (1.1) is the same size as the EW contributions, which implies that an additional $\mathcal{O}(100)$ GeV electroweakly-interacting particle could possibly explain the discrepancy. The number of variants of the 2HDM that can accommodate the muon $g - 2$ anomaly whilst evading existing experimental constraints is limited, and hence we can take a bottom-up approach in this review article.

Within the 2HDMS, the suppression of flavor changing neutral currents (FCNCs) is achieved once a discrete symmetry is imposed under which the two Higgs doublets and

^{#1}Very recently, the CMD-3 collaboration reported new data for the $e^+e^- \rightarrow \pi^+\pi^-$ cross section [18]. The result is consistent with the lattice simulation [8]. However, the cross section is incompatible with other precision data by the KLOE [19] and BaBar [20] collaborations, as well as the previous data of the same collaboration [21] without sufficient explanation.

fermions carry different charges [23].^{#2} A famous example is a discrete \mathbb{Z}_2 symmetry. With the ad-hoc \mathbb{Z}_2 symmetry, depending on the assignment of the \mathbb{Z}_2 charges to the SM fermions, four different types of Yukawa interactions are allowed [32, 33], known as type-I, type-II, type-X and type-Y 2HDMs. Among the four scenarios, only the type-X 2HDM can explain the $g - 2$ anomaly without conflicting with experimental constraints [34, 35]. In this model, the two-loop Barr-Zee diagram, in which a light CP-odd Higgs and the tau lepton are in the loop, gives the dominant contribution to the muon $g - 2$ [36–45].

An alternative method to eliminate the FCNCs is assuming flavor alignment in the Yukawa matrices for each type of right-handed fermions. Such a model is called the flavor-aligned 2HDM (FA2HDM). In the FA2HDM, all Yukawa interactions are proportional to the corresponding fermion mass matrix [46]. Since the model contains the type-X 2HDM as a certain limit, it is also an interesting candidate for explaining the muon $g - 2$ anomaly. In addition to the tau-lepton Barr-Zee diagram, a top-quark Barr-Zee diagram can also contribute to the muon $g - 2$ in the FA2HDM [39, 47].

More complicated discrete symmetries, \mathbb{Z}_4 , are also discussed. In the muon-specific 2HDM (μ 2HDM) based on a \mathbb{Z}_4 symmetry, the additional Higgs doublet strongly couples only to the muon, without the FCNCs [48]. Another viable method introduces a lepton-flavor-violating scalar particle. The magnetic dipole operator requires a chirality flip, which corresponds to the muon mass within the flavor-conserving scenarios. On the other hand, if a model contains a $\mu\tau$ -flavor violating vertex with a neutral particle, the chirality of the virtual tau lepton can be flipped instead and the one-loop contribution is enhanced by a factor of $m_\tau/m_\mu \simeq 17$ compared to the flavor-conserving scenarios [49–69]. A \mathbb{Z}_4 symmetry can naturally realize such a model within the 2HDM [59], and we call it the $\mu\tau$ -flavor violating 2HDM ($\mu\tau$ 2HDM). In this review article, in light of the muon $g - 2$ anomaly, we update the status of these four possibilities: type-X 2HDM, FA2HDM, μ 2HDM, and $\mu\tau$ 2HDM. Variants of the 2HDM with even larger discrete symmetries can also be conceived; however, their implications for the muon $g - 2$ anomaly are mostly the same as in the μ 2HDM and $\mu\tau$ 2HDM.

The outline of the paper is as follows. In Sec. 2, all kinds of minimal 2HDMs which can resolve the muon $g - 2$ anomaly are introduced. In Sec. 3, we examine various experimental and theoretical constraints on their parameter spaces and discuss the future experimental prospects for them. Finally, conclusions are drawn in Sec. 4. In the Appendix, we collect all relevant formulae for the analyses of this article.

^{#2}The suppression of the FCNCs can also be realized by using an additional $U(1)$ gauge symmetry instead of a discrete symmetry, see *e.g.*, Refs. [30, 31].

2 Two-Higgs-doublet models

First, we introduce two different bases in the various 2HDMs: the so-called \mathbb{Z}_2 basis and the Higgs basis, which are mathematically equivalent.^{#3} The \mathbb{Z}_2 basis respects charge assignments of the discrete symmetry. One can straightforwardly track the free parameters of the model in the \mathbb{Z}_2 basis. On the other hand, the Higgs basis which will be used in this article can parametrize more general 2HDMs. In particular, if one assumes the alignment of the SM Higgs boson, any calculations become significantly simpler in the Higgs basis.

Within the 2HDMs, when the Higgs potential is minimized at the EW symmetry breaking vacuum, both neutral components of Higgs doublets acquire vacuum expectation values (VEVs) in general. In the \mathbb{Z}_2 basis, the Higgs potential is given by

$$\begin{aligned} V(\Phi_1, \Phi_2) = & m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 - \left(m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.} \right) \\ & + \frac{\lambda_1^{Z_2}}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2^{Z_2}}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3^{Z_2} (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_4^{Z_2} (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ & + \frac{1}{2} \left(\lambda_5^{Z_2} (\Phi_1^\dagger \Phi_2)^2 + \text{h.c.} \right), \end{aligned} \quad (2.1)$$

where in general m_{11}^2 , m_{22}^2 and $\lambda_{1-4}^{Z_2}$ are real parameters, while m_{12}^2 and $\lambda_5^{Z_2}$ are complex ones. In this article, we assume absence of CP violation in the Higgs potential for simplicity, so that m_{12}^2 and $\lambda_5^{Z_2}$ are treated as real parameters as well. The two Higgs doublets in the \mathbb{Z}_2 basis are defined as

$$\Phi_1 = \begin{pmatrix} \omega_1^+ \\ \frac{v_1 + h_1 + iz_1}{\sqrt{2}} \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} \omega_2^+ \\ \frac{v_2 + h_2 + iz_2}{\sqrt{2}} \end{pmatrix}. \quad (2.2)$$

The VEVs v_1, v_2 can be taken to be real and positive and need to satisfy $v = \sqrt{v_1^2 + v_2^2} \simeq 246 \text{ GeV}$ in order to reproduce the masses of the weak gauge bosons. The ratio of the VEVs is represented by $\tan \beta = v_2/v_1$ ($0 \leq \beta \leq \pi/2$).

By taking a certain linear combination of $\Phi_{1,2}$, one can always work in the Higgs basis [73–75] where only one Higgs doublet obtains a VEV as

$$\begin{pmatrix} H_1 \\ H_2 \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}. \quad (2.3)$$

In the Higgs basis, the doublets can be decomposed as

$$H_1 = \begin{pmatrix} G^+ \\ \frac{v+h+iG}{\sqrt{2}} \end{pmatrix}, \quad H_2 = \begin{pmatrix} H^+ \\ \frac{H+iA}{\sqrt{2}} \end{pmatrix}, \quad (2.4)$$

^{#3}For the parameter conversions between the two bases, see Appendices of Refs. [70, 71] for the Higgs potential and Ref. [72] for the Yukawa sector.

where the fields G^+ , H^+ , h , H , G , and A are linear combinations of $\omega_{1,2}^+$, $h_{1,2}$, and $z_{1,2}$. The Nambu-Goldstone bosons of the spontaneously broken EW gauge symmetry are denoted by G^\pm and G , and H^\pm denotes an additional charged Higgs boson, while A is a neutral CP-odd Higgs boson. In principle, the CP-even scalars h and H in the doublets mix with an angle α to constitute the mass eigenstates. However, since the LHC found that the interactions of the observed scalar boson are currently consistent with the SM Higgs expectations, we consider the case where the mixing between CP-even scalars is negligible corresponding to a conservative choice, *i.e.*, $\sin(\beta - \alpha) \approx 1$, such that h and H are promoted to mass eigenstates.^{#4} Since experimental constraints are commonly weakened in the Higgs alignment limit, we expect the study in this article to be conservative.

In the Higgs basis, the scalar potential is given by

$$\begin{aligned}
V(H_1, H_2) = & M_{11}^2 H_1^\dagger H_1 + M_{22}^2 H_2^\dagger H_2 - \left(M_{12}^2 H_1^\dagger H_2 + \text{h.c.} \right) \\
& + \frac{\lambda_1}{2} (H_1^\dagger H_1)^2 + \frac{\lambda_2}{2} (H_2^\dagger H_2)^2 + \lambda_3 (H_1^\dagger H_1)(H_2^\dagger H_2) + \lambda_4 (H_1^\dagger H_2)(H_2^\dagger H_1) \\
& + \left\{ \frac{\lambda_5}{2} (H_1^\dagger H_2)^2 + \left[\lambda_6 (H_1^\dagger H_1) + \lambda_7 (H_2^\dagger H_2) \right] (H_1^\dagger H_2) + \text{h.c.} \right\}. \quad (2.5)
\end{aligned}$$

By matching to the Higgs potential in the \mathbb{Z}_2 basis, one obtains

$$\lambda_6 = -\frac{1}{2} \sin 2\beta \left(\cos^2 \beta \lambda_1^{Z_2} - \sin^2 \beta \lambda_2^{Z_2} - \cos 2\beta \lambda_{345}^{Z_2} \right), \quad (2.6)$$

$$\lambda_7 = -\frac{1}{2} \sin 2\beta \left(\sin^2 \beta \lambda_1^{Z_2} - \cos^2 \beta \lambda_2^{Z_2} + \cos 2\beta \lambda_{345}^{Z_2} \right), \quad (2.7)$$

with $\lambda_{345}^{Z_2} = \lambda_3^{Z_2} + \lambda_4^{Z_2} + \lambda_5^{Z_2}$. We note that the Higgs alignment condition corresponds to $\lambda_6 = 0$ at the renormalization scale $\mu \approx m_W$ in the Higgs basis, which leads to $M_{12}^2 = 0$ under the stationary condition.

The scalar mass spectrum is important for our discussion and hence it is useful to show the mass relations in the Higgs alignment limit in the Higgs basis:

$$\begin{aligned}
m_h^2 = \lambda_1 v^2, \quad m_A^2 = M_{22}^2 + \frac{\lambda_3 + \lambda_4 - \lambda_5}{2} v^2, \\
m_H^2 = m_A^2 + \lambda_5 v^2, \quad m_{H^\pm}^2 = m_A^2 - \frac{\lambda_4 - \lambda_5}{2} v^2.
\end{aligned} \quad (2.8)$$

The mass differences among the neutral scalars are crucial parameters to discuss the muon $g - 2$ anomaly and constraints from collider physics. The numerical relation is given as

$$\Delta_{H-A} = m_H - m_A \simeq 60 \left(\frac{\lambda_5}{1.0} \right) \left(\frac{1000 \text{ GeV}}{m_H + m_A} \right) \text{ GeV}. \quad (2.9)$$

^{#4}If non-zero mixing is considered, we have more stringent constraints on the model from, *e.g.*, $h \rightarrow \tau^+ \tau^-$.

We also define

$$\lambda_{hAA} = \lambda_3 + \lambda_4 - \lambda_5, \quad (2.10)$$

for later convenience, which corresponds to the hAA coupling in the Higgs alignment limit.

The most general Yukawa sector of the 2HDM in the fermion mass eigenbasis is given as

$$\begin{aligned} -\mathcal{L} = & \bar{Q}_L^i H_1 y_d^i d_R^i + \bar{Q}_L^i H_2 \rho_d^{ij} d_R^j + \bar{Q}_L^i (V^\dagger)^{ij} \tilde{H}_1 y_u^j u_R^j + \bar{Q}_L^i (V^\dagger)^{ij} \tilde{H}_2 \rho_u^{jk} u_R^k \\ & + \bar{L}_L^i H_1 y_e^i e_R^i + \bar{L}_L^i H_2 \rho_e^{ij} e_R^j + \text{h.c.} . \end{aligned} \quad (2.11)$$

Here i, j denote flavor indices and $Q = (V^\dagger u_L, d_L)^T$ and $L = (U \nu_L, e_L)^T$ are $SU(2)_L$ doublets, where V and U are the Cabibbo-Kobayashi-Maskawa (CKM) and the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrices, respectively. In writing Eq. (2.11) we have assumed that neutrino masses are explained by the seesaw mechanism introducing heavy right-handed neutrinos, so that in the low-energy effective theory the left-handed neutrinos have a 3×3 Majorana mass matrix. Note that the Yukawa couplings y_f are expressed in terms of the fermion masses m_f , $y_f = \sqrt{2} m_f / v$. On the other hand, the Yukawa couplings ρ_f^{ij} are a priori arbitrary 3×3 complex matrices and can in general be sources of flavor violation mediated by additional neutral Higgs bosons at tree level. In the Higgs alignment limit, the interactions of H_1 are exactly the same as the ones of the SM Higgs doublet.

Following the notation of Ref. [72], in terms of the mass eigenstates of the Higgs bosons, the Yukawa interactions are represented by

$$\begin{aligned} -\mathcal{L} = & \sum_{f=u,d,e} \sum_{\phi=h,H,A} y_{\phi ij}^f \bar{f}_{Li} \phi f_{Rj} + \text{h.c.} \\ & + \bar{\nu}_{Li} (U^\dagger \rho_e)^{ij} H^+ e_{Rj} + \bar{u}_i (V \rho_d P_R - \rho_u^\dagger V P_L)^{ij} H^+ d_j + \text{h.c.} , \end{aligned} \quad (2.12)$$

where

$$y_{hij}^f = \frac{1}{\sqrt{2}} y_f^i \delta_{ij} = \frac{m_{f_i}}{v} \delta_{ij}, \quad y_{Hij}^f = \frac{1}{\sqrt{2}} \rho_f^{ij}, \quad y_{Aij}^f = \begin{cases} -\frac{i}{\sqrt{2}} \rho_f^{ij} & \text{for } f = u, \\ \frac{i}{\sqrt{2}} \rho_f^{ij} & \text{for } f = d, e. \end{cases} \quad (2.13)$$

The off-diagonal components of the Yukawa couplings ρ_f^{ij} ($i \neq j$) induce FCNCs from decays of the scalar bosons and Higgs-mediated processes. The absence of such FCNCs at experiments^{#5} motivates us to impose a discrete symmetry which distinguishes the two Higgs doublets $\Phi_{1,2}$. The Yukawa structure of ρ_f^{ij} depends on the charge assignment of the Higgs doublets and fermions as well, which we will classify in the following sections.

^{#5}The recent ATLAS and CMS results both show hints of Higgs lepton-flavor violating decays [76–78]. In order to address these slight deviations, one needs a small neutral-scalar mixing.

	Φ_1	Φ_2	Q	u_R	d_R	(L_e, L_μ, L_τ)	(e_R, μ_R, τ_R)
type-I	0	1	0	1	1	0	1
type-II	0	1	0	1	0	0	0
type-X	0	1	0	1	1	0	0
type-Y	0	1	0	1	0	0	1
μ 2HDM	2	0	0	0	0	$(0, -1, 0)$	$(0, 1, 0)$
$\mu\tau$ 2HDM	2	0	0	0	0	$(0, 1, -1)$	$(0, 1, -1)$

Table 1. The charge assignments under the discrete symmetry: the matter fields are transformed as $\psi \rightarrow \exp(2\pi q_\psi i/N) \psi$ under the discrete \mathbb{Z}_N symmetry. We show the \mathbb{Z}_2 charge assignments for the type-I, II, X, and Y 2HDMs (0 is \mathbb{Z}_2 -even, while 1 is \mathbb{Z}_2 -odd) [79] and the \mathbb{Z}_4 charge assignments for the μ 2HDM [48] and the $\mu\tau$ 2HDM [59].

2.1 Type-X 2HDM

We start with the type-X 2HDM. This is one of the four types of 2HDMs with softly-broken \mathbb{Z}_2 symmetry and thus naturally suppresses FCNCs. The \mathbb{Z}_2 assignment is summarized in Table 1. Out of the four \mathbb{Z}_2 -symmetric 2HDMs, only the type-X 2HDM can resolve the muon $g - 2$ anomaly because in all three other 2HDMs a large contribution to a_μ^{NP} is not allowed by flavor and collider constraints and the perturbative unitarity bound. In the type-I and type-Y 2HDMs, both ρ_u^{ii} and ρ_e^{ii} are proportional to $\cot \beta$. Therefore, the upper limit on ρ_u^{tt} from perturbativity indirectly sets an upper limit on $\rho_e^{\mu\mu}$, suppressing any potential contributions to a_μ^{NP} . On the other hand, in the type-II and type-X 2HDMs, ρ_e^{ii} is proportional to $\tan \beta$ while ρ_u^{ii} remains proportional to $\cot \beta$. It is noted that the type-II and type-X 2HDMs cannot accommodate the muon $g - 2$ anomaly at one-loop level with a light CP-even Higgs. If one tries to explain Δa_μ with $\rho_e^{\mu\mu} = \mathcal{O}(1)$, $\rho_e^{\tau\tau}$ immediately becomes non-perturbative because $\rho_e^{\tau\tau}/\rho_e^{\mu\mu}$ is fixed by m_τ/m_μ . Instead, the contributions to a_μ^{NP} can be dominated by the so-called Barr-Zee diagram [80] at two-loop level with a light CP-odd Higgs [36–45]. We exhibit the Feynman diagrams in Fig. 1 and the formula in Appendix A.1.

The difference between the type-II and type-X 2HDMs is the down-type quark Yukawa couplings to the extra Higgs bosons. In the type-II 2HDM, both the down-type quark and charged lepton couplings are enhanced by $\tan \beta$ at the same time. Therefore, the model is severely constrained by B -meson flavor physics and direct searches for extra Higgs bosons. The rare radiative decay $b \rightarrow s\gamma$ gives a lower mass limit for the charged Higgs boson of $m_{H^\pm} \geq 800 \text{ GeV}$ [81]. For the muon $g - 2$ anomaly, we need a $\mathcal{O}(10\text{--}100) \text{ GeV}$ light scalar at the same time. Such a large mass difference




Figure 1. The leading-order contributions to the muon $g-2$ in the μ 2HDM (left), $\mu\tau$ 2HDM (right), and type-X 2HDM and FA2HDM (bottom). The bottom diagram is the two-loop Barr-Zee diagram where $f = \tau, t$.

is troublesome since the theory will be non-perturbative at less than 1 TeV.

On the other hand, only the lepton Yukawa couplings are enhanced by $\tan\beta$ in the type-X 2HDM. Therefore, the constraints from B -meson decays are weaker compared to the type-II ones. The Yukawa structure in the type-X 2HDM is given by

$$\rho_u = \frac{\sqrt{2}m_{u_i}}{v}\xi^{-1}, \quad \rho_d = \frac{\sqrt{2}m_{d_i}}{v}\xi^{-1}, \quad \rho_e = -\frac{\sqrt{2}m_{e_i}}{v}\xi, \quad (2.14)$$

with $\xi = \tan\beta$ and all non-diagonal Yukawa couplings vanishing. Although $\tan\beta$ is conventionally used, we will use the notation ξ in this article in order to allow for an easy comparison with the other 2HDM scenarios.

2.2 Flavor-aligned 2HDM

In the flavor-aligned 2HDM (FA2HDM), it is assumed that the Yukawa interactions of the additional scalars are proportional to mass matrices ($\rho_f \propto y_f$); both the ρ_f and y_f matrices are simultaneously diagonalized in the Lagrangian of Eq. (2.11) [46]. Therefore, tree-level FCNCs are absent in this model.^{#6} Viable UV models for the flavor alignment condition are discussed in Refs. [84, 85].

^{#6}Even if one imposes the flavor alignment at tree-level, FCNCs are in general induced radiatively, in particularly by the renormalization group evolutions (RGEs). However, these RGE-induced FCNCs are shown to be too small to be probed at current experiments [82, 83].

There are three free parameters in the Yukawa interactions, ξ_u , ξ_d , and ξ_e . In other words, the type-X scenario is a special case of the FA2HDM, in which $\xi_u = \xi_d = \xi^{-1}$ and $\xi_e = -\xi$. The additional Yukawa couplings are given as

$$\rho_u = \frac{\sqrt{2}m_{u_i}}{v}\xi_u, \quad \rho_d = \frac{\sqrt{2}m_{d_i}}{v}\xi_d, \quad \rho_e = \frac{\sqrt{2}m_{e_i}}{v}\xi_e. \quad (2.15)$$

In this model, the dominant contribution to a_μ^{NP} comes from the two-loop Barr-Zee diagram with a light CP-odd Higgs and a tau lepton, while the Barr-Zee diagram with a top-quark loop can also contribute [39, 47].

2.3 Muon-specific 2HDM

Another interesting scenario is the so-called muon-specific 2HDM (μ 2HDM). In this model, the additional scalars dominantly couple to muons due to a \mathbb{Z}_4 charge assignment as summarized in Table 1. This coupling structure helps to avoid the constraints from τ decays and loop-induced Z decays [48]. The Yukawa coupling structure of this model is given as

$$\rho_f = \frac{\sqrt{2}m_f}{v}\xi^{-1}, \quad \rho_e^{\mu\mu} = -\frac{\sqrt{2}m_\mu}{v}\xi, \quad (2.16)$$

with $\xi = \tan\beta$, where f denotes all fermions except for μ . The dominant contribution to a_μ^{NP} stems from one-loop diagrams, see Fig. 1 (top-left), and the formula is given in Appendix A.1. A phenomenologically interesting parameter region is $\xi \gg 1$.

2.4 $\mu\tau$ -flavor violating 2HDM

In the $\mu\tau$ -flavor violating 2HDM ($\mu\tau$ 2HDM), a discrete \mathbb{Z}_4 symmetry with charge assignments shown in Table 1 is imposed. Due to the unique charge assignment, after the diagonalization of the charged lepton mass, the only flavor-violating interactions of the additional neutral scalars in the Higgs alignment limit are the $\bar{\mu}_{L/R}\tau_{R/L}H(A)$ interactions. The Yukawa coupling structure of this model is given as

$$\rho_f^{\text{diag}} = \frac{\sqrt{2}m_f}{v}\xi^{-1}, \quad \rho_e^{\mu\tau} \neq 0, \quad \rho_e^{\tau\mu} \neq 0, \quad (2.17)$$

with $\xi \approx \tan\beta$. Here, f denotes all fermions. The dominant contribution to a_μ^{NP} comes from the one-loop diagram with a virtual tau lepton in Fig. 1, and the formula is given in Appendix A.1.

However, even in the case of the Higgs alignment limit, this model also predicts $\tau \rightarrow \mu$ lepton-flavor violating transitions at one-loop level. To avoid their experimental bounds, the limit $\tan\beta \rightarrow \infty$ is a natural solution, corresponding to the original discrete \mathbb{Z}_4 symmetry (a variant of the inert doublet model) proposed in Ref. [59].

	Δa_μ	mass range	precision	LHC	life time
Type-X 2HDM	2 loop	$m_A = \mathcal{O}(10) \text{ GeV} \ll m_H = m_{H^\pm}$	$h \rightarrow AA, Z, \tau$ decays	multi- τ	Run 2
FA2HDM	2 loop	$m_A = \mathcal{O}(10) \text{ GeV} \ll m_H = m_{H^\pm}$	$B_s \rightarrow \mu^+ \mu^-$, $h \rightarrow AA$	multi- τ	Run 2
μ 2HDM	1 loop	$900 \text{ GeV} \leq m_{A,H} \leq 1000 \text{ GeV}$	Z decay	multi- μ	Run 3
$\mu\tau$ 2HDM	1 loop	$500 \text{ GeV} \leq m_{A,H} \leq 1600 \text{ GeV}$	$\tau \rightarrow \mu\nu\bar{\nu}$	$\mu^\pm \mu^\pm \tau^\mp \tau^\mp$	HE-LHC

Table 2. Summary table for all 2HDM scenarios which can accommodate the muon $g - 2$ anomaly. The second column shows the loop order of the dominant contribution to a_μ^{NP} . The third column lists the available mass range of scalars. The fourth and fifth ones show the relevant constraints from precision measurements and important processes at LHC, respectively. The last column summarizes how much data is needed to fully explore the parameter space where the muon $g - 2$ anomaly can be solved at the 1σ level.

Since a_μ^{NP} is insensitive to $\tan\beta$ in this model, we consider the following Yukawa coupling structure

$$\rho_f^{\text{diag}} = 0, \quad \rho_e^{\mu\tau} \neq 0, \quad \rho_e^{\tau\mu} \neq 0. \quad (2.18)$$

Then, we can safely focus on the phenomenology of the additional scalars. Note that for realistic neutrino masses and mixing, breaking of the \mathbb{Z}_4 symmetry is necessary [69].

3 Current status and prospects for muon $g - 2$ interpretations

In this main section, we discuss explanations of the muon $g - 2$ anomaly based on the models introduced in Sec. 2, along with relevant flavor and collider constraints. The future prospects at the (HL-)LHC are also discussed. Table 2 summarizes the interesting mass range and relevant processes.

3.1 Type-X 2HDM

The type-X model explains the muon $g - 2$ with a two-loop Barr-Zee diagram in which a CP-odd scalar A of $\mathcal{O}(20\text{--}40) \text{ GeV}$ propagates. In the large ξ limit, the two-loop Barr-Zee correction with a light A and tau internal loop can generate a large positive contribution to a_μ^{NP} since it is enhanced by m_τ^2/m_μ^2 . The Barr-Zee contribution with H in the loop gives a negative contribution and thus a heavier H is preferred to enhance a_μ^{NP} .

In this mass range, the SM Higgs boson can decay into a pair of light CP-odd scalars, which modifies the Higgs total width. This additional decay channel opens




Figure 2. The parameter plane of ξ and m_A in the type-X 2HDM. The cyan and blue regions can accommodate the muon $g - 2$ anomaly at the 1σ and 2σ levels, respectively. The region to the left of the green, purple, and brown lines is excluded by the τ decay, Z decay, and scalar bremsstrahlung constraints, respectively. The $B_s \rightarrow \mu^+\mu^-$ constraint excludes the region to the left of the red line. We take $m_H = m_{H^\pm} = 250$ (300) GeV on the left (right) panel.

for $m_h \geq 2m_A$ and the tree-level $h \rightarrow AA$ partial decay width is given as

$$\Gamma(h \rightarrow AA) = \frac{\lambda_{hAA}^2 v^2}{32\pi m_h} \sqrt{1 - \frac{4m_A^2}{m_h^2}}. \quad (3.1)$$

Recent Higgs width measurements restrict the trilinear Higgs coupling to [86, 87]

$$|\lambda_{hAA}| \leq 0.03. \quad (3.2)$$

A more stringent limit $|\lambda_{hAA}| \lesssim 0.01$ is obtained for $m_A \leq 21$ GeV based on searches for $h \rightarrow AA \rightarrow \mu^+\mu^-\tau^+\tau^-$ decays [88]. Since the non-discovery of H^\pm prefers large mass differences between the CP-odd and charged scalars, $\mathcal{O}(1)$ couplings in the Higgs potential are necessary, see Eq. (2.8). Therefore, Eq. (3.2) requires parameter tuning at the 1% level.

In Fig. 2, we show the parameter regions where the muon $g - 2$ anomaly can be explained at the 1σ and 2σ level by the cyan and blue regions, respectively. We take $m_H = m_{H^\pm} = 250$ (300) GeV on the left (right) panel.

In the regime of large lepton couplings in the type-X 2HDM, the tau (leptonic and hadronic) decays are modified by tree-level and one-loop corrections from the additional scalars. The tree-level and one-loop corrections have been calculated in Refs. [36, 89]. Relevant formulae and the current experimental data are summarized in Appendix A.2. Since the corrections from the additional scalars are suppressed

by $1/m_{H^\pm}^2$, the τ -decay bound uniformly becomes weaker, the heavier the additional scalar becomes. Since the one-loop correction involving two tau-Yukawa couplings is larger than the one involving a single muon-Yukawa coupling, the box diagram is less important. The excluded regions from tau decays are shown by the green lines in Fig. 2.

In addition, the one-loop correction to the fermion coupling of the Z boson provides an important cross-check in this scenario. Thanks to the hierarchical structure of the Yukawa couplings, the Z -boson interaction with the tau leptons is most sensitive to the additional scalars. For the partial cancellation of systematic uncertainties, taking the ratio of leptonic decay widths improves the sensitivity. The LEP average is given as [90],^{#7}

$$\frac{\Gamma(Z \rightarrow \tau^+\tau^-)}{\Gamma(Z \rightarrow e^+e^-)} = 1.0019 \pm 0.0032, \quad \frac{\Gamma(Z \rightarrow \mu^+\mu^-)}{\Gamma(Z \rightarrow e^+e^-)} = 1.0009 \pm 0.0028, \quad (3.3)$$

with a correlation of 0.63. A larger mass difference of $m_H - m_A$ yields a larger deviation from the SM prediction of the Z -boson interaction. Furthermore, the tau polarization asymmetry in $Z \rightarrow \tau^+\tau^-$ is also precisely measured at the LEP and the average is [93]

$$A_\tau = 0.143 \pm 0.004. \quad (3.4)$$

The corresponding corrections involving additional scalars are summarized in Appendix A.3. The excluded regions from Z -boson decays are shown by the purple lines in Fig. 2. Contrary to the conventional decoupling behaviour, it is known that the additional scalar contributions are enhanced if $m_A \ll m_Z \ll m_H \simeq m_{H^\pm}$ is considered [40]. Therefore, the Z -boson bound becomes stricter for heavier additional scalars, which provides an exclusion region complementary to the τ -decay bound.

Furthermore, the rare leptonic meson decay $B_s \rightarrow \mu^+\mu^-$ gives a lower bound on m_A , which is independent of ξ , since in the diagram with a neutral scalar the ξ dependence is cancelled. In the past, the experimental world average of $\text{BR}(B_s \rightarrow \mu^+\mu^-)$ had deviated from the SM prediction [94–96] by about 2σ [97, 98]. However, the CMS collaboration recently reported the Run 2 full analysis and found the branching ratio to be consistent with the SM prediction [99]. As a result, the latest experimental world average is well consistent with the SM prediction [100]. The dominant contribution to the $B_s \rightarrow \mu^+\mu^-$ comes from the light CP-odd scalar mediated diagram at one-loop, see Appendix A.4 for more details. The new world average thus leads to a relaxed lower bound on m_A compared to the previous world average. We obtain $m_A \gtrsim 27 \text{ GeV}$ in the type-X 2HDM, which is shown by the red lines in Fig. 2.

^{#7}It is noted that the recent result of $\frac{\Gamma(Z \rightarrow \mu^+\mu^-)}{\Gamma(Z \rightarrow e^+e^-)}$ from ATLAS [91], which has a twice larger uncertainty, is consistent with the LEP result. In addition, the uncertainty from LHCb is much larger than that of the LEP [92]. Furthermore, the correlation matrix is not available in the PDG [93]. Therefore we use the LEP average.

The LEP probed electroweak AH production in 4τ final states ($e^+e^- \rightarrow Z^* \rightarrow AH \rightarrow 2\tau^+2\tau^-$) without finding a significant excess over the SM expectation. As a result a lower limit on the sum of the neutral scalar masses, $m_H+m_A \geq 190$ (155) GeV, has been obtained for the case where $\text{BR}(A \rightarrow \tau^+\tau^-) \times \text{BR}(H \rightarrow \tau^+\tau^-) = 1(0.1)$ [101, 102]. Lighter scalars have been also searched for in the scalar bremsstrahlung process ($e^+e^- \rightarrow \tau^+\tau^-A \rightarrow 2\tau^+2\tau^-$) [101]. This constraint is especially stringent for a very light A . An upper bound of $\xi \leq 34$ (83) was obtained for $m_A = 10$ (25) GeV with $\text{BR}(A \rightarrow \tau^+\tau^-) \simeq 1$. Furthermore, there have been searches for a pair of H^\pm in $\tau\nu$, cs and WA decay modes. Again the absence of events exceeding the SM expectation allows us to set the lower mass limit as $m_{H^\pm} \geq 84$ –94 GeV, depending on m_A [103]. The LEP exclusions, which come from the scalar bremsstrahlung process, are shown by the brown lines in Fig. 2.

The LHC is also a powerful tool to search for additional scalars. Due to the nature of a hadron collider, only partial information of a collision is accessible. Thanks to the large statistics and good control of tau-lepton identification, low-mass charged Higgs bosons have recently been excluded [104, 105]. The Run 2 full result for the search for left-handed staus can be directly adapted to draw exclusion plots because the production cross section is the same as the charged-Higgs pair production. The ATLAS measurement excluded $120 \text{ GeV} \leq m_{H^\pm} \leq 390 \text{ GeV}$ [106] while the CMS one excluded $115 \text{ GeV} \leq m_{H^\pm} \leq 340 \text{ GeV}$ [107], assuming $\text{BR}(H^\pm \rightarrow \tau\nu) = 1$. Since the SM background (BG) stems from W^\pm pair production, it is difficult to probe lower values of m_{H^\pm} at the LHC. The low mass window $m_{H^\pm} \approx 100 \text{ GeV}$ will close in the near future [108] once the systematic error scales as $1/\sqrt{L}$. If a large mass difference between the neutral scalars is assumed, searches for same-sign H^\pm become relevant [109]. However, a more dedicated experimental analysis is necessary to estimate the sensitivity. It is worth mentioning that the decay $h \rightarrow AZ$ is absent in the alignment limit considered throughout this review.

In order to precisely interpret the constraints by the LHC searches, the decay properties of the additional scalars are important. In addition to $\text{BR}(A \rightarrow \tau^+\tau^-) = 1$, we summarize here the relevant parameter dependence as follows:

$$\text{BR}(H \rightarrow \tau^+\tau^-) = \frac{\Gamma(H \rightarrow \tau^+\tau^-)}{\Gamma(H \rightarrow \tau^+\tau^-) + \Gamma(H \rightarrow AZ)}, \quad (3.5)$$

$$\text{BR}(H^\pm \rightarrow \tau\nu) = \frac{\Gamma(H^\pm \rightarrow \tau\nu)}{\Gamma(H^\pm \rightarrow \tau\nu) + \Gamma(H^\pm \rightarrow AW^\pm)}, \quad (3.6)$$

$$\Gamma(H \rightarrow \tau^+\tau^-) \simeq \Gamma(H^\pm \rightarrow \tau\nu) = \frac{m_H m_\tau^2}{8\pi v^2} \xi^2, \quad (3.7)$$

$$\Gamma(H \rightarrow AZ) = \frac{m_H^3}{16\pi v^2} \lambda^{3/2} \left(\frac{m_A^2}{m_H^2}, \frac{m_Z^2}{m_H^2} \right), \quad (3.8)$$

$$\Gamma(H^\pm \rightarrow AW^\pm) = \frac{m_{H^\pm}^3}{16\pi v^2} \lambda^{3/2} \left(\frac{m_A^2}{m_{H^\pm}^2}, \frac{m_W^2}{m_{H^\pm}^2} \right), \quad (3.9)$$

where $\lambda(x_1, x_2) = (1 - x_1 - x_2)^2 - 4x_1x_2$ and m_τ is neglected in the phase-space factor. Thus, for large ξ , tauonic scalar decays can make up a significant fraction of the total decay width. For the relevant mass scale, $\xi \sim 100$ yields branching ratios $\sim \mathcal{O}(50\%)$, which means that the branching ratios and the resulting LHC constraints are sensitive in this interesting parameter region.

For the type-X 2HDM interpretation of the muon $g - 2$ anomaly, the favored parameter regions are at large ξ and very light $m_A \sim 30$ GeV, see Fig. 2. In this case, τ -rich signatures at the LHC become relevant, as discussed in Ref. [38]. Although a type-X specific search has not yet been performed, we can recast the current experimental searches for charginos and neutralinos in the MSSM, based on these τ -rich signatures [110–114]. In particular, we consider the ATLAS analysis [114] which provides detailed kinematic cuts. In the analysis, eight signal regions (SRs) are defined, designed such that the sensitivity to $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ and $\tilde{\chi}^\pm \tilde{\chi}_2^0$ events is enhanced. As common features of all the SRs, at least two τ -leptons, a veto on bottom quark jets in order to reject SM top-quark processes, and large stransverse mass [115, 116] $m_{T2} > 70$ –100 GeV are required. Some of the SRs aiming for $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ and $\tilde{\chi}_1^+ \tilde{\chi}_2^0$ require opposite-sign (OS) τ leptons, while the rest requires same-sign taus for $\tilde{\chi}_1^+ \tilde{\chi}_2^0$. As for the SRs with OS taus, a Z/h veto ($m_{\tau\tau} > 120$ GeV) is required to capture the taus from stau decays, otherwise $m_{\tau\tau} \simeq m_h$ is required to capture the $h \rightarrow \tau^+ \tau^-$ in the decay chain. Regarding the missing momentum, both possibilities of $E_{\text{miss}} < 150$ GeV and $E_{\text{miss}} > 150$ GeV are considered to capture the low-mass and the high-mass spectra, respectively. Additional selection cuts are imposed depending on the SRs, for details see Ref. [114]. As a result, 1–14 events were observed in each SR for 139 fb^{-1} at $\sqrt{s} = 13$ TeV, without significant excess. This results in 95% confidence level (CL) upper limits on the non-SM fiducial cross section ($\sigma_{\text{vis}}^{95\%}$) of 0.03–0.1 fb, which put severe bounds even on EW production processes. This stringent constraint compared to the previous one is achieved by the very strong selection cut optimized for the chargino and neutralino searches and due to the large integrated luminosity of 139 fb^{-1} .

We generate EW pair-production events of additional scalars,

$$pp \rightarrow HA, H^\pm A, H^\pm H, H^\pm H^\mp (\rightarrow \text{multi-}\tau), \quad (3.10)$$

in the type-X 2HDM using MADGRAPH5_aMC@NLO [117] + PYTHIA [118] + DELPHES [119] and apply the selection cuts defined for the above SRs [114]. We consider the cases where $m_A \leq m_H = m_{H^\pm}$, and $10 \text{ GeV} < m_A < 200 \text{ GeV}$, $100 \text{ GeV} < m_H < 500 \text{ GeV}$. The total production cross section ranges from 5 fb to 4 pb. For each model point, we generate 100K signal events. Figure 3 shows the contour plots of the ratio $\sigma_{\text{vis}}^{\text{type-X}} / \sigma_{\text{vis}}^{95\%}$ (which we call exclusion factor), that is, how large an event number is expected relative to the 95% CL upper limit. The maximal exclusion factors are mainly from the SRs C1C1-LM and C1N2SS, which were designed to capture chargino pair-production events and chargino-neutralino pair-production




Figure 3. The exclusion factor ($\sigma_{\text{vis}}^{\text{type-X}}/\sigma_{\text{vis}}^{95\%}$) obtained by the MSSM $\tilde{\chi}^{\pm}\tilde{\chi}^0$ searches at the LHC and $\text{BR}(H \rightarrow \tau^+\tau^-)$ are shown in the solid and dashed contours, respectively, in the type-X 2HDM. The parameter ξ is fixed to explain the muon $g-2$ anomaly at the 0σ (left panel) and -2σ (right panel) levels, corresponding to $a_{\mu}^{\text{NP}} = 25.1 \times 10^{-10}$ and 13.3×10^{-10} , respectively. The pale blue region in the bottom-right corner corresponds to $m_H < m_A$.

events with same-sign τ signatures, respectively. We only take the largest of these SRs. We fix $\xi (= \tan\beta)$ as to reproduce Δa_{μ} at the 0σ (left panel) and -2σ levels (right panel), corresponding to $a_{\mu}^{\text{NP}} = 25.1 \times 10^{-10}$ and 13.3×10^{-10} , respectively. The contours in dotted lines show the expected value of $\text{BR}(H \rightarrow \tau^+\tau^-)$. The results show that the interesting regions for the muon $g-2$ anomaly are completely excluded and it is difficult to save this model unless new decay modes are introduced. We also checked that selecting any values of $\text{BR}(H \rightarrow \tau^+\tau^-)$ and $\text{BR}(H^{\pm} \rightarrow \tau\nu)$ results in the whole region of the depicted plane being excluded as long as $\text{BR}(H \rightarrow \tau^+\tau^-) + \text{BR}(H \rightarrow ZA) = 1$ and $\text{BR}(H^{\pm} \rightarrow \tau\nu) + \text{BR}(H^+ \rightarrow W^+A) = 1$. Even if setting $\text{BR}(H \rightarrow \tau^+\tau^-) = \text{BR}(H^{\pm} \rightarrow \tau\nu) = 0$, the lowest value of the exclusion factor is about 1.6. These results are the updated plots of Fig. 7 in Ref. [38].

Finally, we show the same exclusion factor defined above in the ξ - $m_H (= m_{H^{\pm}})$ plane with fixed $m_A = 30 \text{ GeV}$ in Fig. 4. In general, if $(m_H - m_A) > m_Z, m_W$, the exclusion factor becomes stronger if the ratio ξ/m_H increases. This is due to the larger production cross sections and the larger τ -branching ratio, respectively, because the acceptance in the relevant SRs is larger in the tau modes. If $(m_H - m_A) < m_Z, m_W$, specifically at $m_H = 100 \text{ GeV}$ in this plot, the exclusion factor weakens although still larger than ten. The cyan and blue regions can explain the muon $g-2$ anomaly at the 1σ and 2σ levels, respectively. One can also see explicitly that even for $\text{BR}(H \rightarrow \tau^+\tau^-) = \text{BR}(H^{\pm} \rightarrow \tau\nu) = 0$, which is equivalent to the $\xi \rightarrow 0$ limit, the exclusion factor is always more than 1, with the minimum around 2 in the bottom-




Figure 4. The contour plot of the exclusion factor ($\sigma_{\text{vis}}^{\text{type-X}}/\sigma_{\text{vis}}^{95\%}$) with fixed $m_A = 30$ GeV, in the type-X 2HDM. The cyan and blue regions can explain the muon $g - 2$ anomaly at the 1σ and 2σ levels, respectively.

right corner.

In conclusion, the type-X 2HDM interpretation of the muon $g - 2$ anomaly in the mass regions we considered is completely excluded by the chargino pair-production and chargino-neutralino pair-production searches at the LHC Run 2, as shown in Figs. 3 and 4. It is impressive that these searches using the tremendous luminosity accumulated at the LHC already provide such high sensitivities for EW production processes.

3.2 Flavor-aligned 2HDM

In the FA2HDM, the Yukawa couplings ρ_f are proportional to the mass matrix but the overall constants of up-type and down-type quarks and leptons are independent free parameters. Therefore, the model includes the type-X 2HDM and thus has a broader parameter space.

In addition to the Barr-Zee diagram with the tau-lepton in the loop, the one with a top quark in the loop could contribute to the muon $g - 2$. However, the constraint from $B_s \rightarrow \mu^+\mu^-$ sets an upper limit on the top-loop contribution to the muon $g - 2$. This implies a stringent constraint for the light CP-odd scalar scenario. It is worth mentioning that $B_s \rightarrow \mu^+\mu^-$ can also receive “type-II 2HDM”-like contributions proportional to $\xi_d \xi_e$. If ξ_d is of order $\xi_u m_t/m_b$, these contributions can become significant despite the much lighter bottom-quark mass. This can weaken the




Figure 5. The parameter plane spanned by ξ_e and ξ_u in the FA2HDM. The cyan and blue regions accommodate the muon $g - 2$ anomaly at the 1σ and 2σ levels, respectively. The regions above the red lines are excluded by $B_s \rightarrow \mu^+\mu^-$ at the 95% CL. The black contours show the size of the top-loop Barr-Zee contribution in units of Δa_μ . The regions to the right of the green, purple and brown lines are excluded by τ decay, Z decay, and scalar bremsstrahlung constraints, respectively. We take $m_A = 20$ (30) GeV and $m_H = m_{H^\pm} = 250$ (300) GeV on the left (right) panel. The gray dashed lines represent the type-X 2HDM limit.

top-quark loop constraints from $B_s \rightarrow \mu^+\mu^-$, in particular if a fine-tuned cancellation between ξ_d and ξ_u occurs. However, we do not consider such cancellations here and thus neglect contributions involving ξ_d .

In Fig. 5, we show the muon $g - 2$ favored region in the ξ_e vs. ξ_u plane where $m_A = 20$ GeV and $m_A = 30$ GeV are fixed on the left and right panels, respectively. The heavier scalar masses are fixed as $m_H = m_{H^\pm} = 300$ GeV. The red, green, purple and brown lines are excluded by $B_s \rightarrow \mu^+\mu^-$, τ decay, Z decay, and scalar bremsstrahlung constraints, respectively. The black contours show the size of the top-loop Barr-Zee contribution in units of Δa_μ . It is found that the top-quark loop constitutes up to about 1% of the total deviation for $m_A = 20$ GeV. For $m_A = 30$ GeV, the contribution is still less than 2%. The constraint becomes more stringent when A gets lighter. We found that even with $m_A = 50$ GeV the contribution can only be up to 5%. In order to evade the stringent bound from $B_s \rightarrow \mu^+\mu^-$, ξ_u needs to be as small as $\mathcal{O}(0.01)$, implying that this scenario is almost identical to the type-X 2HDM in the context of the muon $g - 2$ anomaly. Note that the gray dashed lines in Fig. 5 represent the type-X 2HDM limit.

If the Yukawa coupling ξ_u is small, other flavor observables and collider processes are less affected, *e.g.*, the single-scalar production via gluon fusion and di-Higgs production cross section are too small to be probed [71]. Therefore, the key probe is the

same as in the type-X 2HDM case and we refer the reader to the discussion in the previous section. As a result, multi- τ signatures at the LHC exclude the explanation of the muon $g - 2$ anomaly within the FA2HDM. We emphasize that the presence of ξ_d cannot significantly reduce the $A \rightarrow \tau^+\tau^-$ branching ratio since ξ_e is already large in any attempt to explain the muon $g - 2$ anomaly. However, the coupling ξ_d could also contribute to the production of scalars and $B_s \rightarrow \mu^+\mu^-$.

3.3 Muon-specific 2HDM

In the μ 2HDM, the one-loop contribution to a_μ^{NP} can explain the muon $g - 2$ anomaly, while the two-loop Barr-Zee contribution is suppressed by the mass of the heavy additional scalars and the electromagnetic coupling constant α . It is known that H should be lighter than A in order to have the correct sign of a_μ^{NP} . In this case $m_H \simeq m_{H^\pm}$ is favored to satisfy both the vacuum stability condition and T -parameter constraint [120]. The size of the mass difference is controlled by the Higgs quartic couplings and thus constrained by the RGE-based perturbative unitarity constraint, see Appendices A.5 and A.6 for details. The black lines of Fig. 6 show the values of $\rho_e^{\mu\mu}$ required to explain the central value of the muon $g - 2$ anomaly (top), -1σ (middle), and -2σ (bottom) levels, corresponding to $a_\mu^{\text{NP}} = 25.1 \times 10^{-10}$, 19.2×10^{-10} , and 13.3×10^{-10} , respectively. The orange region requires $\rho_e^{\mu\mu} \geq \sqrt{4\pi}$ (corresponding to $\xi_e \simeq 5900$), violating perturbativity, and thus we do not consider it.

Since the additional scalars mainly couple to muons, direct searches at the LHC, *e.g.*, searches for smuons or multi-lepton final states, give a lower bound on the masses of the additional scalars. Previously, the authors of Ref. [48] found that the multi-lepton search performed with the 35.9 fb^{-1} data set at $\sqrt{s} = 13 \text{ TeV}$ [121] excludes Higgs masses below $m_H \simeq 620 \text{ GeV}$.^{#8} We updated the analysis with the Run 2 full data [122]. Note that there is a similar search for NP in multi-lepton final states [123], originally motivated by heavy vector-like leptons in the type-III seesaw model. However, we cannot directly use the result since additional jets are required.

We generated 100K signal events for the process $pp \rightarrow \phi H^\pm \rightarrow 3\mu + \nu_\mu$, shown in Fig. 7 (top-left), with MADGRAPH5_aMC@NLO [117] for a given set of H^\pm and ϕ masses at $\sqrt{s} = 13 \text{ TeV}$, where ϕ denotes A or H . Then the scalar sum of charged-lepton p_T (L_T) and missing transverse energy (MET) is calculated and compared to the result in Fig. 3-c of Ref. [122] where both muon and electron are considered. If muon-exclusive data become available, the signal-to-background ratio will be amplified, yielding improved sensitivity. We evaluated the Run 2 exclusion region from the multi-lepton search, which is depicted by the purple shaded region in Fig. 6. Compared to the previous study [48], the lower mass limit is increased by about 200 GeV.

The future sensitivity is estimated by assuming the significance scales as the square

^{#8} $m_A = m_{H^\pm} = m_H + 90 \text{ GeV}$ is assumed to derive the bound.




Figure 6. The λ_5 - m_A parameter plane in the μ 2HDM. The black contours correspond to the value of $\rho_e^{\mu\mu}$ that explains the muon $g - 2$ anomaly at the 0σ (upper), -1σ (middle), and -2σ (bottom) levels. The orange regions violate the perturbative unitarity bound. The purple regions are excluded by the high- p_T multi-lepton searches based on our simulation. The dashed and dotted purple lines correspond to the extended future prospect with an integrated luminosity of 500fb^{-1} and 3ab^{-1} . The solid and dashed colored (blue, green, and red) lines in each panel correspond to the parameter regions where the cutoff scale are 5 TeV and 30 TeV, respectively.




Figure 7. Representative Feynman diagrams for the collider searches in the μ 2HDM (top-left) and $\mu\tau$ 2HDM (the others).

root of the luminosity.^{#9} We point out that better sensitivity would be obtained with larger $L_T + \text{MET}$ bins; however, this would entail a more complicated experimental analysis. Therefore, our procedure gives a conservative estimate regarding sensitivity. We note that smuon searches at the LHC give less stringent constraints [124, 125].

Furthermore, the Landau poles Λ_{LP} are shown as coloured contours in Fig. 6, where we use solid and dashed lines to illustrate poles at $\Lambda_{\text{LP}} = 5 \text{ TeV}$ and 30 TeV . There is still a small region that can explain the central value of the muon $g - 2$ anomaly if one requires the theory to be perturbative at 5 TeV . On the other hand, if $\Lambda_{\text{LP}} \geq 30 \text{ TeV}$ is required, the model cannot explain the muon $g - 2$ anomaly in any region of the parameter space.

We stress that a future 500 fb^{-1} data set, which approximately corresponds to the integrated luminosity of the Run 3 full data, can probe the whole 1σ region once we require that perturbative theory holds at least up to 5 TeV . The current 2σ region can be covered with a data set of 3 ab^{-1} .

3.4 $\mu\tau$ -flavor violating 2HDM

In the $\mu\tau$ 2HDM, the dominant contribution to a_μ^{NP} arises from the one-loop diagram involving a scalar and a tau lepton, see Fig. 1 (right). The contribution receives an

^{#9}Since the high- p_T lepton signal region is currently statistically limited, this treatment is justified.

m_τ/m_μ enhancement factor compared to the μ 2HDM, due to the chirality flip on the internal tau propagator. Thanks to this enhancement, heavy scalars can serve as an explanation for the muon $g - 2$ anomaly, in contrast to the μ 2HDM.

Again the mass difference $m_H - m_A$ needs to be large to explain the muon $g - 2$, see Appendix A.1. Note that the mass difference is determined by λ_5 in the Higgs alignment limit, see Eq. (2.9). This implies large scalar couplings in the Higgs potential, and thus the RGEs become important, see Appendices A.5 and A.6 for details. Once we require that the model remains perturbative up to 30 TeV (5 TeV), we obtain an upper limit on the scalar mass scale of 1250 GeV (1650 GeV).

For this model, direct searches at the LHC can effectively put constraints on the available parameter space. The additional scalars are quark-phobic (see Eq. (2.18)) and thus the main production mechanism at hadron colliders is EW pair production. The heavy neutral scalar dominantly decays into $\mu\tau$. The unique double $\mu\tau$ -flavor violating resonances can result in two same-sign muons and two oppositely charged same-sign tau leptons in the final state, shown in Fig. 7 (top-right), while the SM background can be neglected to a good approximation. There is so far no experimental analysis for this channel, however theoretical sensitivity studies are available [126]. In these studies, single-scalar production diagrams via Yukawa interactions are taken into account, shown in Fig. 7 (bottom diagrams), in addition to EW pair production, enhancing the sensitivity in the scenario of large masses.

The black contour in Fig. 8 shows the value of $\rho_e^{\mu\tau}\rho_e^{\tau\mu}$ required to explain the central value of the muon $g - 2$ anomaly with the assumption of $|\rho_e^{\mu\tau}| = |\rho_e^{\tau\mu}|$. For given masses $m_{H,A}$, the NP effect in a_μ is large if both couplings $\rho_e^{\mu\tau}$ and $\rho_e^{\tau\mu}$ are large while the Landau pole resides at a high-energy scale. As we are interested in the heaviest scenario, we thus set $|\rho_e^{\mu\tau}| = |\rho_e^{\tau\mu}|$. The τ decay constraint, which mainly comes from the charged-scalar tree-level correction, is depicted by the blue region. It is known that the systematic uncertainty is already the dominant one [127]. The Belle II experiment will improve the sensitivity of the τ decays in the future while its detailed prospect is not available. It should be added that the constraint from Z -boson decays is weaker in this scenario [55].

The dashed magenta contours in Fig. 8 show the HL-LHC reach, where the region to the left of the contour can be probed. A future high energy (HE)-LHC taking data at $\sqrt{s} = 27$ TeV with 3 ab^{-1} can cover the complete region in the plane depicted in Fig. 8, see Ref. [126] for more details. Since the main production mechanism is via EW processes which are insensitive to $\rho_e^{\mu\tau}$ and $\rho_e^{\tau\mu}$, the result does not change drastically even if the anomaly needs to be explained at the -1σ level.

We note that decays into gauge bosons such as $H^+ \rightarrow W^+H$ are not kinematically allowed in the $\mathcal{O}(1)$ TeV scenario since the squared mass difference is $\mathcal{O}(v^2)$, see Eq. (2.9). Even if Δa_μ decreases, the distinctive signal cross section is controlled by the gauge coupling and thus the proposed $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ final state would serve as a smoking gun signal.




Figure 8. The λ_5 - m_A parameter plane in the $\mu\tau$ 2HDM. The black contour corresponds to the value of $\rho_e^{\mu\tau}\rho_e^{\tau\mu}$ that explains $a_\mu^{\text{NP}} = 25.1 \times 10^{-10}$. The cutoff scale is depicted by the purple contour. The blue shaded region is excluded by the tau decay constraint. The dashed magenta line corresponds to the future prospect of the HL-LHC, the region to the left of which will be covered. A future HE-LHC will cover the complete parameter region on the plane. The underlying figure is taken from Ref. [126].

4 Summary and discussion

The current deviation in the muon anomalous magnetic moment could be a long-awaited hint of new physics. In this review article, we revisited the muon $g - 2$ anomaly within the two-Higgs-doublet models. Despite the fact that the 2HDM is one of the simplest extensions of the SM, the model has very rich phenomenology and constitutes the scalar sector of several UV-completing models. In addition to the \mathbb{Z}_2 -based type-X 2HDM, a flavor-aligned 2HDM and the \mathbb{Z}_4 -based muon-specific 2HDM and $\mu\tau$ -flavor violating 2HDM were considered. These models have been known to be a possible solution to the muon $g - 2$ anomaly. We updated the collider constraints which give crucial bounds and clarified the available parameter space.

We found that due to the updated constraint from $B_s \rightarrow \mu^+\mu^-$ the up-type Yukawa coupling in the FA2HDM cannot be large unless the contribution is cancelled by a down-type Yukawa coupling. If we do not rely on this tuning, the Barr-Zee contribution with a top quark cannot explain more than 5% of the discrepancy. Therefore, the scenario is effectively the same as the type-X 2HDM in this case.

Although tau-rich signatures at the LHC provide a distinctive test of the type-X

2HDM and also the FA2HDM interpretation, a detailed Run 2 analysis has not yet been performed for this signature. Based on the latest chargino-neutralino searches with the Run 2 full data, we found that the muon $g - 2$ anomaly favored parameter region is certainly excluded in the type-X 2HDM and FA2HDM. It should be emphasised that even if we employ the cancellation in $B_s \rightarrow \mu^+ \mu^-$, as long as tauonic and bosonic scalar decays are dominant, the interesting region is excluded.

We also revisited the μ 2HDM and found that the Run 2 data pushed up the lower mass bound by 200 GeV compared to the previous analysis and the model encounters the Landau pole at less than 5 TeV if the central value of the muon $g - 2$ anomaly is required. We also found that the complete 1σ region satisfying $\Lambda_{\text{LP}} \geq 5$ TeV can be probed with a near future 500 fb^{-1} data set, which approximately corresponds to the integrated luminosity of the Run 3 full data. The status and prospect of the $\mu\tau$ 2HDM was also summarized. In this model a distinctive $\mu^\pm \mu^\pm \tau^\mp \tau^\mp$ final state at the LHC is a key prediction. We found that the HL-LHC can probe scenarios with scalars of up to 1.1 TeV. Together with the Landau pole constraint, a future upgrade of the HL-LHC to energies of 27 TeV could cover the complete parameter space relevant for the muon $g - 2$ anomaly. The summary of the relevant flavor and collider constraints for those four kinds of 2HDMs is shown in Table 2.

In this review, we focused on a simple extension that features only one additional Higgs doublet. It is known that a further extension, *e.g.*, 2HDM+vector-like lepton, can explain the muon $g - 2$ anomaly (see Ref. [128] for instance), since the heavier vector-like lepton mass can be used to flip the chirality, leading to a much larger mass enhancement factor.^{#10} Further collider searches for the vector-like lepton scenario will shed light on a possible realization of these extended models [129–131].

Acknowledgements

The research of S. I. and M. L. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762-TRR 257. T. K. is supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan, No. 21K03572. The work of T. K. is supported by the Japan Society for the Promotion of Science (JSPS) Core-to-Core Program, No. JPJSCCA20200002. M.T. is supported by the Fundamental Research Funds for the Central Universities, the One Hundred Talent Program of Sun Yat-sen University, China, and by the JSPS KAKENHI Grant, the Grant-in-Aid for Scientific Research C, Grant No. 18K03611.

^{#10}However, it is known that with only vector-like leptons one can explain the muon $g - 2$ anomaly even without introducing an additional scalar doublet.

A Explicit formulae

In this appendix, we collect all relevant formulae required in the analyses of this article.

A.1 Muon $g - 2$

The leading-order Feynman diagrams contributing to possible solutions of the muon $g - 2$ anomaly are shown in Fig. 1.

The one-loop flavor conserving contribution is given as [48]

$$\delta a_\mu^\mu \simeq \frac{G_F v^2 m_\mu^2}{8\sqrt{2}\pi^2} (\rho_e^{\mu\mu})^2 \left[\frac{1}{m_A^2} \left(\frac{11}{6} + \log \frac{m_\mu^2}{m_A^2} \right) - \frac{1}{m_H^2} \left(\frac{7}{6} + \log \frac{m_\mu^2}{m_H^2} \right) - \frac{1}{6m_{H^\pm}^2} \right]. \quad (\text{A.1})$$

The contributions from A and H have opposite sign and H^\pm always gives a negative yet tiny contribution to the muon $g - 2$. A positive shift is realized for $m_H \lesssim m_A$.

Next, the one-loop $\mu\tau$ -flavor violating contribution receives the tau-mass chirality enhancement factor as [55],

$$\delta a_\mu^\tau \simeq \frac{m_\mu^2}{16\pi^2} \rho_e^{\mu\tau} \rho_e^{\tau\mu} \frac{m_\tau}{m_\mu} \left(\frac{\log \frac{m_H^2}{m_\tau^2} - \frac{3}{2}}{m_H^2} - \frac{\log \frac{m_A^2}{m_\tau^2} - \frac{3}{2}}{m_A^2} \right), \quad (\text{A.2})$$

where the H^\pm -loop contribution does not have the tau-mass enhancement and can thus be neglected. Again the contributions from A and H have opposite signs. A positive contribution to the muon $g - 2$ requires $m_H \leq m_A$ for $\rho_e^{\mu\tau} \rho_e^{\tau\mu} > 0$ and $m_H \geq m_A$ for $\rho_e^{\mu\tau} \rho_e^{\tau\mu} < 0$.

Finally, the contribution from the two-loop Barr-Zee diagram is given as [47]

$$\delta a_\mu^{\text{BZ}} \simeq \frac{\alpha m_\mu}{16\pi^3} \left\{ \frac{4\rho_u^{tt} \rho_e^{\mu\mu}}{3m_t} [F_1(x_{tH}) - F_2(x_{tA})] + \frac{\rho_e^{\tau\tau} \rho_e^{\mu\mu}}{m_\tau} [F_1(x_{\tau H}) + F_2(x_{\tau A})] \right\}, \quad (\text{A.3})$$

where $x_{f\phi} = m_f^2/m_\phi^2$ and the loop functions are defined as

$$F_1(x) = x \int_0^1 dy \frac{2y(1-y) - 1}{x - y(1-y)} \log \left[\frac{x}{y(1-y)} \right], \quad (\text{A.4})$$

$$F_2(x) = x \int_0^1 dy \frac{1}{x - y(1-y)} \log \left[\frac{x}{y(1-y)} \right]. \quad (\text{A.5})$$

A.2 Tau decays

The treatment of the constraint arising from tau-lepton decays is crucial in order to judge the type-X 2HDM interpretation. There are five precision observables, the

correlations among which should be taken into account. The HFLAV constraints on the tau effective couplings are summarized as [132],

$$\begin{aligned} \left(\frac{g_\tau}{g_\mu}\right)_\tau &= 1.0009 \pm 0.0014, & \left(\frac{g_\tau}{g_e}\right)_\tau &= 1.00027 \pm 0.0014, \\ \left(\frac{g_\mu}{g_e}\right)_\tau &= 1.0019 \pm 0.0014, & \left(\frac{g_\tau}{g_\mu}\right)_\pi &= 0.9959 \pm 0.0038, \\ \left(\frac{g_\tau}{g_\mu}\right)_K &= 0.9855 \pm 0.0075, \end{aligned} \quad (\text{A.6})$$

where the symmetric correlation matrix is given by

$$\begin{pmatrix} 1 & & & & \\ 0.51 & 1 & & & \\ -0.50 & 0.49 & 1 & & \\ 0.16 & 0.18 & 0.01 & 1 & \\ 0.12 & 0.11 & -0.01 & 0.07 & 1 \end{pmatrix}. \quad (\text{A.7})$$

The effective couplings for leptonic tau decays are defined as

$$\left(\frac{g_\tau}{g_\mu}\right)_\tau \propto \frac{\Gamma(\tau \rightarrow e\nu_\tau\bar{\nu}_e)}{\Gamma(\mu \rightarrow e\nu_\mu\bar{\nu}_e)}, \quad \left(\frac{g_\tau}{g_e}\right)_\tau \propto \frac{\Gamma(\tau \rightarrow \mu\nu_\tau\bar{\nu}_\mu)}{\Gamma(\mu \rightarrow e\nu_\mu\bar{\nu}_e)}, \quad \left(\frac{g_\mu}{g_e}\right)_\tau \propto \frac{\Gamma(\tau \rightarrow \mu\nu_\tau\bar{\nu}_\mu)}{\Gamma(\tau \rightarrow e\nu_\tau\bar{\nu}_e)}, \quad (\text{A.8})$$

while the ones for hadronic tau decays ($h = \pi, K$) are defined as

$$\left(\frac{g_\tau}{g_\mu}\right)_h \propto \frac{\Gamma(\tau \rightarrow h\nu_\tau)}{\Gamma(h \rightarrow \mu\bar{\nu}_\mu)}. \quad (\text{A.9})$$

These ratios are normalized by the phase spaces so that the SM predictions are 1.

Tree-level and one-loop corrections to the tau effective couplings are calculated in Refs. [36, 89]. The contributions are given by

$$\left(\frac{g_\tau}{g_\mu}\right)_{\tau, \pi, K} \simeq 1 + \delta_\tau^{\text{loop}} - \delta_\mu^{\text{loop}}, \quad (\text{A.10})$$

$$\left(\frac{g_\tau}{g_e}\right)_\tau \simeq 1 + \delta_\tau^{\text{tree}} + \delta_\tau^{\text{loop}} - \delta_\mu^{\text{loop}}, \quad (\text{A.11})$$

$$\left(\frac{g_\mu}{g_e}\right)_\tau \simeq 1 + \delta_\tau^{\text{tree}}, \quad (\text{A.12})$$

with

$$\delta_\tau^{\text{tree}} = \frac{1}{2} \left[-\frac{v^2}{m_{H^\pm}^2} \rho_e^{\mu\mu} \rho_e^{\tau\tau} \frac{m_\mu g(m_\mu^2/m_\tau^2)}{m_\tau f(m_\mu^2/m_\tau^2)} + \frac{v^4}{16m_{H^\pm}^4} (\rho_e^{\mu\mu} \rho_e^{\tau\tau})^2 \right], \quad (\text{A.13})$$

$$\delta_\ell^{\text{loop}} = \frac{(\rho_e^{\ell\ell})^2}{32\pi^2} \left\{ 1 + \frac{1}{4} \left[h \left(\frac{m_A^2}{m_{H^\pm}^2} \right) + h \left(\frac{m_H^2}{m_{H^\pm}^2} \right) \right] \right\}. \quad (\text{A.14})$$

A contribution from the tree-level H^\pm exchange in $\tau \rightarrow \mu\nu_\tau\bar{\nu}_\mu$ is represented by $\delta_\tau^{\text{tree}}$, while radiative corrections to the W - ℓ - ν_ℓ couplings are denoted by $\delta_\ell^{\text{loop}}$. The loop functions are given by

$$f(x) = 1 - 8x + 8x^3 - x^4 - 12x^2 \log x, \quad (\text{A.15})$$

$$g(x) = 1 + 9x - 9x^2 - x^3 + 6x(1+x) \log x, \quad (\text{A.16})$$

$$h(x) = \frac{1+x}{1-x} \log x. \quad (\text{A.17})$$

To investigate the exclusion region from tau decays, the χ^2 is constructed based on these five observables including their correlations. For the $\mu\tau$ 2HDM, the tau decay constraints are less significant due to the heavy additional scalar masses, even if one includes the Michel parameters [133].

A.3 Z decays

Sizable lepton Yukawa couplings change the Z -boson vertices at the one-loop level [36, 40, 59]. The corrections to the ratios of the leptonic decay widths are given by

$$\frac{\Gamma(Z \rightarrow \tau^+\tau^-)}{\Gamma(Z \rightarrow e^+e^-)} \simeq 1 + 2 \frac{g_L^e \text{Re}(\delta g_L) + g_R^e \text{Re}(\delta g_R)}{(g_L^e)^2 + (g_R^e)^2}, \quad (\text{A.18})$$

$$\frac{\Gamma(Z \rightarrow \mu^+\mu^-)}{\Gamma(Z \rightarrow e^+e^-)} \simeq 1 + 2 \frac{g_L^e \text{Re}(\delta g_L) + g_R^e \text{Re}(\delta g_R)}{(g_L^e)^2 + (g_R^e)^2} \left(\frac{\rho_e^{\mu\mu}}{\rho_e^{\tau\tau}} \right)^2, \quad (\text{A.19})$$

with the vertex corrections

$$\delta g_L = \frac{(\rho_e^{\tau\tau})^2}{32\pi^2} \left\{ -\frac{1}{2} [B_Z(x_A) + B_Z(x_H) + 4C_Z(x_A, x_H)] \right. \\ \left. + s_W^2 [B_Z(x_A) + B_Z(x_H) + \tilde{C}_Z(x_A) + \tilde{C}_Z(x_H)] \right\}, \quad (\text{A.20})$$

$$\delta g_R = \frac{(\rho_e^{\tau\tau})^2}{32\pi^2} \left\{ \frac{1}{2} [4C_Z(x_A, x_H) - 4C_Z(x_{H^\pm}, x_{H^\pm}) + 2\tilde{C}_Z(x_{H^\pm}) - \tilde{C}_Z(x_A) - \tilde{C}_Z(x_H)] \right. \\ \left. + s_W^2 [B_Z(x_A) + B_Z(x_H) + 2B_Z(x_{H^\pm}) + 4C_Z(x_{H^\pm}, x_{H^\pm}) + \tilde{C}_Z(x_A) + \tilde{C}_Z(x_H)] \right\}, \quad (\text{A.21})$$

where $x_\phi = m_\phi^2/m_Z^2$, $s_W \equiv \sin \theta_W$, and $g_{L,R} = T_3 - Qs_W^2$ ($g_L^e \simeq -0.27$ and $g_R^e \simeq 0.23$). The loop functions are defined as

$$B_Z(x) = -\frac{1}{2\bar{\epsilon}} - \frac{1}{4} + \frac{1}{2} \log x, \quad (\text{A.22})$$

$$C_Z(x, y) = \frac{1}{4\bar{\epsilon}} - \frac{1}{2} \int_0^1 dz_1 \int_0^{z_1} dz_2 \log [z_1 z_2 + y(1-z_1) + (x-1)z_2], \quad (\text{A.23})$$

$$\tilde{C}_Z(x) = \frac{1}{2\bar{\epsilon}} + \frac{1}{2} - x(1 + \log x) + x^2 [\log x \log(1+x^{-1}) - \text{Li}_2(-x^{-1})] \\ - \frac{i\pi}{2} [1 - 2x + 2x^2 \log(1+x^{-1})], \quad (\text{A.24})$$




Figure 9. The dominant Feynman diagram for $B_s \rightarrow \mu^+ \mu^-$ in the type-X 2HDM and FA2HDM.

where Li_2 denotes the di-logarithm function and the $1/\bar{\epsilon}$ poles cancel in the sum. We confirm that the forward-backward asymmetry and the tau polarization asymmetry in $Z \rightarrow \tau^+ \tau^-$ give less stringent constraints.

For $\mu\tau$ 2HDM, $(\rho_e^{\tau\tau})^2$ is replaced by $(\rho_e^{\mu\tau})^2$ in δg_L and by $(\rho_e^{\tau\mu})^2$ in δg_R , respectively [59]. For $|\rho_e^{\mu\tau}| = |\rho_e^{\tau\mu}|$ the constraint is less stringent.

A.4 $B_s \rightarrow \mu^+ \mu^-$

In this section, we discuss the constraint from $B_s \rightarrow \mu^+ \mu^-$. In Ref. [134], the calculation of the full one-loop Wilson coefficients contributing to $B_s \rightarrow \mu^+ \mu^-$ has been performed within the FA2HDM. The recent CMS result [99] is consistent with the SM prediction [94–96]. Since the type-X 2HDM increases the branching ratio of $B_s \rightarrow \mu^+ \mu^-$, the recent shift of the experimental world average [100] weakens the m_A bound stemming from $B_s \rightarrow \mu^+ \mu^-$ compared to the previous world average [97, 98]. This bound is relevant for the type-X 2HDM and FA2HDM, since the dominant contribution comes from the one-loop diagram with the light CP-odd scalar mediation, shown in Fig. 9. We adopted the formulae from Ref. [134] and derived the constraint.

Since $\rho_e^{\tau\tau}$ is larger than $\rho_e^{\mu\mu}$, one might think that $B_s \rightarrow \tau^+ \tau^-$ could be a good decay process in order to probe the 2HDMs. However, this does not hold and the NP sensitivity would be the same as $B_s \rightarrow \mu^+ \mu^-$, because the SM amplitude (W -box and Z -penguin diagrams) is also proportional to m_τ . Moreover, $B_s \rightarrow \tau^+ \tau^-$ has not yet been observed.

A.5 Perturbative unitarity and vacuum stability

In this section we discuss theoretical constraints imposed on the couplings in the scalar potential by the requirement of perturbative unitarity and the vacuum stability. The constraints from perturbativity can be derived from the consideration of scattering amplitudes of the Higgs bosons. Following Ref. [135] where longitudinally polarized gauge bosons are replaced with the corresponding Nambu-Goldstone bosons, we only

consider the scattering processes involving scalars and gauge bosons. The full set of scattering amplitudes is expressed as a 22×22 matrix, which falls apart into four decoupled sub-matrices [36, 136–141]. The perturbative unitarity bound is imposed on the 12 distinct eigenvalues of the matrix as

$$|e_j| < 8\pi \quad (j = 1, \dots, 12), \quad (\text{A.25})$$

where

$$e_{1,2} = \lambda_3 \pm \lambda_4, \quad e_{3,4} = \lambda_3 \pm \lambda_5, \quad e_{5,6} = \lambda_3 + 2\lambda_4 \pm 3\lambda_5, \quad (\text{A.26})$$

$$e_{7,8} = \frac{1}{2} \left[(\lambda_1 + \lambda_2) \pm \sqrt{(\lambda_1 - \lambda_2)^2 + 4\lambda_4^2} \right], \quad (\text{A.27})$$

$$e_{9,10} = \frac{1}{2} \left[3(\lambda_1 + \lambda_2) \pm \sqrt{9(\lambda_1 - \lambda_2)^2 + 4(2\lambda_3 + \lambda_4)^2} \right], \quad (\text{A.28})$$

$$e_{11,12} = \frac{1}{2} \left[(\lambda_1 + \lambda_2) \pm \sqrt{(\lambda_1 - \lambda_2)^2 + 4\lambda_5^2} \right], \quad (\text{A.29})$$

where all λ_j are running couplings. Here, the contributions from λ_6 and λ_7 are discarded; $\lambda_6 = 0$ is fixed by the Higgs alignment condition and λ_7 is suppressed by large $\tan\beta$, see Eq. (2.7). We define our cutoff scale Λ_{LP} as the minimum scale at which either the vacuum stability or perturbative unitarity condition breaks down when evolving the couplings with the RGEs from an input scale to a high-energy scale. It should be noted that the quartic couplings are also bounded from below by the conditions [142, 143],

$$\lambda_1, \lambda_2 \geq 0, \quad \sqrt{\lambda_1\lambda_2} + \lambda_3 \geq 0, \quad \sqrt{\lambda_1\lambda_2} + \lambda_3 + \lambda_4 - |\lambda_5| \geq 0, \quad (\text{A.30})$$

but these conditions are always satisfied in the parameter region of our interest.

A.6 Renormalization group equations

The RGEs of the scalar quartic couplings in the μ 2HDM and $\mu\tau$ 2HDM are given in the form of

$$\frac{d\lambda_j}{d\log\mu} = \frac{\beta_{\lambda_j}}{(4\pi)^2}, \quad (\text{A.31})$$

where μ is the renormalization scale. The RGE running effect is important for these two 2HDMs since the scalars are heavy and thus the Yukawa couplings are large.

At the one-loop level, the beta functions β_{λ_j} of the scalar potential in Eq. (2.5)

are given by [143, 144]

$$\begin{aligned} \beta_{\lambda_1} = & 12\lambda_1^2 + 4\lambda_3^2 + 4\lambda_3\lambda_4 + 2\lambda_4^2 + 2\lambda_5^2 + 24\lambda_6^2 + \frac{3}{4}(3g^4 + g'^4 + 2g^2g'^2) \\ & - 3\lambda_1(3g^2 + g'^2) + 12\lambda_1y_t^2 - 12y_t^4, \end{aligned} \quad (\text{A.32})$$

$$\begin{aligned} \beta_{\lambda_2} = & 12\lambda_2^2 + 4\lambda_3^2 + 4\lambda_3\lambda_4 + 2\lambda_4^2 + 2\lambda_5^2 + 24\lambda_7^2 + \frac{3}{4}(3g^4 + g'^4 + 2g^2g'^2) - 3\lambda_2(3g^2 + g'^2) \\ & + 4\lambda_2[(\rho_e^{\ell\ell})^2 + (\rho_e^{\mu\tau})^2 + (\rho_e^{\tau\mu})^2] - 4[(\rho_e^{\ell\ell})^4 + (\rho_e^{\mu\tau})^4 + (\rho_e^{\tau\mu})^4], \end{aligned} \quad (\text{A.33})$$

$$\begin{aligned} \beta_{\lambda_3} = & 2(\lambda_1 + \lambda_2)(3\lambda_3 + \lambda_4) + 4\lambda_3^2 + 2\lambda_4^2 + 2\lambda_5^2 + 4\lambda_6^2 + 16\lambda_6\lambda_7 + 4\lambda_7^2 - 3\lambda_3(3g^2 + g'^2) \\ & + \frac{3}{4}(3g^4 + g'^4 - 2g^2g'^2) + 2\lambda_3[3y_t^2 + (\rho_e^{\ell\ell})^2 + (\rho_e^{\tau\mu})^2 + (\rho_e^{\mu\tau})^2], \end{aligned} \quad (\text{A.34})$$

$$\begin{aligned} \beta_{\lambda_4} = & 2\lambda_4(\lambda_1 + \lambda_2 + 4\lambda_3 + 2\lambda_4) + 8\lambda_5^2 + 10\lambda_6^2 + 4\lambda_6\lambda_7 + 10\lambda_7^2 + 3g^2g'^2 \\ & - 3\lambda_4(3g^2 + g'^2) + 2\lambda_4[3y_t^2 + (\rho_e^{\ell\ell})^2 + (\rho_e^{\tau\mu})^2 + (\rho_e^{\mu\tau})^2], \end{aligned} \quad (\text{A.35})$$

$$\begin{aligned} \beta_{\lambda_5} = & 2\lambda_5(\lambda_1 + \lambda_2 + 4\lambda_3 + 6\lambda_4) + 10\lambda_6^2 + 4\lambda_6\lambda_7 + 10\lambda_7^2 - 3\lambda_5(3g^2 + g'^2) \\ & + 2\lambda_5[3y_t^2 + (\rho_e^{\ell\ell})^2 + (\rho_e^{\tau\mu})^2 + (\rho_e^{\mu\tau})^2], \end{aligned} \quad (\text{A.36})$$

$$\begin{aligned} \beta_{\lambda_6} = & 12\lambda_1\lambda_6 + 6\lambda_3(\lambda_6 + \lambda_7) + 8\lambda_4\lambda_6 + 4\lambda_4\lambda_7 + 10\lambda_5\lambda_6 + 2\lambda_5\lambda_7 - 3\lambda_6(3g^2 + g'^2) \\ & + 3\lambda_6[3y_t^2 + (\rho_e^{\ell\ell})^2 + (\rho_e^{\tau\mu})^2 + (\rho_e^{\mu\tau})^2], \end{aligned} \quad (\text{A.37})$$

$$\begin{aligned} \beta_{\lambda_7} = & 12\lambda_2\lambda_7 + 6\lambda_3(\lambda_6 + \lambda_7) + 4\lambda_4\lambda_6 + 8\lambda_4\lambda_7 + 2\lambda_5\lambda_6 + 10\lambda_5\lambda_7 - 3\lambda_7(3g^2 + g'^2) \\ & + \lambda_7[3y_t^2 + (\rho_e^{\ell\ell})^2 + (\rho_e^{\tau\mu})^2 + (\rho_e^{\mu\tau})^2], \end{aligned} \quad (\text{A.38})$$

where the Yukawa couplings are defined by $y_f = \sqrt{2}m_f/v$. The RGEs of the gauge and Yukawa couplings are defined in the same way, with the beta functions given by

$$\beta_{g_j} = b_j g_j^3, \quad b_j = \{7, -3, -7\} \quad (g_j = \{g', g, g_s\}), \quad (\text{A.39})$$

$$\beta_{y_t} = y_t \left(-\frac{17}{12}g'^2 - \frac{9}{4}g^2 - 8g_s^2 + \frac{9}{2}y_t^2 \right), \quad (\text{A.40})$$

$$\beta_{\rho_e^{\ell\ell}} = \rho_e^{\ell\ell} \left[\frac{5}{2}(\rho_e^{\ell\ell})^2 - \frac{9}{4} \left(\frac{5}{3}g'^2 + g^2 \right) \right], \quad (\text{A.41})$$

$$\beta_{\rho_e^{\tau\mu}} = \rho_e^{\tau\mu} \left[(\rho_e^{\mu\tau})^2 + \frac{5}{2}(\rho_e^{\tau\mu})^2 - \frac{9}{4} \left(\frac{5}{3}g'^2 + g^2 \right) \right], \quad (\text{A.42})$$

$$\beta_{\rho_e^{\mu\tau}} = \rho_e^{\mu\tau} \left[(\rho_e^{\tau\mu})^2 + \frac{5}{2}(\rho_e^{\mu\tau})^2 - \frac{9}{4} \left(\frac{5}{3}g'^2 + g^2 \right) \right]. \quad (\text{A.43})$$

References

- [1] T. Aoyama *et al.*, “The anomalous magnetic moment of the muon in the Standard Model,” *Phys. Rept.* **887** (2020) 1–166 [[arXiv:2006.04822](#)].
- [2] **Muon g-2** Collaboration, “Measurement of the positive muon anomalous magnetic moment to 0.7 ppm,” *Phys. Rev. Lett.* **89** (2002) 101804 [[hep-ex/0208001](#)]. [Erratum: *Phys.Rev.Lett.* 89, 129903 (2002)].

- [3] **Muon g-2** Collaboration, “Measurement of the negative muon anomalous magnetic moment to 0.7 ppm,” *Phys. Rev. Lett.* **92** (2004) 161802 [[hep-ex/0401008](#)].
- [4] **Muon g-2** Collaboration, “Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL,” *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)].
- [5] **Muon g-2** Collaboration, “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm,” *Phys. Rev. Lett.* **126** (2021) 141801 [[arXiv:2104.03281](#)].
- [6] G. Lafferty and S. Soldner-Rembold, eds., “Measurement of muon g-2 and EDM with an ultra-cold muon beam at J-PARC,” *Nucl. Phys. B Proc. Suppl.* **218** (2011) 242–246.
- [7] M. Abe *et al.*, “A New Approach for Measuring the Muon Anomalous Magnetic Moment and Electric Dipole Moment,” *PTEP* **2019** (2019) 053C02 [[arXiv:1901.03047](#)].
- [8] S. Borsanyi *et al.*, “Leading hadronic contribution to the muon magnetic moment from lattice QCD,” *Nature* **593** (2021) 51–55 [[arXiv:2002.12347](#)].
- [9] M. Cè *et al.*, “Window observable for the hadronic vacuum polarization contribution to the muon g-2 from lattice QCD,” *Phys. Rev. D* **106** (2022) 114502 [[arXiv:2206.06582](#)].
- [10] C. Alexandrou *et al.*, “Lattice calculation of the short and intermediate time-distance hadronic vacuum polarization contributions to the muon magnetic moment using twisted-mass fermions.” [arXiv:2206.15084](#).
- [11] A. Bazavov *et al.*, “Light-quark connected intermediate-window contributions to the muon $g - 2$ hadronic vacuum polarization from lattice QCD.” [arXiv:2301.08274](#).
- [12] T. Blum *et al.*, “An update of Euclidean windows of the hadronic vacuum polarization.” [arXiv:2301.08696](#).
- [13] **RBC, UKQCD** Collaboration, “Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment,” *Phys. Rev. Lett.* **121** (2018) 022003 [[arXiv:1801.07224](#)].
- [14] H. Wittig, “Progress on $(g - 2)_\mu$ from Lattice QCD.” Presentation at the 57th Rencontres de Moriond EW 2023, 21 March 2023.

- [15] A. Crivellin, M. Hoferichter, C. A. Manzari, and M. Montull, “Hadronic Vacuum Polarization: $(g - 2)_\mu$ versus Global Electroweak Fits,” *Phys. Rev. Lett.* **125** (2020) 091801 [[arXiv:2003.04886](#)].
- [16] A. Keshavarzi, W. J. Marciano, M. Passera, and A. Sirlin, “Muon $g - 2$ and $\Delta\alpha$ connection,” *Phys. Rev. D* **102** (2020) 033002 [[arXiv:2006.12666](#)].
- [17] G. Colangelo, M. Hoferichter, and P. Stoffer, “Constraints on the two-pion contribution to hadronic vacuum polarization,” *Phys. Lett. B* **814** (2021) 136073 [[arXiv:2010.07943](#)].
- [18] **CMD-3** Collaboration, “Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section from threshold to 1.2 GeV with the CMD-3 detector.” [arXiv:2302.08834](#).
- [19] **KLOE-2** Collaboration, “Combination of KLOE $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ measurements and determination of $a_\mu^{\pi^+\pi^-}$ in the energy range $0.10 < s < 0.95$ GeV²,” *JHEP* **03** (2018) 173 [[arXiv:1711.03085](#)].
- [20] **BaBar** Collaboration, “Precise Measurement of the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ Cross Section with the Initial-State Radiation Method at BABAR,” *Phys. Rev. D* **86** (2012) 032013 [[arXiv:1205.2228](#)].
- [21] **CMD-2** Collaboration, “High-statistics measurement of the pion form factor in the rho-meson energy range with the CMD-2 detector,” *Phys. Lett. B* **648** (2007) 28–38 [[hep-ex/0610021](#)].
- [22] P. Athron, *et al.*, “New physics explanations of a_μ in light of the FNAL muon $g - 2$ measurement,” *JHEP* **09** (2021) 080 [[arXiv:2104.03691](#)].
- [23] S. L. Glashow and S. Weinberg, “Natural Conservation Laws for Neutral Currents,” *Phys. Rev. D* **15** (1977) 1958.
- [24] G. L. Kane, ed., “A Supersymmetry primer,” *Adv. Ser. Direct. High Energy Phys.* **18** (1998) 1–98 [[hep-ph/9709356](#)].
- [25] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, “Towards high precision predictions for the MSSM Higgs sector,” *Eur. Phys. J. C* **28** (2003) 133–143 [[hep-ph/0212020](#)].
- [26] R. N. Mohapatra and J. C. Pati, “A Natural Left-Right Symmetry,” *Phys. Rev. D* **11** (1975) 2558.
- [27] J. C. Pati and A. Salam, “Lepton Number as the Fourth Color,” *Phys. Rev. D* **10** (1974) 275–289. [Erratum: *Phys.Rev.D* 11, 703–703 (1975)].

- [28] N. Arkani-Hamed, A. G. Cohen, E. Katz, and A. E. Nelson, “The Littlest Higgs,” *JHEP* **07** (2002) 034 [[hep-ph/0206021](#)].
- [29] I. Low, W. Skiba, and D. Tucker-Smith, “Little Higgses from an antisymmetric condensate,” *Phys. Rev. D* **66** (2002) 072001 [[hep-ph/0207243](#)].
- [30] P. Ko, Y. Omura, and C. Yu, “A Resolution of the Flavor Problem of Two Higgs Doublet Models with an Extra $U(1)_H$ Symmetry for Higgs Flavor,” *Phys. Lett. B* **717** (2012) 202–206 [[arXiv:1204.4588](#)].
- [31] S. Banik, A. Crivellin, S. Iguro, and T. Kitahara, “Asymmetric Di-Higgs Signals of the N2HDM- $U(1)$.” [arXiv:2303.11351](#).
- [32] V. D. Barger, J. L. Hewett, and R. J. N. Phillips, “New Constraints on the Charged Higgs Sector in Two Higgs Doublet Models,” *Phys. Rev. D* **41** (1990) 3421–3441.
- [33] Y. Grossman, “Phenomenology of models with more than two Higgs doublets,” *Nucl. Phys. B* **426** (1994) 355–384 [[hep-ph/9401311](#)].
- [34] A. Broggio, E. J. Chun, M. Passera, K. M. Patel, and S. K. Vempati, “Limiting two-Higgs-doublet models,” *JHEP* **11** (2014) 058 [[arXiv:1409.3199](#)].
- [35] L. Wang and X.-F. Han, “A light pseudoscalar of 2HDM confronted with muon $g - 2$ and experimental constraints,” *JHEP* **05** (2015) 039 [[arXiv:1412.4874](#)].
- [36] T. Abe, R. Sato, and K. Yagyu, “Lepton-specific two Higgs doublet model as a solution of muon $g - 2$ anomaly,” *JHEP* **07** (2015) 064 [[arXiv:1504.07059](#)].
- [37] A. Hektor, K. Kannike, and L. Marzola, “Muon $g - 2$ and Galactic Centre γ -ray excess in a scalar extension of the 2HDM type-X,” *JCAP* **10** (2015) 025 [[arXiv:1507.05096](#)].
- [38] E. J. Chun, Z. Kang, M. Takeuchi, and Y.-L. S. Tsai, “LHC τ -rich tests of lepton-specific 2HDM for $(g - 2)_\mu$,” *JHEP* **11** (2015) 099 [[arXiv:1507.08067](#)].
- [39] T. Han, S. K. Kang, and J. Sayre, “Muon $g - 2$ in the aligned two Higgs doublet model,” *JHEP* **02** (2016) 097 [[arXiv:1511.05162](#)].
- [40] E. J. Chun and J. Kim, “Leptonic Precision Test of Leptophilic Two-Higgs-Doublet Model,” *JHEP* **07** (2016) 110 [[arXiv:1605.06298](#)].
- [41] A. Cherchiglia, D. Stöckinger, and H. Stöckinger-Kim, “Muon $g-2$ in the 2HDM: maximum results and detailed phenomenology,” *Phys. Rev. D* **98** (2018) 035001 [[arXiv:1711.11567](#)].

- [42] L. Wang, J. M. Yang, M. Zhang, and Y. Zhang, “Revisiting lepton-specific 2HDM in light of muon $g - 2$ anomaly,” *Phys. Lett. B* **788** (2019) 519–529 [[arXiv:1809.05857](#)].
- [43] E. J. Chun, J. Kim, and T. Mondal, “Electron EDM and Muon anomalous magnetic moment in Two-Higgs-Doublet Models,” *JHEP* **12** (2019) 068 [[arXiv:1906.00612](#)].
- [44] A. Jueid, J. Kim, S. Lee, and J. Song, “Type-X two-Higgs-doublet model in light of the muon $g-2$: Confronting Higgs boson and collider data,” *Phys. Rev. D* **104** (2021) 095008 [[arXiv:2104.10175](#)].
- [45] P. M. Ferreira, B. L. Gonçalves, F. R. Joaquim, and M. Sher, “ $(g-2)_\mu$ in the 2HDM and slightly beyond: An updated view,” *Phys. Rev. D* **104** (2021) 053008 [[arXiv:2104.03367](#)].
- [46] A. Pich and P. Tuzon, “Yukawa Alignment in the Two-Higgs-Doublet Model,” *Phys. Rev. D* **80** (2009) 091702 [[arXiv:0908.1554](#)].
- [47] V. Ilisie, “New Barr-Zee contributions to $(g - 2)_\mu$ in two-Higgs-doublet models,” *JHEP* **04** (2015) 077 [[arXiv:1502.04199](#)].
- [48] T. Abe, R. Sato, and K. Yagyu, “Muon specific two-Higgs-doublet model,” *JHEP* **07** (2017) 012 [[arXiv:1705.01469](#)].
- [49] S. Nie and M. Sher, “The Anomalous magnetic moment of the muon and Higgs mediated flavor changing neutral currents,” *Phys. Rev. D* **58** (1998) 097701 [[hep-ph/9805376](#)].
- [50] R. A. Diaz, R. Martinez, and J. A. Rodriguez, “Bounds for lepton flavor violation and the pseudoscalar Higgs boson in the general two Higgs doublet model using the $g - 2$ muon factor,” *Phys. Rev. D* **64** (2001) 033004 [[hep-ph/0010339](#)].
- [51] S. Baek, N. G. Deshpande, X. G. He, and P. Ko, “Muon anomalous $g - 2$ and gauged $L_\mu - L_\tau$ models,” *Phys. Rev. D* **64** (2001) 055006 [[hep-ph/0104141](#)].
- [52] E. O. Iltan and H. Sundu, “Anomalous magnetic moment of muon in the general two Higgs doublet model,” *Acta Phys. Slov.* **53** (2003) 17 [[hep-ph/0103105](#)].
- [53] Y.-L. Wu and Y.-F. Zhou, “Muon anomalous magnetic moment in the standard model with two Higgs doublets,” *Phys. Rev. D* **64** (2001) 115018 [[hep-ph/0104056](#)].

- [54] K. A. Assamagan, A. Deandrea, and P.-A. Delsart, “Search for the lepton flavor violating decay $A^0/H^0 \rightarrow \tau^\pm \mu^\mp$ at hadron colliders,” *Phys. Rev. D* **67** (2003) 035001 [[hep-ph/0207302](#)].
- [55] Y. Omura, E. Senaha, and K. Tobe, “Lepton-flavor-violating Higgs decay $h \rightarrow \mu\tau$ and muon anomalous magnetic moment in a general two Higgs doublet model,” *JHEP* **05** (2015) 028 [[arXiv:1502.07824](#)].
- [56] Y. Omura, E. Senaha, and K. Tobe, “ τ - and μ -physics in a general two Higgs doublet model with $\mu - \tau$ flavor violation,” *Phys. Rev. D* **94** (2016) 055019 [[arXiv:1511.08880](#)].
- [57] P. S. B. Dev, R. N. Mohapatra, and Y. Zhang, “Lepton Flavor Violation Induced by a Neutral Scalar at Future Lepton Colliders,” *Phys. Rev. Lett.* **120** (2018) 221804 [[arXiv:1711.08430](#)].
- [58] S. Iguro and Y. Omura, “Status of the semileptonic B decays and muon $g-2$ in general 2HDMs with right-handed neutrinos,” *JHEP* **05** (2018) 173 [[arXiv:1802.01732](#)].
- [59] Y. Abe, T. Toma, and K. Tsumura, “A μ - τ -philic scalar doublet under Z_n flavor symmetry,” *JHEP* **06** (2019) 142 [[arXiv:1904.10908](#)].
- [60] L. Wang and Y. Zhang, “ μ - τ -philic Higgs doublet model confronted with the muon $g-2$, τ decays, and LHC data,” *Phys. Rev. D* **100** (2019) 095005 [[arXiv:1908.03755](#)].
- [61] S. Iguro, Y. Omura, and M. Takeuchi, “Testing the 2HDM explanation of the muon $g - 2$ anomaly at the LHC,” *JHEP* **11** (2019) 130 [[arXiv:1907.09845](#)].
- [62] A. Crivellin, D. Müller, and C. Wiegand, “ $b \rightarrow s\ell^+\ell^-$ transitions in two-Higgs-doublet models,” *JHEP* **06** (2019) 119 [[arXiv:1903.10440](#)].
- [63] S. Iguro, Y. Omura, and M. Takeuchi, “Probing $\mu\tau$ flavor-violating solutions for the muon $g - 2$ anomaly at Belle II,” *JHEP* **09** (2020) 144 [[arXiv:2002.12728](#)].
- [64] S. Jana, V. P. K., and S. Saad, “Resolving electron and muon $g - 2$ within the 2HDM,” *Phys. Rev. D* **101** (2020) 115037 [[arXiv:2003.03386](#)].
- [65] H.-X. Wang, L. Wang, and Y. Zhang, “Muon $g - 2$ anomaly and μ - τ -philic Higgs doublet with a light CP-even component,” *Eur. Phys. J. C* **81** (2021) 1007 [[arXiv:2104.03242](#)].

- [66] N. Ghosh and J. Lahiri, “Generalized 2HDM with wrong-sign lepton-Yukawa coupling, in light of $g_\mu - 2$ and lepton flavor violation at the future LHC,” *Eur. Phys. J. C* **81** (2021) 1074 [[arXiv:2103.10632](#)].
- [67] W.-S. Hou, R. Jain, C. Kao, G. Kumar, and T. Modak, “Collider Prospects for Muon $g - 2$ in General Two Higgs Doublet Model,” *Phys. Rev. D* **104** (2021) 075036 [[arXiv:2105.11315](#)].
- [68] X.-F. Han, F. Wang, L. Wang, J. M. Yang, and Y. Zhang, “Joint explanation of W-mass and muon $g-2$ in the 2HDM*,” *Chin. Phys. C* **46** (2022) 103105 [[arXiv:2204.06505](#)].
- [69] K. Asai, C. Miyao, S. Okawa, and K. Tsumura, “Scalar dark matter with a $\mu\tau$ flavored mediator,” *Phys. Rev. D* **106** (2022) 035017 [[arXiv:2205.08998](#)].
- [70] M. Aiko and S. Kanemura, “New scenario for aligned Higgs couplings originated from the twisted custodial symmetry at high energies,” *JHEP* **02** (2021) 046 [[arXiv:2009.04330](#)].
- [71] S. Iguro, T. Kitahara, Y. Omura, and H. Zhang, “Chasing the two-Higgs doublet model in the di-Higgs production,” *Phys. Rev. D* **107** (2023) 075017 [[arXiv:2211.00011](#)].
- [72] S. Iguro and K. Tobe, “ $R(D^{(*)})$ in a general two Higgs doublet model,” *Nucl. Phys. B* **925** (2017) 560–606 [[arXiv:1708.06176](#)].
- [73] H. Georgi and D. V. Nanopoulos, “Suppression of Flavor Changing Effects From Neutral Spinless Meson Exchange in Gauge Theories,” *Phys. Lett. B* **82** (1979) 95–96.
- [74] J. F. Donoghue and L. F. Li, “Properties of Charged Higgs Bosons,” *Phys. Rev. D* **19** (1979) 945.
- [75] S. Davidson and H. E. Haber, “Basis-independent methods for the two-Higgs-doublet model,” *Phys. Rev. D* **72** (2005) 035004 [[hep-ph/0504050](#)]. [Erratum: *Phys.Rev.D* 72, 099902 (2005)].
- [76] **CMS** Collaboration, “Search for lepton-flavor violating decays of the Higgs boson in the $\mu\tau$ and $e\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *Phys. Rev. D* **104** (2021) 032013 [[arXiv:2105.03007](#)].
- [77] **ATLAS** Collaboration, “Searches for lepton-flavour-violating decays of the Higgs boson into $e\tau$ and $\mu\tau$ in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector.” [arXiv:2302.05225](#).

- [78] **CMS** Collaboration, “Search for the lepton flavor violating decay of a Higgs boson in the $e\mu$ final state in proton-proton collisions at $\sqrt{s} = 13$ TeV.” <https://cds.cern.ch/record/2851512>.
- [79] M. Aoki, S. Kanemura, K. Tsumura, and K. Yagyu, “Models of Yukawa interaction in the two Higgs doublet model, and their collider phenomenology,” *Phys. Rev. D* **80** (2009) 015017 [[arXiv:0902.4665](https://arxiv.org/abs/0902.4665)].
- [80] S. M. Barr and A. Zee, “Electric Dipole Moment of the Electron and of the Neutron,” *Phys. Rev. Lett.* **65** (1990) 21–24. [Erratum: *Phys.Rev.Lett.* 65, 2920 (1990)].
- [81] M. Misiak, A. Rehman, and M. Steinhauser, “Towards $\overline{B} \rightarrow X_s \gamma$ at the NNLO in QCD without interpolation in m_c ,” *JHEP* **06** (2020) 175 [[arXiv:2002.01548](https://arxiv.org/abs/2002.01548)].
- [82] C. B. Braeuninger, A. Ibarra, and C. Simonetto, “Radiatively induced flavour violation in the general two-Higgs doublet model with Yukawa alignment,” *Phys. Lett. B* **692** (2010) 189–195 [[arXiv:1005.5706](https://arxiv.org/abs/1005.5706)].
- [83] S. Gori, H. E. Haber, and E. Santos, “High scale flavor alignment in two-Higgs doublet models and its phenomenology,” *JHEP* **06** (2017) 110 [[arXiv:1703.05873](https://arxiv.org/abs/1703.05873)].
- [84] S. Knapen and D. J. Robinson, “Disentangling Mass and Mixing Hierarchies,” *Phys. Rev. Lett.* **115** (2015) 161803 [[arXiv:1507.00009](https://arxiv.org/abs/1507.00009)].
- [85] A. Peñuelas and A. Pich, “Flavour alignment in multi-Higgs-doublet models,” *JHEP* **12** (2017) 084 [[arXiv:1710.02040](https://arxiv.org/abs/1710.02040)].
- [86] **CMS** Collaboration, “Measurement of the Higgs boson width and evidence of its off-shell contributions to ZZ production,” *Nature Phys.* **18** (2022) 1329–1334 [[arXiv:2202.06923](https://arxiv.org/abs/2202.06923)].
- [87] **ATLAS** Collaboration, “Evidence of off-shell Higgs boson production from ZZ leptonic decay channels and constraints on its total width with the ATLAS detector.” [arXiv:2304.01532](https://arxiv.org/abs/2304.01532).
- [88] **CMS** Collaboration, “Search for a light pseudoscalar Higgs boson in the boosted $\mu\mu\tau\tau$ final state in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **08** (2020) 139 [[arXiv:2005.08694](https://arxiv.org/abs/2005.08694)].
- [89] M. Krawczyk and D. Temes, “2HDM(II) radiative corrections in leptonic tau decays,” *Eur. Phys. J. C* **44** (2005) 435–446 [[hep-ph/0410248](https://arxiv.org/abs/hep-ph/0410248)].

- [90] **ALEPH, DELPHI, L3, OPAL, SLD, LEP Electroweak Working Group, SLD Electroweak Group, SLD Heavy Flavour Group** Collaboration, “Precision electroweak measurements on the Z resonance,” *Phys. Rept.* **427** (2006) 257–454 [[hep-ex/0509008](#)].
- [91] **ATLAS** Collaboration, “Precision measurement and interpretation of inclusive W^+ , W^- and Z/γ^* production cross sections with the ATLAS detector,” *Eur. Phys. J. C* **77** (2017) 367 [[arXiv:1612.03016](#)].
- [92] **LHCb** Collaboration, “Measurement of $Z \rightarrow \tau^+\tau^-$ production in proton-proton collisions at $\sqrt{s} = 8$ TeV,” *JHEP* **09** (2018) 159 [[arXiv:1806.05008](#)].
- [93] **Particle Data Group** Collaboration, “Review of Particle Physics,” *PTEP* **2022** (2022) 083C01.
- [94] C. Bobeth, *et al.*, “ $B_{s,d} \rightarrow l^+l^-$ in the Standard Model with Reduced Theoretical Uncertainty,” *Phys. Rev. Lett.* **112** (2014) 101801 [[arXiv:1311.0903](#)].
- [95] M. Beneke, C. Bobeth, and R. Szafron, “Enhanced electromagnetic correction to the rare B -meson decay $B_{s,d} \rightarrow \mu^+\mu^-$,” *Phys. Rev. Lett.* **120** (2018) 011801 [[arXiv:1708.09152](#)].
- [96] A. J. Buras and E. Venturini, “The exclusive vision of rare K and B decays and of the quark mixing in the standard model,” *Eur. Phys. J. C* **82** (2022) 615 [[arXiv:2203.11960](#)].
- [97] **CMS, LHCb, ATLAS** Collaboration, “Combination of the ATLAS, CMS and LHCb results on the $B_{(s)}^0 \rightarrow \mu^+\mu^-$ decays.” <https://cds.cern.ch/record/2727216>.
- [98] W. Altmannshofer and P. Stangl, “New physics in rare B decays after Moriond 2021,” *Eur. Phys. J. C* **81** (2021) 952 [[arXiv:2103.13370](#)].
- [99] **CMS** Collaboration, “Measurement of $B_s^0 \rightarrow \mu^+\mu^-$ decay properties and search for the $B^0 \rightarrow \mu\mu$ decay in proton-proton collisions at $\sqrt{s} = 13$ TeV.” <https://cds.cern.ch/record/2815334>.
- [100] **HFLAV** Collaboration. “Average of $\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$ for End of January 2023” at https://rare-decays-jan23-raredecays-hflav.app.cern.ch/detail/BR_Bs0_mu+_mu-/.
- [101] **DELPHI** Collaboration, “Searches for neutral higgs bosons in extended models,” *Eur. Phys. J. C* **38** (2004) 1–28 [[hep-ex/0410017](#)].

- [102] **ALEPH, DELPHI, L3, OPAL, LEP Working Group for Higgs Boson Searches** Collaboration, “Search for neutral MSSM Higgs bosons at LEP,” *Eur. Phys. J. C* **47** (2006) 547–587 [[hep-ex/0602042](#)].
- [103] **ALEPH, DELPHI, L3, OPAL, LEP** Collaboration, “Search for Charged Higgs bosons: Combined Results Using LEP Data,” *Eur. Phys. J. C* **73** (2013) 2463 [[arXiv:1301.6065](#)].
- [104] **CMS** Collaboration, “Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV,” *JINST* **13** (2018) P10005 [[arXiv:1809.02816](#)].
- [105] **ATLAS** Collaboration, “Identification of hadronic tau lepton decays using neural networks in the ATLAS experiment.” <http://cds.cern.ch/record/2688062>.
- [106] **ATLAS** Collaboration, “Search for direct stau production in events with two hadronic τ -leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector,” *Phys. Rev. D* **101** (2020) 032009 [[arXiv:1911.06660](#)].
- [107] **CMS** Collaboration, “Search for direct pair production of supersymmetric partners of τ leptons in the final state with two hadronically decaying τ leptons and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV.” [arXiv:2207.02254](#).
- [108] S. Iguro, S. Okawa, and Y. Omura, “Light lepton portal dark matter meets the LHC,” *JHEP* **03** (2023) 010 [[arXiv:2208.05487](#)].
- [109] M. Aiko, S. Kanemura, and K. Mawatari, “Exploring the global symmetry structure of the Higgs potential via same-sign pair production of charged Higgs bosons,” *Phys. Lett. B* **797** (2019) 134854 [[arXiv:1906.09101](#)].
- [110] **CMS** Collaboration, “Search for electroweak production of charginos in final states with two τ leptons in pp collisions at $\sqrt{s} = 8$ TeV,” *JHEP* **04** (2017) 018 [[arXiv:1610.04870](#)].
- [111] **ATLAS** Collaboration, “Search for the direct production of charginos and neutralinos in final states with tau leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector,” *Eur. Phys. J. C* **78** (2018) 154 [[arXiv:1708.07875](#)].
- [112] **ATLAS** Collaboration, “Search for chargino–neutralino pair production in final states with three leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector,” *Eur. Phys. J. C* **81** (2021) 1118 [[arXiv:2106.01676](#)].

- [113] **CMS** Collaboration, “Search for supersymmetry in final states with two or three soft leptons and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **04** (2022) 091 [[arXiv:2111.06296](#)].
- [114] **ATLAS** Collaboration, “Search for the direct production of charginos and neutralinos in final states with tau leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector.”. <http://cds.cern.ch/record/2815678>.
- [115] C. G. Lester and D. J. Summers, “Measuring masses of semiinvisibly decaying particles pair produced at hadron colliders,” *Phys. Lett. B* **463** (1999) 99–103 [[hep-ph/9906349](#)].
- [116] A. Barr, C. Lester, and P. Stephens, “m(T2): The Truth behind the glamour,” *J. Phys. G* **29** (2003) 2343–2363 [[hep-ph/0304226](#)].
- [117] J. Alwall, *et al.*, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)].
- [118] P. Skands, S. Carrazza, and J. Rojo, “Tuning PYTHIA 8.1: the Monash 2013 Tune,” *Eur. Phys. J. C* **74** (2014) 3024 [[arXiv:1404.5630](#)].
- [119] **DELPHES 3** Collaboration, “DELPHES 3, A modular framework for fast simulation of a generic collider experiment,” *JHEP* **02** (2014) 057 [[arXiv:1307.6346](#)].
- [120] J. M. Gerard and M. Herquet, “A Twisted custodial symmetry in the two-Higgs-doublet model,” *Phys. Rev. Lett.* **98** (2007) 251802 [[hep-ph/0703051](#)].
- [121] **CMS** Collaboration, “Search for evidence of Type-III seesaw mechanism in multilepton final states in pp collisions at $\sqrt{s} = 13$ TeV.”. <https://cds.cern.ch/record/2256657>.
- [122] **CMS** Collaboration, “Search for new physics in multilepton final states in pp collisions at $\sqrt{s} = 13$ TeV.”. <http://cds.cern.ch/record/2668721>.
- [123] **ATLAS** Collaboration, “Search for type-III seesaw heavy leptons in dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Eur. Phys. J. C* **81** (2021) 218 [[arXiv:2008.07949](#)].
- [124] **CMS** Collaboration, “Search for supersymmetric partners of electrons and muons in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *Phys. Lett. B* **790** (2019) 140–166 [[arXiv:1806.05264](#)].

- [125] **ATLAS** Collaboration, “Search for new phenomena in dijet events using 37 fb⁻¹ of pp collision data collected at $\sqrt{s}=13$ TeV with the ATLAS detector,” *Phys. Rev. D* **96** (2017) 052004 [[arXiv:1703.09127](#)].
- [126] M. Blanke and S. Iguro, “Collider probe of heavy additional Higgs bosons solving the muon $g - 2$.” [arXiv:2210.13508](#).
- [127] E. Kou and P. Urquijo, eds., “The Belle II Physics Book,” *PTEP* **2019** (2019) 123C01 [[arXiv:1808.10567](#)]. [Erratum: *PTEP* 2020, 029201 (2020)].
- [128] E. J. Chun and T. Mondal, “Explaining $g - 2$ anomalies in two Higgs doublet model with vector-like leptons,” *JHEP* **11** (2020) 077 [[arXiv:2009.08314](#)].
- [129] **CMS** Collaboration, “Search for physics beyond the standard model in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **03** (2020) 051 [[arXiv:1911.04968](#)].
- [130] **ATLAS** Collaboration, “Search for type-III seesaw heavy leptons in leptonic final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Eur. Phys. J. C* **82** (2022) 988 [[arXiv:2202.02039](#)].
- [131] **CMS** Collaboration, “Inclusive nonresonant multilepton probes of new phenomena at $\sqrt{s}=13$ TeV,” *Phys. Rev. D* **105** (2022) 112007 [[arXiv:2202.08676](#)].
- [132] **HFLAV** Collaboration, “Averages of b -hadron, c -hadron, and τ -lepton properties as of 2021,” *Phys. Rev. D* **107** (2023) 052008 [[arXiv:2206.07501](#)].
- [133] K. Tobe, “Michel parameters for τ decays $\tau \rightarrow l\nu\bar{\nu}$ ($l = e, \mu$) in a general two Higgs doublet model with $\mu - \tau$ flavor violation,” *JHEP* **10** (2016) 114 [[arXiv:1607.04447](#)].
- [134] X.-Q. Li, J. Lu, and A. Pich, “ $B_{s,d}^0 \rightarrow \ell^+\ell^-$ Decays in the Aligned Two-Higgs-Doublet Model,” *JHEP* **06** (2014) 022 [[arXiv:1404.5865](#)].
- [135] B. W. Lee, C. Quigg, and H. B. Thacker, “Weak Interactions at Very High-Energies: The Role of the Higgs Boson Mass,” *Phys. Rev. D* **16** (1977) 1519.
- [136] S. Kanemura, T. Kubota, and E. Takasugi, “Lee-Quigg-Thacker bounds for Higgs boson masses in a two doublet model,” *Phys. Lett. B* **313** (1993) 155–160 [[hep-ph/9303263](#)].
- [137] A. G. Akeroyd, A. Arhrib, and E.-M. Naimi, “Note on tree level unitarity in the general two Higgs doublet model,” *Phys. Lett. B* **490** (2000) 119–124 [[hep-ph/0006035](#)].

- [138] A. Arhrib, “Unitarity constraints on scalar parameters of the standard and two Higgs doublets model,” in *Workshop on Noncommutative Geometry, Superstrings and Particle Physics*. 2000. [hep-ph/0012353](#).
- [139] I. F. Ginzburg and I. P. Ivanov, “Tree level unitarity constraints in the 2HDM with CP violation.” [hep-ph/0312374](#).
- [140] I. F. Ginzburg and I. P. Ivanov, “Tree-level unitarity constraints in the most general 2HDM,” *Phys. Rev. D* **72** (2005) 115010 [[hep-ph/0508020](#)].
- [141] J. Horejsi and M. Kladiva, “Tree-unitarity bounds for THDM Higgs masses revisited,” *Eur. Phys. J. C* **46** (2006) 81–91 [[hep-ph/0510154](#)].
- [142] M. Maniatis, A. von Manteuffel, O. Nachtmann, and F. Nagel, “Stability and symmetry breaking in the general two-Higgs-doublet model,” *Eur. Phys. J. C* **48** (2006) 805–823 [[hep-ph/0605184](#)].
- [143] P. M. Ferreira and D. R. T. Jones, “Bounds on scalar masses in two Higgs doublet models,” *JHEP* **08** (2009) 069 [[arXiv:0903.2856](#)].
- [144] A. Goudelis, B. Herrmann, and O. Stål, “Dark matter in the Inert Doublet Model after the discovery of a Higgs-like boson at the LHC,” *JHEP* **09** (2013) 106 [[arXiv:1303.3010](#)].