$SU(3)_F$ sum rules for CP asymmetry of $D_{(s)}$ decays

Syuhei Iguro,^{1,2,3,4,5,*} Ulrich Nierste,^{1,†} Emil Overduin,^{1,6,‡} and Maurice Schüßler^{1,6,§}

¹Institute for Theoretical Particle Physics (TTP), Karlsruhe Institute of Technology (KIT),

Wolfgang-Gaede-Str. 1, 76131 Karlsruhe, Germany

²Institute for Astroparticle Physics (IAP), Karlsruhe Institute of Technology (KIT),

Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany

³Institute for Advanced Research (IAR), Nagoya University, Nagoya 464–8601, Japan

⁴Kobayashi-Maskawa Institute (KMI) for the Origin of Particles and the Universe,

Nagoya University, Nagoya 464-8602, Japan

⁵KEK Theory Center, IPNS, KEK, Tsukuba 305-0801, Japan

⁶Institute for Theoretical Physics (ITP), Karlsruhe Institute of Technology (KIT),

Wolfgang-Gaede-Str. 1, 76131 Karlsruhe, Germany

Charge-parity (CP) asymmetries in charm decays are extremely suppressed in the Standard Model and may well be dominated by new-physics contributions. The LHCb collaboration reported the results of direct CP asymmetry measurements in $D^0 \rightarrow K^+ K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays with unprecedented accuracy: $a_{\rm CP}(K^+K^-) = (7.7 \pm 5.7) \times 10^{-4}$ and $a_{\rm CP}(\pi^+\pi^-) = (23.2 \pm 6.1) \times 10^{-4}$, with the latter quantity inferred from the precise measurement of $\Delta a_{\rm CP} = a_{\rm CP}(K^+K^-) - a_{\rm CP}(\pi^+\pi^-) = (-15.7 \pm 2.9) \times 10^{-4}$. When interpreted within the Standard Model, these values indicate a breakdown of the approximate U-spin symmetry of QCD. If, however, this symmetry holds and the data stem from new physics, other CP asymmetries should be enhanced as well. We derive CP asymmetry sum rules based on SU(3) flavor symmetry for D meson decays into a pair of pseudoscalar mesons as well as a pair of a pseudoscalar and a vector meson for two generic scenarios, with $\Delta U = 0$ and $|\Delta U| = 1$ interactions, respectively. The correlations implied by the sum rules can be used to check the consistency between different measurements and to discriminate between these scenarios with future data. For instance, we find $a_{\rm CP}(\pi^+K^{*0}) + a_{\rm CP}(K^+\overline{K}^{*0}) = 0$ for $\Delta U = 0$ new physics and the opposite relative sign for the $|\Delta U| = 1$ case. One sum rule, connecting four decay modes, holds in both scenarios. We further extend our sum rules to certain differences of CP asymmetries from which the D production asymmetries drop out.

KEYWORDS: CP asymmetry, charm physics, D meson decay, sum rule, SU(3) flavor symmetry

I. INTRODUCTION

In 2019 the LHCb collaboration reported the discovery of charm CP violation (CPV) in the measurement of the difference of two CP asymmetries stating [1]

$$\Delta a_{\rm CP}^{2019} = a_{\rm CP} (K^+ K^-) - a_{\rm CP} (\pi^+ \pi^-)$$

= (-15.7 ± 2.9) × 10⁻⁴, (1)

where $a_{\rm CP}(f)$ is the time-integrated direct CP asymmetry in $D^0 \to f$. Strictly speaking, the measurement in Eq. (1) contains a small contribution from (the yet undiscovered) mixing-induced CP violation, because the average decay times of the $D^0 \to K^+K^-$ and $D^0 \to \pi^+\pi^$ data samples are different. In this paper we assign the measured values completely to direct CP asymmetry; subtracting the maximal experimentally allowed contribution from mixing-induced CP violation changes the central value of the direct CP asymmetry difference in Eq. (1) by as little as $+0.3 \times 10^{-4}$ [1].

In 2022 LHCb presented the corresponding measurement of the individual CP asymmetry $a_{\rm CP}(K^+K^-)$ and combined it with Eq. (1) and previous measurements to find [2]:

$$a_{\rm CP}(K^+K^-) = (7.7 \pm 5.7) \times 10^{-4},$$
 (2)

$$a_{\rm CP}(\pi^+\pi^-) = (23.2 \pm 6.1) \times 10^{-4},$$
 (3)

with a correlation of $\rho = 0.88$.

It is difficult to calculate Standard-Model (SM) predictions for the penguin amplitudes feeding Eqs. (2) and (3). However, the strong parametric suppression stemming from tiny off-diagonal elements of the Cabbibo-Kobayashi-Maskawa (CKM) matrix [3, 4] makes these CP asymmetries highly sensitive probes of new physics. Virtual effects of multi-TeV mass heavy particles can easily dominate over the SM contribution, so that even SM predictions with $\mathcal{O}(100\%)$ uncertainty can constrain beyond-SM (BSM) models in a meaningful way. A SM prediction based on QCD sum rules is [5, 6]

$$|\Delta a_{\rm CP}^{\rm SM}| = (2.4 \pm 1.2) \times 10^{-4}, \tag{4}$$

which is smaller than the measured value in Eq. (1) by a factor of more than six. While QCD sum rules are a

^{*} igurosyuhei@gmail.com

[†] ulrich.nierste@kit.edu

[‡] overduin.emil@gmail.com

[§] maurice.schuessler@gmail.com

sound, field-theoretic method with a plethora of successful predictions in *B* physics, little is known about their applicability to charm physics. The discrepancy between Eqs. (1) and (4) (as well as already the earlier, less significant measurement $\Delta a_{\rm CP} = (-82 \pm 21 \pm 11) \times 10^{-4}$ [7]) has stimulated many theory papers addressing either BSM physics [6, 8–21] or invoking a SM explanation in terms of an enhanced SM penguin amplitude. The latter papers have postulated a QCD enhancement ad-hoc [22, 23] or by invoking unflavored resonances which are almost mass degenerate with the D^0 [24, 25].

The approximate SU(3) flavor symmetry of QCD $(SU(3)_F)$ can be used to derive relations between various CP asymmetries and we expect to predict other non-vanishing direct CP asymmetries from the non-zero CP asymmetry in Eq. (3). To this end we employ the subgroup of SU(3)_F corresponding to SU(2) rotations of the U-spin doublet $(s, d)^T$. The U-spin symmetry breaking parameter is $(m_s - m_d)/\Lambda_{\rm QCD}$, so that one expects U-spin relations to hold up to corrections of order 30%. SU(3)_F analyses of branching ratios (BR) and CP asymmetries can be found in Refs. [26–36]. However, within the SM one finds the sum rule

$$a_{\rm CP}(\pi^+\pi^-) = -a_{\rm CP}(K^+K^-), \tag{5}$$

valid in the limit of exact U-spin symmetry. As seen the sum rule predicts opposite signs of $a_{\rm CP}(\pi^+\pi^-)$ and $a_{\rm CP}(K^+K^-)$, and deviates from the measurement by about 2.7 σ [2].

Thus the experimental results in Eqs. (2) and (3) imply that

- (i) U-spin breaks down in charm CP asymmetries
- (ii) or the dominant contribution to at least one of the two CP asymmetries in Eqs. (2) and (3) stems from new physics (NP) [21, 37]
- (iii) or future measurements will find different values for $a_{\rm CP}(K^+K^-)$ and/or $\Delta a_{\rm CP}$; this possibility necessarily implies a shift of $a_{\rm CP}(K^+K^-)$ by more than 2σ with a change of the sign. (We do not consider the possibility that $\Delta a_{\rm CP}$ will change by far more than 5σ to comply with $a_{\rm CP}(K^+K^-) > 0$).

Although the short-distance partonic level quark transitions can be evaluated perturbatively, the hadronic D meson decays and its CPV parameters involve hadronic matrix elements such as $\langle K^+K^-|(\bar{u}\gamma_\mu P_L s)(\bar{s}\gamma^\mu P_L c)|D^0\rangle$, which are not easily evaluated.

In this paper, we rely on the approximate $SU(3)_F$ symmetry of the QCD Lagrangian to correlate the amplitudes of different decay modes with the goal of discriminating between the three explanations listed above. For example, if (ii) is the correct explanation while U-spin holds, the pattern of Eqs. (2) and (3) will have imprints on other decay modes. Furthermore, the comparisons of CP asymmetries in $D_{(s)}$ decays to two pseudoscalar mesons with



FIG. 1. The experimental result and theory predictions on $a_{\rm CP}(K^+K^-)$ vs $a_{\rm CP}(\pi^+\pi^-)$ plane. See the text at the end of Sec. II for details.

those in decays to a pseudoscalar/vector meson pair will give insight into the Dirac structure of the underlying BSM couplings. We will derive $SU(3)_F$ sum rules for both classes of decays. Previously sum rules for amplitudes and decay rates of charmed meson decays were derived in Ref. [30].^{#1}

The outline of this paper is as follows. In Sec. II, we explain the setup and in Sec. III the CP asymmetry sum rules are derived. In Sec. IV, we extend our sum rules to differences of CP asymmetries modeled after Eq. (1) in order to eliminate experimental production asymmetries. We conclude in Sec. V.

II. FRAMEWORK

Within the SM the decays of interest are induced by the singly Cabibbo suppressed (SCS) charm decays $c \rightarrow uq\bar{q}$ with q = d, s at tree-level and q = u, d, s in the loop-induced penguin contribution. The relevant $|\Delta C| =$ 1 effective Hamiltonian is given at the interaction scale

^{#1} See also Ref. [38] and references therein. The earlier work of Ref. [32] discusses CP violation in charmless B decays based on SU(3)_F.

 $(\mu = m_c)$ as

$$\mathcal{H}_{\rm SM}^{\rm eff} = \frac{4G_F}{\sqrt{2}} \sum_{q=s,d} \lambda_q \left(C_1(\bar{q}^{\alpha} \gamma^{\mu} P_L c^{\alpha}) (\bar{u}^{\beta} \gamma_{\mu} P_L q^{\beta}) + C_2(\bar{q}^{\alpha} \gamma^{\mu} P_L c^{\beta}) (\bar{u}^{\beta} \gamma_{\mu} P_L q^{\alpha}) \right)$$
$$\equiv \lambda_s h_{\rm SM}^s + \lambda_d h_{\rm SM}^d, \tag{6}$$

where $\lambda_q = V_{uq}V_{cq}^*$ and α, β are color indices. Defining $A_{\pi} = \langle \pi^+\pi^- | h_{\rm SM}^d - h_{\rm SM}^s | D^0 \rangle$, $A_K = \langle K^+K^- | h_{\rm SM}^s - h_{\rm SM}^d | D^0 \rangle$, $P_{\pi} = \langle \pi^+\pi^- | h_{\rm SM}^s | D^0 \rangle$ and $P_K = \langle K^+K^- | h_{\rm SM}^d | D^0 \rangle$, the decay amplitudes are expressed as

$$\mathcal{A}(D^0 \to \pi^+ \pi^-) = \langle \pi^+ \pi^- | \mathcal{H}_{\rm SM}^{\rm eff} | D^0 \rangle$$
$$= \lambda_d A_\pi - \lambda_b P_\pi, \tag{7}$$

$$\mathcal{A}(D^0 \to K^+ K^-) = \lambda_s A_K - \lambda_b P_K, \tag{8}$$

where the CKM unitarity relation $\lambda_d + \lambda_s + \lambda_b = 0$ is used. The *b* quark is integrated out leading to penguin operators with tiny coefficients in $\mathcal{H}_{\text{SM}}^{\text{eff}}$ which come with λ_b and are omitted in Eq. (6). These small terms contribute equally to h_{SM}^s and h_{SM}^d .

From Eqs. (7) and (8) one notes that the meson pair is produced in a U = 1 state in the limit $\lambda_b = 0$. It is straightforward to calculate the direct CP asymmetry

$$a_{\rm CP} \equiv \frac{|\mathcal{A}_{i \to f}|^2 - |\mathcal{A}_{\overline{i} \to \overline{f}}|^2}{|\mathcal{A}_{i \to f}|^2 + |\mathcal{A}_{\overline{i} \to \overline{f}}|^2},\tag{9}$$

and we obtain

$$a_{\rm CP}(\pi^+\pi^-) \simeq 2\,{\rm Im}\,\frac{\lambda_b}{\lambda_d}{\rm Im}\,\left(\frac{P_\pi}{A_\pi}\right),\qquad(10)$$

$$a_{\rm CP}(K^+K^-) \simeq 2 \,{\rm Im} \,\frac{\lambda_b}{\lambda_s} {\rm Im} \,\left(\frac{P_K}{A_K}\right),$$
 (11)

where we neglected $\mathcal{O}(\lambda_b^2)$ terms. Given that $\lambda_s = -\lambda_d + \mathcal{O}(\lambda_b)$ holds thanks to CKM unitarity, *U*-spin symmetry ensures $A_{\pi} = A_K$ and $P_{\pi} = P_K$, reproducing the famous CP asymmetry sum rule of Eq. (5).

In reality SU(3)_F is broken and the breaking effect is found to be $\simeq 30\%$ in the dominant decay amplitude $\propto \lambda_{d,s}$ in measurements of BRs [30]. In this paper, we employ SU(3)_F at the leading order and neglect *U*spin symmetry violation since the observed violation in Eqs. (2) and (3) is huge. In the SM the penguin contributions P_{π} and P_K are $\Delta U = 0$ amplitudes. $\Delta U = 0$ NP contributes to $a_{\rm CP}(\pi^+\pi^-)$ and $a_{\rm CP}(K^+K^-)$ with opposite sign as the SM one, so that a NP explanation of Eqs. (2) and (3) requires a $|\Delta U| = 1$ contribution. If such a contribution is observed in future measurements of other CP asymmetries, this will corroborate the NP interpretation. To this end, we derive CP sum rules for decays in which non-zero CP asymmetries are not yet observed. Generic NP four-quark $\Delta S = 0$ interactions can be described by amending the effective SM Hamiltonian in Eq. (6) with

$$\Delta \mathcal{H}_{\rm NP}^{\rm eff} = \frac{G_F}{\sqrt{2}} (\overline{u} \Gamma c) \left(a_u \, \overline{u} \Gamma u + a_d \, \overline{d} \Gamma d + a_s \, \overline{s} \Gamma s \right)$$
$$\equiv a_u \mathcal{O}'_u + a_d \mathcal{O}'_d + a_s \mathcal{O}'_s \,, \tag{12}$$

where Γ represents an arbitrary Dirac structure. While several such terms with different Dirac structures could be present, our symmetry-based analyses will not be changed compared to the case in Eq. (12) with a single Dirac structure. The same remark applies to the two possible color structures; color indices are not shown in Eq. (12).

Returning to $D^0 \to \pi^+\pi^-$ and K^+K^- , we set $a_u = 0$ until the end of this section, because a_u contributes only through penguin or annihilation diagrams to these decays which are likely to be smaller than tree-level NP effects involving a_d or a_s . The contributing amplitudes in the presence of NP effects, $\mathcal{H}_{\rm NP}^{\rm eff} = \mathcal{H}_{\rm SM}^{\rm eff} + \Delta \mathcal{H}_{\rm NP}^{\rm eff}$ are expressed as,

$$A^{\rm NP}(D^0 \to \pi^+\pi^-) = \lambda_d A_\pi + a_d \ \mathcal{Q}_d^\pi + a_s \mathcal{Q}_s^\pi, \qquad (13)$$

$$A^{\rm NP}(D^0 \to K^+ K^-) = \lambda_s A_K + a_d \ \mathcal{Q}_d^K + a_s \mathcal{Q}_s^K, \quad (14)$$

where $Q_q^M = \langle M\overline{M}|\mathcal{O}'_q|D^0\rangle$ is defined. For instance, if we introduce NP which only couples to d quarks, this corresponds to $a_d \neq 0$ and $a_s = 0$. We emphasize that as long as the involved hadronic matrix elements cannot be determined accurately, the $\Delta U = 0$ NP contribution $\propto a_d + a_s$ cannot be disentangled from the SM contribution. Similarly to the SM case we obtain

$$a_{\rm CP}(\pi^+\pi^-) \simeq -2\,{\rm Im}\,\frac{a_d}{\lambda_d}\,{\rm Im}\,\frac{\mathcal{Q}_d^\pi}{A} - 2\,{\rm Im}\,\frac{a_s}{\lambda_d}\,{\rm Im}\,\frac{\mathcal{Q}_s^\pi}{A},\qquad(15)$$

$$a_{\rm CP}(K^+K^-) \simeq -2 \operatorname{Im} \frac{a_d}{\lambda_s} \operatorname{Im} \frac{\mathcal{Q}_d^K}{A} - 2 \operatorname{Im} \frac{a_s}{\lambda_s} \operatorname{Im} \frac{\mathcal{Q}_s^K}{A},$$
 (16)

working to leading order in λ_b and $a_{d,s}$ and using the *U*-spin symmetry which holds approximately, $A \equiv A_{\pi} = A_K$, in the SM part. The maximal *U*-spin breaking in $\Delta a_{\rm CP}$ corresponds to ${\rm Im} (a_d + a_s) = 0$. In this scenario $a_{\rm CP}(\pi^+\pi^-) = a_{\rm CP}(K^+K^-)$ holds, however, this relation also does not fit the recent data.

In Fig. 1 we show the experimental status and theory predictions in the $a_{\rm CP}(K^+K^-)$ vs $a_{\rm CP}(\pi^+\pi^-)$ plane. The 2019 LHCb result for $\Delta a_{\rm CP}$ is shown in orange with 1σ uncertainty. We show the latest LHCb result of 1, 2, and 3σ in blue solid, dashed, and dotted ellipses. The U-spin limit, $a_{\rm CP}(K^+K^-) = -a_{\rm CP}(\pi^+\pi^-)$ as well as the limit of maximal U-spin violation, $a_{\rm CP}(K^+K^-) =$ $+a_{\rm CP}(\pi^+\pi^-)$ are shown in light green and red, respectively. To account for U-spin violating effects based on the ratio of BR $(D^0 \rightarrow K^+K^-)$ and BR $(D^0 \rightarrow$ $\pi^+\pi^-)$ [30], we allow 30% deviation from the U-spin limit corresponding to the green band. The magenta lines correspond to the reference points, $a_{\rm CP}(\pi^+\pi^-) =$ $\pm 3 a_{\rm CP}(K^+K^-)$. These reference points are motivated by the case $a_d \neq 0$ with $a_s = 0$ and the observation that \mathcal{Q}_d^K is color-suppressed w.r.t. \mathcal{Q}_d^{π} . The phase difference between \mathcal{Q}_d^K/A and \mathcal{Q}_d^{π}/A can be anything and the two signs in $a_{\rm CP}(\pi^+\pi^-) = \pm 3 a_{\rm CP}(K^+K^-)$ are the limiting cases if the color suppression is at its nominal value of $1/N_c = 1/3$.

Assuming Im $\left(\frac{Q_d^{-}}{A}\right) = 1$, the distance between any two points next to each other on the lines corresponds to $\Delta \text{Im}(a_d) = 0.05 \times 10^{-3}$. It is evident that $|\Delta U| =$ 1 *i.e.* maximal U-spin violation cannot explain the data either while the central value of the recent LHCb data can be reproduced with $a_{\text{CP}}(\pi^+\pi^-) = +3 a_{\text{CP}}(K^+K^-)$.

III. CP ASYMMETRY SUM RULES

In this section, we derive sum rules connecting new CP asymmetries, valid for $\Delta U = 0$ and $|\Delta U| = 1$ interactions, respectively. The derivation is similar to that of the amplitude sum rule performed in Ref. [30], however, it is hard to find CP asymmetry sum rules in general, since CP asymmetries involve interference between two amplitudes, and thus the number of independent relations is smaller. For demonstration, we start with a generic decay amplitude of

$$\langle M_1 M_2 | \mathcal{H} | D \rangle = \lambda A^{(M_1 M_2)} + a P^{(M_1 M_2)}.$$
 (17)

This leads to a CP asymmetry of

$$a_{\rm CP}(M_1M_2) \simeq$$

 $+2 \frac{{\rm Im}(a)}{\lambda} \frac{{\rm Im}\left(A^{(M_1M_2)} P^{(M_1M_2)*}\right)}{|A^{(M_1M_2)}|^2}, \quad (18)$

where terms of order a^2 and higher are neglected, and the relative complex phase is put in a while λ is chosen real. It is helpful to decompose the amplitude via the Wigner-Eckart theorem, which allows us to rewrite the amplitude in terms of Clebsch-Gordan (CG) coefficients [39, 40] and reduced matrix elements, to construct the desired CP asymmetry sum rules. The relevant coefficients are summarized in Tab. I and Tabs. II, III for $D \to PP$ and $D \to PV$, where P and V stands for a pseudoscalar meson and vector meson, respectively.

Schematically, we decompose the amplitudes $A^{(M_1M_2)}$ and $P^{(M_1M_2)}$ as,

$$A^{(M_1M_2)} = \sum_{i=1}^{n} c_i (M_1M_2) x_i, \qquad (19)$$

$$P^{(M_1M_2)*} = \sum_{j=1}^{m} c'_j(M_1M_2)y_j,$$
(20)

where x_i , y_j are in general independent reduced matrix elements and $c_i(M_1M_2)$, $c'_j(M_1M_2)$ correspond to the CG coefficients where the relevant entries are summarized in Appendix A. The product is expressed as

$$\delta_{\rm CP}(M_1 M_2) \equiv A^{(M_1 M_2)} P^{(M_1 M_2)*}$$

= $\sum_{i=1}^n \sum_{j=1}^m c_i (M_1 M_2) c'_j (M_1 M_2) x_i y_j$
= $(c_1 c'_1, c_1 c'_2, ..., c_n c'_m) \cdot \begin{pmatrix} x_1 y_1 \\ \vdots \\ x_n y_m \end{pmatrix}$
= $\mathbf{c} (M_1 M_2)^T \cdot \mathbf{x}$. (21)

In this convention, CP asymmetries are expressed as

$$a_{\rm CP}(M_1M_2) = \chi(M_1M_2) \operatorname{Im} \left(\delta_{\rm CP}(M_1M_2)\right) \operatorname{Im} (a), \quad (22)$$

where $\chi(M_1M_2) = +2/(\lambda |A^{(M_1M_2)}|^2)$ is defined. We can express $|A^{(M_1M_2)}|^2$ in terms of the decay rate Γ , which is experimentally found from the measured BR:

$$\lambda^{2} |A^{(M_{1}M_{2})}|^{2} = \Gamma(D \to M_{1}M_{2})$$

= BR $(D \to M_{1}M_{2})/\tau_{D}$, (23)

where the different lifetimes τ_D for $D = D^0, D^+, D_s^+$ must be taken into account. Eq. (23) holds, because $|P^{(M_1M_2)}|$ is too small to have an effect on BR $(D \rightarrow M_1M_2)$. The phase space factor is absorbed into the definition of $A^{(M_1M_2)}$; the phase space factors of different two-body D decays are equal in the SU(3)_F symmetry limit and indeed do not differ much from each other. We have

$$\chi(M_1 M_2) = \frac{2 \lambda \tau_D}{\text{BR}(D \to M_1 M_2)},$$
(24)

and hence

$$\operatorname{Im}\left(\delta_{\mathrm{CP}}(M_1 M_2)\right) \lambda \operatorname{Im}(a) = a_{\mathrm{CP}}(M_1 M_2) \frac{\operatorname{BR}(D \to M_1 M_2)}{2 \tau_D}.$$
 (25)

We will quote sum rules in terms of the $\delta_{CP}(M_1M_2)$'s; to relate these to the measured CP asymmetries, BRs, and lifetimes one must use Eq. (25). Since the sum rules are linear in the $\delta_{CP}(M_1M_2)$'s, the overall normalisation does not matter. For this reason, also the theoretical parameter $\lambda \text{ Im } (a)$ drops out from the sum rules and cannot be determined.

Next, we consider a vector of all CP asymmetries

$$\mathbf{a}_{CP}^{T} = (a_{CP,1}, a_{CP,2}...a_{CP,n}).$$
 (26)

Then constructing sum rules is equivalent to finding a vector \vec{v} orthogonal to \mathbf{a}_{CP} , which satisfies

$$\mathbf{v}^T \cdot \mathbf{a}_{\rm CP} = 0. \tag{27}$$

If a sum rule involves only two modes, we can directly construct the $a_{\rm CP}$ sum rule from the $\delta_{\rm CP}$ sum rule, see, Appendix B for detail.

The general procedure discussed above can also be performed incorporating higher orders of $SU(3)_F$ breaking. The cost is a larger number of involved reduced matrix elements. We note that since the matrix **c** grows with the number of reduced matrix elements squared, it will be more difficult to find sum rules for CP asymmetries incorporating $SU(3)_F$ breaking effect.

To incorporate the generic interaction we consider the following general amplitude of the pseudoscalar decays,

$$\langle M_1 M_2 | \mathcal{H} | D \rangle = \lambda_{SM} A^{(M_1 M_2)} + a_0 P_0^{(M_1 M_2)} + a_1 P_1^{(M_1 M_2)}, \quad (28)$$

where $P_0^{(M_1M_2)}$ and $P_1^{(M_1M_2)}$ correspond to the $\Delta U = 0$ and $|\Delta U| = 1$ contributions, respectively. The SM $|\Delta U| = 1$ amplitude with the CKM factor $\lambda_{SM} = \lambda_{d,s}$ is $A^{(M_1M_2)}$. Both SM penguin and NP $\Delta U = 0$ contribution are contained in $P_0^{(M_1M_2)}$, while $P_1^{(M_1M_2)}$ stems solely from NP. Keeping terms up to linear order in a_0 and a_1 , the contributions to the CP asymmetry can be separated into two parts as

$$a_{\rm CP} = a_{\rm CP}^{\Delta U=0} + a_{\rm CP}^{\Delta U=1}$$

\$\approx 2 \Im \frac{a_0}{\lambda_{SM}} \frac{\Im(A \P_0^*)}{|A|^2} + 2 \Im \frac{a_1}{\lambda_{SM}} \frac{\Im(A \P_1^*)}{|A|^2}. (29)\$

It is difficult to find vectors **v** satisfying Eq. (27) if both a_0 and a_1 are non-zero, because the CG coefficients c'_j in Eq. (20) are different for $\Delta U = 0$ and $|\Delta U| = 1$ matrix elements in general.

In the following two sub-sections (Secs. III A and III B) we present CP asymmetry sum rules for $D \rightarrow PP$ and $D \rightarrow PV$, respectively, considering the cases $a_1 = 0$ and $a_0 = 0$. Specifically, we consider two scenarios, characterised by U-spin U and isospin I:

- I. $\Delta U = \Delta I = 0$: We assume $\Delta \mathcal{H}_{\text{NP}}^{\text{eff}}$ in Eq. (12) to be an SU(3) singlet, $a_0 \propto a_u = a_d = a_s$ and $a_1 = 0$. This NP scenario mimics the SM penguin contribution, but with a_0 unrelated to λ_b .
- II. $|\Delta U| = 1$ with $a_0 = a_u = 0$ and $a_1 \propto a_s = -a_d$. This NP scenario is motivated by a heavy new charged particle, such as a charged Higgs boson, though such a particle will also involve $\Delta U = 0$ interactions (and effects on DCS decays) as well. Also, a neutral particle with FCNC $\bar{u}c$ coupling could produce this situation, if the coupling to up quarks is suppressed.

A. $D \rightarrow PP$

1. $\Delta U = 0$ New Physics

First, we present CP asymmetry sum rules which hold for $\Delta U = 0$ interactions $(a_0 \neq 0, a_1 = 0)$. The following two sum rules are well known and are found by a naive interchange of d and s quarks.

$$a_{\rm CP}^{\Delta U=0}(K^-K^+) + a_{\rm CP}^{\Delta U=0}(\pi^-\pi^+) = 0,$$
 (30)

$$a_{\rm CP}^{\Delta U=0}(K^0\pi^+) + a_{\rm CP}^{\Delta U=0}(\overline{K}^0K^+) = 0.$$
 (31)

To clarify our notation, remember that SCS decays do not change the strangeness S, so that in this section all $a_{\rm CP}(M_1^0M_2^+)$'s in which the final state $M_1^0M_2^+$ has S = 1 (like those in Eq. (31)) stem from $D_s^+ \to M_1M_2^+$.

The first sum rule Eq. (30) is the same as Eq. (5) and found to be violated by the latest measurements. The experimental data leads to

$$a_{\rm CP}(K^-K^+) + a_{\rm CP}(\pi^-\pi^+) = (30.9 \pm 11.4) \times 10^{-4},$$
(32)

which deviates from the $\Delta U = 0$ sum rule by more than 2σ , as seen in Fig. 1. This is already an interesting hint that there may be more contributions beyond the $\Delta U = 0$ penguin interaction. The last four sum rules do not contain two modes,

$$a_{\rm CP}^{\Delta U=0}(\pi^0 \pi^+) = 0, \qquad (33)$$

$$\delta_{\rm CP}^{\Delta U=0}(\eta_8 \eta_8) + \delta_{\rm CP}^{\Delta U=0}(\pi^0 \pi^0) + 2 \, \delta_{\rm CP}^{\Delta U=0}(\eta_8 \pi^0) = 0, \qquad (34)$$

$$\delta_{\rm CP}^{\Delta U=0}(\eta_8 K^+) + \delta_{\rm CP}^{\Delta U=0}(\eta_8 \pi^+) + \delta_{\rm CP}^{\Delta U=0}(\pi^0 K^+) = 0,$$
(35)

$$3\,\delta_{\rm CP}^{\Delta U=0}(\eta_8 K^+) - 3\,\delta_{\rm CP}^{\Delta U=0}(\pi^0 K^+) + \delta_{\rm CP}^{\Delta U=0}(K^0 \pi^+) = 0.$$
(36)

Eq. (33) is, of course, a well-known null test of the SM, which is not violated if the NP contribution is pure $\Delta I = 0$ as in the considered SU(3) singlet NP scenario. η_8 is the octet η meson. The physical η meson is dominantly η_8 plus a smaller admixture of the singlet state η_0 . The associated mixing angle vanishes in the limit of the exact SU(3)_F symmetry; since we neglect SU(3)_F breaking, the sum rules quoted in this paper can be used with the replacement $\eta_8 \to \eta$.

We emphasize that any linear combination of the above sum rules holds as well. With Eq. (25) one finds the sum rules for the CP asymmetries from the ones quoted for the $\delta_{\rm CP}$'s, with the overall factor $\lambda_{SM} {\rm Im}(a_0)$ dropping out. If in future measurements these sum rules are violated significantly beyond the nominal ~ 30% U-spin breaking, this will be evidence of $|\Delta U| = 1$ NP.

2. $|\Delta U| = 1$ New Physics

Next, we consider CP asymmetry sum rules for $a_0 = 0$ and $a_1 \neq 0$. There are two $|\Delta U| = 1$ contributions, one from the SM amplitude $\lambda_{SM} A^{(M_1M_2)}$ and one from the NP amplitude $a_1 P_1^{(M_1M_2)}$ carrying a different CP phase. Here we can again use the described procedure in Appendix B to find the two-mode sum rules. This time there are four sum rules containing only two modes,

$$a_{\rm CP}^{\Delta U=1}(K^-K^+) - a_{\rm CP}^{\Delta U=1}(\pi^-\pi^+) = 0, \qquad (37)$$
$$a_{\rm CP}^{\Delta U=1}(K^0\pi^+) - a_{\rm CP}^{\Delta U=1}(\overline{K}^0K^+) = 0, \qquad (38)$$

$$a_{CP}^{2D} = (K^{*}\pi^{+}) - a_{CP}^{2D} = (K^{*}K^{+}) = 0, \quad (38)$$
$$a_{CP}^{2D} = (m_{s}m_{s}) - a_{CP}^{2D} = (m_{s}m_{s}^{0}) = 0. \quad (39)$$

$$a_{\rm CP}^{\Delta U=1}(\eta_8\eta_8) - a_{\rm CP}^{\Delta U=1}(\eta_8\pi^0) = 0.$$
(40)

The last two sum rules contain four modes and are given by

$$\begin{split} &\delta_{\rm CP}^{\Delta U=1}(\eta_8 K^+) - \delta_{\rm CP}^{\Delta U=1}(\pi^0 \pi^+) - \delta_{\rm CP}^{\Delta U=1}(\eta_8 \pi^+) \\ &+ \delta_{\rm CP}^{\Delta U=1}(\pi^0 K^+) = 0, \end{split} \tag{41} \\ &6 \, \delta_{\rm CP}^{\Delta U=1}(\pi^0 \pi^+) - 3 \, \delta_{\rm CP}^{\Delta U=1}(\eta_8 K^+) + 3 \, \delta_{\rm CP}^{\Delta U=1}(\pi^0 K^+) \\ &- \delta_{\rm CP}^{\Delta U=1}(K^0 \pi^+) = 0. \end{split}$$

The sum rule in Eq. (37) cannot explain the LHCb measurement of,

$$\Delta a_{\rm CP} = a_{\rm CP} (K^- K^+) - a_{\rm CP} (\pi^- \pi^+)$$

= (-15.7 ± 2.9) × 10⁻⁴, (43)

because $|\Delta U| = 1$ contributions drop out from $\Delta a_{\rm CP}$, their sole effect is to shift $a_{\rm CP}(K^-K^+)$ and $a_{\rm CP}(\pi^-\pi^+)$ into the region compatible with the U-spin symmetric matrix elements.

Interestingly, the sum rule in Eq. (42) holds in both of our two scenarios; for the $\Delta U = 0$ case it is constructed as $6 \times \text{Eq.}(33) - \text{Eq.}(36)$. While current experimental data do not allow us to verify this sum rule due to large experimental uncertainties, it will give useful insight into the quality of SU(3) symmetry of the hadronic matrix elements in the future: In case that data will comply well with Eq. (42), one will gain confidence in the SU(3) method and use the other sum rules to discriminate between the scenarios. Note that a future establishment of $a_{\text{CP}}(\pi^0\pi^+) \neq 0$ will establish isospin-breaking NP and thereby falsify the SM and our scenario I; in our scenario II then at least one other CP asymmetry entering Eq. (42) will be sizable because of the factor 6 in front of $\delta_{\text{CP}}^{\Delta U=1}(\pi^0\pi^+)$.

Above we separately derived the CP asymmetry sum rules for $|\Delta U| = 1$ and $\Delta U = 0$ interactions assuming specific $a_{d,s,u}$ combinations for each. As it is seen in Fig. 1, neither single $|\Delta U| = 1$ nor $\Delta U = 0$ interactions can fully address the current data. Current data prefer the ratio $a_s: a_d = 1: -3$ which leads to

$$(a_s + a_d) : (a_s - a_d) = -1 : 2.$$
(44)

Generally, one can combine the CP-asymmetry sum rules in this proportion and confront these weighted sum rules with data. However, one should keep in mind that a_u enters our scenarios differently: Modifying scenario I by choosing $a_u = 0$ to comply with scenario II will still be a $\Delta U = 0$ scenario, but now with isospin breaking, invalidating Eq. (33). Then two possibilities must be considered: If $a_{\rm CP}(\pi^0\pi^+) \neq 0$ is measured, this will directly establish NP with $a_u \neq a_d$. Yet if $a_{\rm CP}(\pi^0\pi^+)$ is measured compatible with zero, this means that either Im $a_u \approx \text{Im } a_d$ or that the strong phase between the SM tree amplitude and the NP amplitude $\propto a_u - a_d$ is small, see Eq. (16). In the latter case the effect of $a_u - a_d \neq 0$ drops out from the $\Delta U = 0$ sum rules and the abovementioned weighted sum rules are meaningful.

B.
$$D \rightarrow PV$$

Next, we consider the CP sum rule for $D \rightarrow PV$ decays valid for the $\Delta U = 0$ interactions of our scenario I. Considering the penguin operators for SU(3), the following sum rules hold,

$$a_{\rm CP}^{\Delta U=0}(K^0 \overline{K}^{*0}) + a_{\rm CP}^{\Delta U=0}(\overline{K}^0 K^{*0}) = 0, \tag{45}$$

$$a_{\rm CP}^{\Delta C^{-0}}(K^{K^+}) + a_{\rm CP}^{\Delta c^{-0}}(\pi^{-\rho^+}) = 0, \tag{46}$$

$$a_{\rm CP}^{\Delta C^{-0}}(K^+K^{*-}) + a_{\rm CP}^{\Delta C^{-0}}(\pi^+\rho^-) = 0, \tag{47}$$

$$a_{\rm CP}^{\Delta U=0}(K^0 \rho^+) + a_{\rm CP}^{\Delta U=0}(\overline{K}^0 K^{*+}) = 0, \tag{48}$$

$$a_{\rm CP}^{\Delta U=0}(\pi^+ K^{*0}) + a_{\rm CP}^{\Delta U=0}(K^+ \overline{K}^{*0}) = 0, \tag{49}$$

$$\delta_{\rm CP}^{\Delta U=0}(\eta_8 \omega_8) + \delta_{\rm CP}^{\Delta U=0}(\eta_8 \rho^0) + \delta_{\rm CP}^{\Delta U=0}(\pi^0 \omega_8)$$

$$+ \delta_{\rm CP}^{\Delta U=0}(\pi^0 \rho^0) = 0, \tag{50}$$

$$\delta_{\rm CP}^{\Delta U=0}(\pi^0 K^{*+}) + \delta_{\rm CP}^{\Delta U=0}(\eta_8 K^{*+}) + \delta_{\rm CP}^{\Delta U=0}(\pi^0 \rho^+) + \delta_{\rm CP}^{\Delta U=0}(\eta_8 \rho^+) = 0.$$
(51)

$$\delta_{\rm CP}^{\Delta U=0}(K^+\rho^0) + \delta_{\rm CP}^{\Delta U=0}(K^+\omega_8) + \delta_{\rm CP}^{\Delta U=0}(\pi^+\rho^0)$$

$$+ \delta_{\rm CP}^{\Delta U=0} (\pi^+ \omega_8) = 0, \tag{52}$$

$$\epsilon_5 \Delta U=0 (\pi^+ \omega_8) - \epsilon_5 \Delta U=0 (-0, +) - \epsilon_5 \Delta U=0 (\pi^- e^+)$$

$$b\delta_{\rm CP}^{\Delta U} = (\eta_8 \rho^*) - b\delta_{\rm CP}^{\Delta U} = (\pi^* \omega_8) - 5\delta_{\rm CP}^{\Delta U} = (\eta_8 \rho^+) - 3\delta_{\rm CP}^{\Delta U=0}(\pi^0 \rho^+) - \delta_{\rm CP}^{\Delta U=0}(\overline{K}^0 K^{*+}) + \delta_{\rm CP}^{\Delta U=0}(K^+ \overline{K}^{*0}) + 3\delta_{\rm CP}^{\Delta U=0}(\pi^+ \omega_8) + \delta_{\rm CP}^{\Delta U=0}(\pi^+ \rho^0) - 2\delta_{\rm CP}^{\Delta U=0}(\eta_8 K^{*+}) - 2\delta_{\rm CP}^{\Delta U=0}(K^+ \rho^0) = 0.$$
(53)

The first five sum rules only involve two modes, while the remaining three rules involve four decay modes. The last one involves 10 decay modes.

On the other hand, once we assume that NP enters via $|\Delta U| = 1$ operators, the sum rules for CP-asymmetries

can be written as

$$a_{\rm CP}^{\Delta U=1}(\pi^0 \rho^0) - a_{\rm CP}^{\Delta U=1}(\eta_8 \omega_8) = 0, \tag{54}$$

$$a_{\rm CP}^{\Delta U=1}(\overline{K}^0 K^{*0}) - a_{\rm CP}^{\Delta U=1}(K^0 \overline{K}^{*0}) = 0, \tag{55}$$

$$a_{\rm CP}^{\Delta U=1}(\pi^-\rho^+) - a_{\rm CP}^{\Delta U=1}(K^-K^{*+}) = 0, \tag{56}$$

$$a_{\rm CP}^{\Delta U=1}(\pi^+\rho^-) - a_{\rm CP}^{\Delta U=1}(K^+K^{*-}) = 0, \tag{57}$$

$$a_{\rm CP}^{\Delta U=1}(K^0 \rho^+) - a_{\rm CP}^{\Delta U=1}(\overline{K}^0 K^{*+}) = 0, \tag{58}$$

$$a_{\rm CP}^{\Delta U=1}(\pi^+ K^{*0}) - a_{\rm CP}^{\Delta U=1}(K^+ \overline{K}^{*0}) = 0, \tag{59}$$

$$\delta_{\rm CP}^{\Delta U=1}(\pi^0 K^{*+}) + \delta_{\rm CP}^{\Delta U=1}(\eta_8 K^{*+}) - \delta_{\rm CP}^{\Delta U=1}(\pi^0 \rho^+)$$

$$-\delta_{\rm CP}^{\Delta U=1}(\eta_8 \rho^+) = 0, \tag{60}$$

$$\delta_{\rm CP}^{\Delta U=1}(K^+\rho^0) + \delta_{\rm CP}^{\Delta U=1}(K^+\omega_8) - \delta_{\rm CP}^{\Delta U=1}(\pi^+\rho^0) - \delta_{\rm CP}^{\Delta U=1}(\pi^+\omega_8) = 0.$$
(61)

The other sum rules can be obtained by multiplying the individual $\delta_{\rm CP}$ with the corresponding χ , see Eq. (24). In the case of the $D \to PV$ decays no sum rule holds for $\Delta U = 0$ and 1 simultaneously.

IV. EXTENDED SUM RULE

Since except for $a_{\rm CP}(\pi^-\pi^+)$ all CP asymmetries [41] are currently measured consistent with zero (see, table IV of Appendix C), it is difficult to test sum rules at the present stage. However, in the future Belle II and LHCb will reduce uncertainties by a factor of ~ 5 - 10 compared to the current measurements [2, 41-43] and could find more hints of CP violation.

To facilitate these discoveries we will next define differences $\Delta a_{\rm CP}$ of CP asymmetries in such a way that experimental production and detection asymmetries drop out. In the past such considerations led to the measurement of $\Delta a_{\rm CP}$ in Eq. (1). Our new $\Delta a_{\rm CP}$ combine each SCS CP asymmetry with another one in a Cabbibo favored (CF) and doubly Cabbibo suppressed (DCS) decays. Sizable NP contributions to CF decays are not possible and only contrived models can generate a CP asymmetry in DCS decays [44] (see also Ref. [45]). Therefore the $\Delta a_{\rm CP}$'s obey the same sum rules as the corresponding SCS CP asymmetry. out of SCS decay and CF or DCS decay, which can be measured well due to the cancellation of experimental uncertainties such as tagging and hence will provide another important cross check. We find

$$\Delta a_{\rm CP,1}(D_s^+) = a_{\rm CP}(K^0\pi^+) - a_{\rm CP,CF}(\overline{K}^0K^+), \quad (62)$$

$$\Delta a_{\rm CP,2}(D_s^+) = a_{\rm CP}(K^0\pi^+) - a_{\rm CP,DCS}(K^0K^+), \quad (63)$$

$$\Delta a_{\rm CP,3}(D^+) = a_{\rm CP}(\overline{K}^0 K^+) - a_{\rm CP,CF}(\overline{K}^0 \pi^+), \quad (64)$$

$$\Delta a_{\rm CP,4}(D^+) = a_{\rm CP}(\overline{K}^0 K^+) - a_{\rm CP,DCS}(K^0 \pi^+), \quad (65)$$

for $D \to PP$. In reality, one does not observe a K^0 or \bar{K}^0 , but a pair of two pions with the invariant mass of a kaon, i.e. a final state which approximately corresponds to a K_S . One must therefore subtract the effect of kaon

CP violation from the data [46]. This feature also leads to an interference in the CF and DCS decays, for example, the CP asymmetries for $D_s^+ \to K_S K^+$ are non-zero. The resulting CP asymmetries are proportional to the imaginary part of the ratio $V_{cd}^* V_{us}/V_{cs}^* V_{ud}$ in the SM and furthermore unlikely to be large even in the presence of NP [44] and thus negligible compared to the SCS CP asymmetries of interest.

As a result, these sum rules turn out to be very powerful because within the SM and SCS NP scenarios, the CP asymmetries for CF and DCS decays are highly suppressed. Thus these differences essentially coincide with the CP asymmetries in the SCS decays of $D_s^+ \to K^0 \pi^+$ and $D^+ \to \overline{K}^0 K^+$. As said, the differences $\Delta a_{\rm CP}$ in Eqs. (62)-(65) are only taken for experimental reasons to eliminate the production asymmetries of D^+ , D_s^+ , which can fake CP asymmetries. Thus we expect that the $\Delta a_{\rm CP,j}$'s in Eqs. (62)–(65) can be measured more precisely than the single CP asymmetries. But DCS decays might pose additional challenges since the amplitudes are further CKM suppressed and hence these decays are more difficult to access experimentally, so that $\Delta a_{\rm CP,1}$ and $\Delta a_{\rm CP,3}$ with CF-decays might be easier to measure.

Similarly we can construct further $\Delta a_{\rm CP}$ observables for $D \to PV$ decays as

$$\begin{aligned} \Delta a_{\rm CP,5}(D_s^+) &= a_{\rm CP}(K^{*0}\pi^+) - a_{\rm CP,DCS}(K^0K^{*+}), \ (66) \\ \Delta a_{\rm CP,6}(D_s^+) &= a_{\rm CP}(K^{*0}\pi^+) - a_{\rm CP,DCS}(K^{*0}K^+), \ (67) \\ \Delta a_{\rm CP,7}(D_s^+) &= a_{\rm CP}(K^0\rho^+) - a_{\rm CP,DCS}(K^0K^{*+}), \ (68) \\ \Delta a_{\rm CP,8}(D_s^+) &= a_{\rm CP}(K^0\rho^+) - a_{\rm CP,DCS}(K^{*0}K^+), \ (69) \\ \Delta a_{\rm CP,9}(D^+) &= a_{\rm CP}(\overline{K}^{*0}K^+) - a_{\rm CP,CF}(\overline{K}^{*0}\pi^+), \ (70) \\ \Delta a_{\rm CP,10}(D^+) &= a_{\rm CP}(\overline{K}^{*0}K^+) - a_{\rm CP,CF}(\overline{K}^{*0}\pi^+), \ (71) \\ \Delta a_{\rm CP,11}(D^+) &= a_{\rm CP}(\overline{K}^0K^{*+}) - a_{\rm CP,CF}(\overline{K}^{*0}\pi^+), \ (72) \\ \Delta a_{\rm CP,12}(D^+) &= a_{\rm CP}(\overline{K}^{*0}K^+) - a_{\rm CP,DCS}(K^0\rho^+), \ (73) \\ \Delta a_{\rm CP,13}(D^+) &= a_{\rm CP}(\overline{K}^{*0}K^+) - a_{\rm CP,DCS}(\overline{K}^{*0}\pi^+), \ (74) \\ \Delta a_{\rm CP,14}(D^+) &= a_{\rm CP}(\overline{K}^{*0}K^+) - a_{\rm CP,CF}(\overline{K}^{*0}\rho^+), \ (75) \\ \Delta a_{\rm CP,15}(D^+) &= a_{\rm CP}(\overline{K}^{0}K^{*+}) - a_{\rm CP,CF}(\overline{K}^{0}\rho^+), \ (76) \\ \Delta a_{\rm CP,16}(D^+) &= a_{\rm CP}(\overline{K}^{0}K^{*+}) - a_{\rm CP,CF}(\overline{K}^{0}\rho^+). \ (77) \end{aligned}$$

The comments made for $D \to PP$ decays also apply to $\Delta a_{\rm CP,5-16}$, potential CP asymmetries in the CF and DCS decays can be neglected. Note that in $\Delta a_{\rm CP,6}$ and $\Delta a_{\rm CP,9}$ also the $K^{*0} \to K^+\pi^-$ detection asymmetry cancels.

Precise measurements of these $\Delta a_{\rm CP,j}$'s will serve to test the sum rules in Eq. (38), (58) and (59) for scenario II with $|\Delta U| = 1$ NP and $a_0 = 0$. As a result, in total, there are only three independent values of $\Delta a_{\rm CP}$, e.g. $\Delta a_{\rm CP,1}$, $\Delta a_{\rm CP,5}$ and $\Delta a_{\rm CP,7}$. One finds

$$\Delta a_{\rm CP,1} = \Delta a_{\rm CP,2} = \Delta a_{\rm CP,3} = \Delta a_{\rm CP,4}$$
(78)
$$\Delta a_{\rm CP,5} = \Delta a_{\rm CP,6} = \Delta a_{\rm CP,9} = \Delta a_{\rm CP,10}$$

$$a_{\rm CP,5} = \Delta a_{\rm CP,6} = \Delta a_{\rm CP,9} = \Delta a_{\rm CP,10}$$
$$= \Delta a_{\rm CP,13} = \Delta a_{\rm CP,14}, \tag{79}$$

$$\Delta a_{\rm CP,7} = \Delta a_{\rm CP,8} = \Delta a_{\rm CP,11} = \Delta a_{\rm CP,12}$$
$$= \Delta a_{\rm CP,15} = \Delta a_{\rm CP,16}.$$
(80)

for vanishing CP asymmetries in CF and DCS decays. These relations can be useful to test the experimental consistency and the flavor structure of NP.

V. CONCLUSION

In this paper we revisited CP violation in hadronic two-body D meson decays, motivated by the LHCb measurements of $a_{\rm CP}(D^0 \to K^+K^-)$ and $a_{\rm CP}(D^0 \to \pi^+\pi^-)$. The data can only be accommodated within the Standard Model if the approximate SU(3)_F symmetry of QCD fails for the penguin matrix elements entering these CP asymmetries and furthermore a yet unknown mechanism enhances the size of the penguin matrix elements in $D^0 \to \pi^+\pi^-$. We have studied the hypothesis that the measured asymmetries are instead dominated by new physics (NP) assuming that SU(3)_F works. To test this hypothesis we invoked two scenarios characterized by the U-spin quantum number of the NP interaction. We have derived U-spin sum rules between different CP asymmetries which can discriminate between our $\Delta U = 0$ and $|\Delta U| = 1$ scenarios, for both $D \to PP$ and $D \to PV$ decays. The second scenario is qualitatively different from the SM case; we find six $|\Delta U| = 1$ sum rules for $D \to PP$ and eight ones for $D \to PV$ decays. This large number of experimentally testable relations will help to discriminate between NP effects and a SM explanation invoking the breakdown of U-spin symmetry. One of our CP sum rules holds for both the $\Delta U = 0$ and $|\Delta U| = 1$ scenarios. This sum rule could be useful to assess the quality of U-spin symmetry irrespective of the presence of NP. We have also proposed to form differences $\Delta a_{\rm CP}$ between the CP asymmetries in the SCS of interest and those in CF and DCS decays to eliminate experimental production and detection asymmetries. To test our $SU(3)_F$ sum rules more precise data and new measurements are important [42, 43].

ACKNOWLEDGEMENTS

The authors thank S. Schacht and M. Lang for valuable discussion and comment. The research in this paper was supported by the BMBF grant 05H21VKKBA, *Theoretische Studien für Belle II und LHCb* and JSPS KAK-ENHI Grant Number 24K22879 and JPJSCCA20200002.

Appendix A: $\mathcal{O}(1)$ Wigner-Eckart invariants of SCS decays

The effective Hamiltonian transforms as the product

$$\bar{\mathbf{3}} \otimes \mathbf{3} \otimes \bar{\mathbf{3}} = (\mathbf{1} \oplus \mathbf{8}) \otimes \bar{\mathbf{3}} = \bar{\mathbf{3}}_1 \oplus \bar{\mathbf{3}}_2 \oplus \mathbf{6} \oplus \bar{\mathbf{15}},\tag{A1}$$

which can be reduced into a direct sum of irreducible representations [30]. For instance, the $\Delta U = 0$ contribution proportional to $(\bar{s}s - \bar{d}d)(\bar{u}c)$ contains no operators of $\bar{\mathbf{3}}$. These thus lead to penguin contributions in $(\bar{s}s + \bar{d}d)(\bar{u}c)$. Thanks to the Wigner-Eckart theorem, we can systematically express the symmetry properties of the final and initial states as well as the Hamilton operator, and reduce the number of free parameters in the hadronic matrix elements. For SU(3) it has a similar structure as for SU(2) and it follows

$$\langle P_1 P_2 | \mathcal{H} | D \rangle = \sum_w C_w(D, P_1, P_2) X_w , \qquad (A2)$$

where the CG coefficients C_w are also called Wigner-Eckart invariants and X_w are the reduced matrix elements. We label the Wigner-Eckart invariants as $w = \begin{bmatrix} \mathbf{R} \\ i \end{bmatrix}$, see Ref. [30] for details, where **R** is the generating operator in \mathcal{H} and *i* labels the *i*th reduced element. Indices of meson representation are dropped while they are clear from the corresponding decays. The $D \to PP$ and $D \to PV$ Wigner-Eckart invariants of the SCS decay are summarized in Tab. I as well as Tabs. II and III which are taken from Ref. [30].

Appendix B: Two mode sum rules

Thanks to the Wigner-Eckart theorem we can relate the different decay modes based on the group theoretical decomposition and contraction. Here we explain the relation between the amplitude sum rule and a_{CP} sum rule in the

$PP \mod$	$\left(\begin{bmatrix} 3\\1 \end{bmatrix}, \begin{bmatrix} 3\\2 \end{bmatrix} \right)$	$\left(\begin{bmatrix} 6 \\ 1 \end{bmatrix}, \begin{bmatrix} 15 \\ 1 \end{bmatrix}, \begin{bmatrix} 15 \\ 2 \end{bmatrix} \right)$	$\Delta U = 0$ sum rule	$\Delta U = 1$ sum rule
$D^0 \to K^- K^+$	$\frac{\lambda_b}{4}(0,1)$	$+\frac{\lambda}{2}(1,2,1)$	Eq. (30)	$\operatorname{Eq.}(37)$
$D^0 \to \pi^- \pi^+$	$\frac{\lambda_b}{4}(0,1)$	$-\frac{\lambda}{2}(1,2,1)$	Eq. (30)	$\operatorname{Eq.}(37)$
$D^0 \to \pi^0 \pi^0$	$\frac{\lambda_b}{4}(0,1)$	$\frac{\lambda}{2}(1,2,-1)$	Eq. (34)	Eq. (39)
$D^0 \to \eta_8 \eta_8$	$-\frac{\lambda_b}{12}(2,-3)$	$\frac{\lambda}{2}(1,2,-1)$	Eq. (34)	Eqs. $(39, 40)$
$D^0 \to \eta_8 \pi^0$	$\frac{\lambda_b}{4\sqrt{3}}(1,0)$	$\frac{\lambda}{2\sqrt{3}}(1,2,-1)$	Eq. (34)	Eq.(40)
$D^+ \to \pi^0 \pi^+$	(0, 0)	$\frac{\lambda}{\sqrt{2}}(0,0,1)$	Eq. (33)	Eqs. $(41, 42)$
$D^+ \to \eta_8 \pi^+$	$\frac{\lambda_b}{2\sqrt{6}}(1,0)$	$\frac{\lambda}{\sqrt{6}}(1, -2, -2)$	Eq. (35)	Eq.(41)
$D^+ \to \overline{K}^0 K^+$	$\frac{\dot{\lambda}_b}{4}(1,0)$	$\frac{\lambda}{2}(1,-2,1)$	Eq. (31)	Eq. (38)
$D_s^+ \to \pi^0 K^+$	$\frac{\lambda_b}{4\sqrt{2}}(1,0)$	$-\frac{\lambda}{2\sqrt{2}}(1,-2,-1)$	Eqs. $(35, 36)$	Eqs. $(41, 42)$
$D_s^+ \to K^0 \pi^+$	$\frac{\dot{\lambda}_b}{4}(1,0)$	$-\frac{\lambda}{2}(1,-2,1)$	Eqs. $(31, 36)$	Eqs. (38, 42)
$D_s^+ \to \eta_8 K^+$	$-\frac{\lambda_b}{4\sqrt{6}}(1,0)$	$\frac{\lambda}{2\sqrt{6}}(1,-2,-5)$	Eqs. $(35, 36)$	Eqs. $(41, 42)$

. 1

TABLE I. The SCS decay Wigner-Eckart invariants and occurrence of the sum rules for $D_{(s)} \rightarrow PP$.

PV mode	$\left(\left[\frac{\overline{3}}{1}\right], \left[\frac{\overline{3}}{2}\right], \left[\frac{\overline{3}}{3}\right]\right)$	$\left(\begin{bmatrix} 6 \\ 1 \end{bmatrix}, \begin{bmatrix} 6 \\ 2 \end{bmatrix}, \begin{bmatrix} 6 \\ 3 \end{bmatrix}, \begin{bmatrix} \overline{15} \\ 1 \end{bmatrix}, \begin{bmatrix} \overline{15} \\ 2 \end{bmatrix}, \begin{bmatrix} \overline{15} \\ 3 \end{bmatrix}, \begin{bmatrix} \overline{15} \\ 4 \end{bmatrix} \right)$	$\Delta U=0$ sum rule	$\Delta U = 1$ sum rule
$D^0 \to \eta_8 \omega_8$	$-\frac{\lambda_b}{48}(4,-6,-1)$	$\frac{\lambda}{4}(0,1,-1,2,1,1,-1)$	$\operatorname{Eq.}(50)$	Eq. (54)
$D^0 o \eta_8 ho^0$	$\frac{\lambda_b}{16\sqrt{3}}(2,0,1)$	$-\frac{\lambda}{4\sqrt{3}}(2,1,3,-2,-5,-1,-1)$	Eqs. $(50, 53)$	
$D^0 \to K^0 \overline{K}^{*0}$	$-\frac{\lambda_b}{8}(1,-1,0)$	$\frac{\lambda}{2}(1,0,0,0,1,-1,0)$	$\operatorname{Eq.}(45)$	$\operatorname{Eq.}(55)$
$D^0 \to \pi^0 \omega_8$	$\frac{\lambda_b}{16\sqrt{3}}(2,0,1)$	$\frac{\lambda}{4\sqrt{3}}(2,3,1,2,-3,1,-3)$	Eqs. $(50, 53)$	
$D^0 o \pi^0 \rho^0$	$\frac{\lambda_b}{16}(0,2,1)$	$-rac{\lambda}{4}(0,1,-1,2,1,1,-1)$	$\operatorname{Eq.}(50)$	$\operatorname{Eq.}(54)$
$D^0 \to \overline{K}^0 K^{*0}$	$-\frac{\lambda_b}{8}(1,-1,0)$	$-rac{\lambda}{2}(1,0,0,0,1,-1,0)$	$\operatorname{Eq.}(45)$	$\operatorname{Eq.}(55)$
$D^0 \to K^- K^{*+}$	$\frac{\lambda_{b}}{8}(0,1,0)$	$rac{\lambda}{2}(0,1,0,1,1,1,1)$	$\operatorname{Eq.}(46)$	$\operatorname{Eq.}(56)$
$D^0 \to \pi^- \rho^+$	$\frac{\lambda_b}{8}(0,1,0)$	$-rac{\lambda}{2}(0,1,0,1,1,1,1)$	$\operatorname{Eq.}(46)$	$\operatorname{Eq.}(56)$
$D^0 \to K^+ K^{*-}$	$\frac{\lambda_b}{8}(0,1,1)$	$-rac{\lambda}{2}(0,0,1,-1,0,0,0)$	$\operatorname{Eq.}(47)$	$\operatorname{Eq.}(57)$
$D^0 \to \pi^+ \rho^-$	$\frac{\lambda_b}{8}(0,1,1)$	$\frac{\lambda}{2}(0,0,1,-1,0,0,0)$	$\operatorname{Eq.}(47)$	Eq. (57)

TABLE II. The SCS decay Wigner-Eckart invariants and occurrence of the sum rules for $D^0 \rightarrow PV$.

case where only two decay modes are involved. For the relations involving three or more modes, there is in general no simple formula. We start from the relations

$$P_0^{(P_1 P_2)} = c_P P_0^{(Q_1 Q_2)}, \quad A^{(P_1 P_2)} = c_A A^{(Q_1 Q_2)}, \tag{B1}$$

where c_A and c_P are real coefficients which can be read from the tables in Appendix A. We can simply replace the amplitudes P_0, A_1 in Eq. (29) and find the two-modes sum rule

$$a_{\rm CP}^{\Delta U=0}(P_1 P_2) = \frac{c_P}{c_A} a_{\rm CP}^{\Delta U=0}(Q_1 Q_2).$$
 (B2)

This can of course also be done for the $|\Delta U| = 1$ case but it is always necessary that both P and A obey sum rules as in Eq. (B1). For instance sum rules in Eqs. (30, 31) follow from $c_P = 1$ and $c_A = -1$ as seen from table I.

Appendix C: Current experimental status and future sensitivity

Tab. IV shows the current status and future sensitivity of the CP asymmetries measurements [41-43]. For the future sensitivity an integrated luminosity of 50 ab^{-1} and 300 fb^{-1} is assumed for Belle II and LHCb, respectively.

PV mode	$\left(\left[\begin{array}{c} \overline{3}\\1\end{array}\right], \left[\begin{array}{c} \overline{3}\\2\end{array}\right], \left[\begin{array}{c} \overline{3}\\3\end{array}\right]\right)$	$\left(\begin{bmatrix}6\\1\end{bmatrix},\begin{bmatrix}6\\2\end{bmatrix},\begin{bmatrix}6\\3\end{bmatrix},\begin{bmatrix}15\\1\end{bmatrix},\begin{bmatrix}\mathbf{\overline{15}}\\2\end{bmatrix},\begin{bmatrix}\mathbf{\overline{15}}\\3\end{bmatrix},\begin{bmatrix}\mathbf{\overline{15}}\\4\end{bmatrix}\right)$	$\Delta U = 0$ sum rule	$\Delta U = 1$ sum rule
$D^+ \to \eta_8 \rho^+$	$\frac{\lambda_b}{8\sqrt{6}}(2,0,1)$	$-rac{\lambda}{2\sqrt{6}}(2,1,3,2,-1,1,1)$	Eqs. $(51, 53)$	Eq. (60)
$D^+ \to \pi^0 \rho^+$	$\frac{\lambda_b}{8\sqrt{2}}(0,0,1)$	$\frac{\lambda}{2\sqrt{2}}(0,1,1,0,1,-1,1)$	Eqs. $(51, 53)$	Eq. (60)
$D^+ \to \overline{K}^0 K^{*+}$	$\frac{\lambda_b}{8}(1,0,0)$	$rac{\lambda}{2}(1,1,0,-1,0,0,1)$	Eqs. (48, 53)	$\operatorname{Eq.}(58)$
$D^+ \to K^+ \overline{K}^{*0}$	$\frac{\lambda_b}{8}(1,0,1)$	$-rac{\lambda}{2}(1,0,1,1,1,1,0)$	Eqs. $(49, 53)$	$\operatorname{Eq.}(59)$
$D^+ \to \pi^+ \omega_8$	$\frac{\lambda_b}{8\sqrt{6}}(2,0,1)$	$\frac{\lambda}{2\sqrt{6}}(2,3,1,-2,-3,-1,-3)$	Eqs. $(52, 53)$	Eq. (61)
$D^+ \to \pi^+ \rho^0$	$-\frac{\lambda_b}{8\sqrt{2}}(0,0,1)$	$-rac{\lambda}{2\sqrt{2}}(0,1,1,0,1,-1,-1)$	Eqs. $(52, 53)$	Eq. (61)
$D_s^+ \to \eta_8 K^{*+}$	$-\frac{\lambda_b}{8\sqrt{6}}(1,0,-1)$	$-rac{\lambda}{2\sqrt{6}}(1,2,3,1,1,-1,2)$	Eqs. $(51, 53)$	Eq. (60)
$D_s^+ \to K^0 \rho^+$	$\frac{\lambda_b}{8}(1,0,0)$	$-rac{\lambda}{2}(1,1,0,-1,0,0,1)$	Eq. (48)	$\operatorname{Eq.}(58)$
$D_s^+ \to \pi^0 K^{*+}$	$\frac{\lambda_b}{8\sqrt{2}}(1,0,1)$	$rac{\lambda}{2\sqrt{2}}(1,0,1,1,-1,1,0)$	$\operatorname{Eq.}(51)$	Eq. (60)
$D_s^+ \to K^+ \omega_8$	$-\frac{\lambda_b}{8\sqrt{6}}(1,0,2)$	$\frac{\lambda}{2\sqrt{6}}(1,3,2,-1,0,-2,-3)$	Eq.(52)	Eq. (61)
$D_s^+ \to K^+ \rho^0$	$\frac{\lambda_b}{8\sqrt{2}}(1,0,0)$	$-rac{\lambda}{2\sqrt{2}}(1,1,0,-1,-2,0,-1)$	Eqs. $(52, 53)$	Eq. (61)
$D_s^+ \to \pi^+ K^{*0}$	$\frac{\lambda_b}{8}(1,0,1)$	$\frac{\lambda}{2}(1,0,1,1,1,1,0)$	Eq. (49)	Eq. (59)

TABLE III. The SCS decay Wigner-Eckart invariants and occurrence of the sum rules for $D^+_{(s)} \to PV$.

Decay Mode	PDG $a_{\rm CP}$ [%]	Belle	Belle II $(50 \mathrm{ab}^{-1})$	LHCb	LHCb $(300 {\rm fb}^{-1})$	$\Delta U = 0$	$\Delta U = 1$
$D^0 \to K^+ K^-$	-0.07 ± 0.11	-0.32 ± 0.23	± 0.03	0.077 ± 0.057	± 0.007	Eq. (30)	Eq. (37)
$D^0 \to \pi^+ \pi^-$	0.13 ± 0.14	0.55 ± 0.37	± 0.05	0.232 ± 0.061	± 0.007	Eq. (30)	Eq. (37)
$D^0 \to \pi^0 \pi^0$	0.0 ± 0.6	-0.03 ± 0.65	± 0.09	—	—	Eq. (34)	Eq. (39)
$D^+ \to \pi^0 \pi^+$	0.4 ± 1.3	2.31 ± 1.26	± 0.17	-1.3 ± 1.1	—	Eq. (33)	Eqs. $(41, 42)$
$D^+ \to \eta \pi^+$	0.3 ± 0.8	1.74 ± 1.15	± 0.14	-0.2 ± 0.9		Eq. (35)	Eq. (41)
$D^+ \to \eta' \pi^+$	-0.6 ± 0.7	-0.12 ± 1.13	± 0.14	-0.61 ± 0.9	—	Eq. (35)	Eq. (41)
$D^+ \to K^0_s K^+$	-0.01 ± 0.07	-0.25 ± 0.31	± 0.04	-0.004 ± 0.076	—	Eq. (31)	Eq. (38)
$D_s^+ \to K_s^0 \pi^+$	0.20 ± 0.18	5.45 ± 2.52	± 0.29	0.16 ± 0.18		Eqs. (31, 35)	Eqs. (38, 42)
$D_s^+ \to K^+ \eta$	1.8 ± 1.9	2.1 ± 2.1		0.9 ± 3.9	—	Eqs. $(35, 36)$	Eqs. $(41, 42)$
$D_s^+ \to K^+ \eta'$	6.0 ± 18.9				—	Eqs. $(35, 36)$	Eqs. $(41, 42)$

TABLE IV. Current experimental status and future sensitivity taken from Refs. [2, 41–43].

[arXiv:1903.08726].

- [2] LHCb Collaboration, Measurement of the Time-Integrated CP Asymmetry in D0→K-K+ Decays, Phys. Rev. Lett. 131 (2023) 091802 [arXiv:2209.03179].
- [3] N. Cabibbo, Unitary Symmetry and Leptonic Decays, Phys. Rev. Lett. 10 (1963) 531–533.
- [4] M. Kobayashi and T. Maskawa, CP Violation in the Renormalizable Theory of Weak Interaction, Prog. Theor. Phys. 49 (1973) 652–657.
- [5] A. Khodjamirian and A. A. Petrov, Direct CP asymmetry in $D \to \pi^- \pi^+$ and $D \to K^- K^+$ in QCD-based approach, Phys. Lett. B **774** (2017) 235-242 [arXiv:1706.07780].
- [6] M. Chala, A. Lenz, A. V. Rusov, and J. Scholtz, ΔA_{CP} within the Standard Model and beyond, JHEP 07 (2019) 161 [arXiv:1903.10490].
- [7] **LHCb** Collaboration, Evidence for CP violation in time-integrated $D^0 \rightarrow h^-h^+$ decay rates, Phys. Rev. Lett. **108** (2012) 111602 [arXiv:1112.0938].
- [8] Y. Grossman, A. L. Kagan, and Y. Nir, New physics and CP violation in singly Cabibbo suppressed D decays, Phys. Rev. D 75 (2007) 036008 [hep-ph/0609178].
- [9] G. Isidori, J. F. Kamenik, Z. Ligeti, and G. Perez, Implications of the LHCb Evidence for Charm CP

Violation, Phys. Lett. B **711** (2012) 46–51 [arXiv:1111.4987].

- [10] Y. Grossman, A. L. Kagan, and J. Zupan, Testing for new physics in singly Cabibbo suppressed D decays, Phys. Rev. D85 (2012) 114036 [arXiv:1204.3557].
- [11] G. F. Giudice, G. Isidori, and P. Paradisi, Direct CP violation in charm and flavor mixing beyond the SM, JHEP 04 (2012) 060 [arXiv:1201.6204].
- [12] G. Hiller, Y. Hochberg, and Y. Nir, Supersymmetric ΔA_{CP}, Phys. Rev. D 85 (2012) 116008 [arXiv:1204.1046].
- [13] W. Altmannshofer, R. Primulando, C.-T. Yu, and F. Yu, New Physics Models of Direct CP Violation in Charm Decays, JHEP 04 (2012) 049 [arXiv:1202.2866].
- [14] C.-H. Chen, C.-Q. Geng, and W. Wang, Direct CP Violation in Charm Decays due to Left-Right Mixing, Phys. Lett. B 718 (2013) 946–950 [arXiv:1206.5158].
- [15] B. Keren-Zur, et al., On Partial Compositeness and the CP asymmetry in charm decays, Nucl. Phys. B 867 (2013) 394-428 [arXiv:1205.5803].
- [16] C. Delaunay, J. F. Kamenik, G. Perez, and L. Randall, Charming CP Violation and Dipole Operators from RS Flavor Anarchy, JHEP 01 (2013) 027 [arXiv:1207.0474].

- [17] A. Dery and Y. Nir, Implications of the LHCb discovery of CP violation in charm decays, JHEP 12 (2019) 104 [arXiv:1909.11242].
- [18] L. Calibbi, T. Li, Y. Li, and B. Zhu, Simple model for large CP violation in charm decays, B-physics anomalies, muon g - 2 and dark matter, JHEP 10 (2020) 070 [arXiv:1912.02676].
- [19] R. Bause, H. Gisbert, M. Golz, and G. Hiller, *Exploiting CP-asymmetries in rare charm decays*, Phys. Rev. D 101 (2020) 115006 [arXiv:2004.01206].
- [20] A. J. Buras, P. Colangelo, F. De Fazio, and F. Loparco, *The charm of 331*, JHEP **10** (2021) 021 [arXiv:2107.10866].
- [21] R. Bause, et al., U-spin-CP anomaly in charm, Phys. Rev. D 108 (2023) 035005 [arXiv:2210.16330].
- [22] J. Brod, Y. Grossman, A. L. Kagan, and J. Zupan, A Consistent Picture for Large Penguins in $D \rightarrow \pi^+\pi^-, K^+K^-$, JHEP **10** (2012) 161 [arXiv:1203.6659].
- [23] Y. Grossman and S. Schacht, The emergence of the $\Delta U = 0$ rule in charm physics, JHEP **07** (2019) 020 [arXiv:1903.10952].
- [24] A. Soni, Resonance enhancement of Charm CP. arXiv: 1905.00907.
- [25] S. Schacht and A. Soni, Enhancement of charm CP violation due to nearby resonances, Phys. Lett. B 825 (2022) 136855 [arXiv:2110.07619].
- [26] D. Pirtskhalava and P. Uttayarat, CP Violation and Flavor SU(3) Breaking in D-meson Decays, Phys. Lett. B712 (2012) 81-86 [arXiv:1112.5451].
- [27] B. Bhattacharya, M. Gronau, and J. L. Rosner, CP asymmetries in singly-Cabibbo-suppressed D decays to two pseudoscalar mesons, Phys. Rev. D 85 (2012) 054014 [arXiv:1201.2351].
- [28] T. Feldmann, S. Nandi, and A. Soni, Repercussions of Flavour Symmetry Breaking on CP Violation in D-Meson Decays, JHEP 06 (2012) 007 [arXiv:1202.3795].
- [29] B. Bhattacharya, M. Gronau, and J. L. Rosner, Direct CP Violation in D Decays in view of LHCb and CDF Results. arXiv: 1207.0761.
- [30] Y. Grossman and D. J. Robinson, SU(3) Sum Rules for Charm Decay, JHEP 04 (2013) 067 [arXiv:1211.3361].
- [31] G. Hiller, M. Jung, and S. Schacht, SU(3)-flavor anatomy of nonleptonic charm decays, Phys. Rev. D 87 (2013) 014024 [arXiv:1211.3734].

- [32] Y. Grossman, Z. Ligeti, and D. J. Robinson, More Flavor SU(3) Tests for New Physics in CP Violating B Decays, JHEP 01 (2014) 066 [arXiv:1308.4143].
- [33] G. Hiller, M. Jung, and S. Schacht, SU(3)_F in nonleptonic charm decays, PoS EPS-HEP2013 (2013) 371 [arXiv:1311.3883].
- [34] S. Müller, U. Nierste, and S. Schacht, Topological amplitudes in D decays to two pseudoscalars: A global analysis with linear SU(3)_F breaking, Phys. Rev. D 92 (2015) 014004 [arXiv:1503.06759].
- [35] S. Müller, U. Nierste, and S. Schacht, Sum Rules of Charm CP Asymmetries beyond the SU(3)_F Limit, Phys. Rev. Lett. **115** (2015) 251802 [arXiv:1506.04121].
- [36] M. Gavrilova, Y. Grossman, and S. Schacht, The mathematical structure of U-spin amplitude sum rules, JHEP 08 (2022) 278 [arXiv:2205.12975].
- [37] S. Schacht, A U-spin anomaly in charm CP violation, JHEP 03 (2023) 205 [arXiv:2207.08539].
- [38] U. Nierste and S. Schacht, Neutral $D \to KK^*$ decays as discovery channels for charm CP violation, Phys. Rev. Lett. **119** (2017) 251801 [arXiv:1708.03572].
- [39] J. J. de Swart, The Octet model and its Clebsch-Gordan coefficients, Rev. Mod. Phys. 35 (1963) 916–939.
 [Erratum: Rev.Mod.Phys. 37, 326–326 (1965)].
- [40] T. A. Kaeding, *Tables of SU(3) isoscalar factors*, Atom. Data Nucl. Data Tabl. **61** (1995) 233-288
 [nucl-th/9502037].
- [41] Particle Data Group Collaboration, Review of Particle Physics, PTEP 2022 (2022) 083C01.
- [42] Belle-II Collaboration, The Belle II Physics Book, PTEP 2019 (2019) 123C01 [arXiv:1808.10567].
 [Erratum: PTEP 2020, 029201 (2020)].
- [43] LHCb Collaboration, Physics case for an LHCb Upgrade II - Opportunities in flavour physics, and beyond, in the HL-LHC era. arXiv:1808.08865.
- [44] S. Bergmann and Y. Nir, New physics effects in doubly Cabibbo suppressed D decays, JHEP 09 (1999) 031 [hep-ph/9909391].
- [45] I. I. Y. Bigi and H. Yamamoto, Interference between Cabibbo allowed and doubly forbidden transitions in D -> K(S), K(L) + pi's decays, Phys. Lett. B 349 (1995) 363-366 [hep-ph/9502238].
- [46] Y. Grossman and Y. Nir, *CP Violation in* $\tau^{\pm} \to \pi^{\pm} K_S \nu$ and $D^{\pm} \to \pi^{\pm} K_S$: The Importance of $K_S - K_L$ Interference, JHEP **04** (2012) 002 [arXiv:1110.3790].