

PREPARED FOR SUBMISSION TO JHEP

TIF-UNIMI-2025-8, TTP25-006, P3H-25-017, MPP-2025-40, SLAC-PUB-250317

Towards a general subtraction formula for NNLO QCD corrections to processes at hadron colliders: final states with quarks and gluons

**Federica Devoto,^a Kirill Melnikov,^b Raoul Röntschi,^c Chiara Signorile-Signorile,^d
Davide Maria Tagliabue,^{b,c} Matteo Tresoldi^b**

^a*SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94039, USA*

^b*Institute for Theoretical Particle Physics, KIT, Wolfgang-Gaede-Straße 1, 76131, Karlsruhe, Germany*

^c*Tif Lab, Dipartimento di Fisica, Università di Milano and INFN, Sezione di Milano, Via Celoria 16, I-20133 Milano, Italy*

^d*Max-Planck-Institut für Physik, Boltzmannstrasse 8, 85748 Garching, Germany*

E-mail: federica@slac.stanford.edu, kirill.melnikov@kit.edu,
raoul.rontsch@unimi.it, signoril@mpp.mpg.de,
davide.tagliabue@kit.edu, matteo.tresoldi@partner.kit.edu

ABSTRACT: We describe the calculation of integrated subtraction terms in the nested soft-collinear subtraction scheme for hadron collider processes with quarks and gluons, thereby extending the results presented in Ref. [1]. Although this extension eventually proves to be straightforward, it requires a more careful treatment of certain collinear limits to achieve a compact and physically-transparent final result. We also show that the cancellation of infrared divergences can be organized in such a way that, once soft contributions are removed, it occurs independently for each of the external partons. We consider these results to be important stepping stones on the way to deriving finite remainders of the integrated subtraction terms for fully-general hadron collider processes in the context of the nested soft-collinear subtraction scheme.

KEYWORDS: QCD corrections, hadronic colliders, NNLO calculations

Contents

1	Introduction	1
2	The leading order process	3
3	NLO QCD corrections	4
4	NNLO QCD corrections: general framework	10
5	Simplifications of the double-unresolved contributions	16
5.1	Simplifying virtual and soft corrections	17
5.2	Soft-regulated double-collinear contributions	19
6	Cancellation of poles	24
6.1	Single-unresolved terms	24
6.2	Double-boosted contribution	26
6.3	Single-boosted contributions	26
6.4	Unboosted terms	28
7	Finite remainder of the NNLO contribution to the cross section	32
8	Conclusions	36
A	Collection of relevant constant, functions and operators	38
A.1	Useful constants	39
A.2	Tree-level, one-loop and Altarelli-Parisi splitting functions	41
A.3	Generalized splitting functions and anomalous dimensions	42
A.3.1	Tree-level	42
A.3.2	Spin-correlations	44
A.3.3	One-loop	45
A.4	Definitions and relevant properties of I -operators	46
A.5	Double-unresolved contributions	48

1 Introduction

The reliable theoretical description of particle production processes through high-energy collisions is an essential element of the Large Hadron Collider (LHC) physics program, which aims at probing the Standard Model (SM) of particle physics at the shortest distance scales. Providing these descriptions requires the development of techniques to calculate multi-loop amplitudes, as well as those to organize the cancellation of infrared (IR)

singularities which appear at intermediate stages of the computation. With the large and ever-increasing dataset from the LHC allowing us to probe ever more complicated processes, these techniques should be broadly applicable – i.e. they should be able to accommodate processes with a large number of final state particles, including jets – as well as amenable to efficient implementation in a numerical code.

In this paper, we focus on the problem of IR singularities. While efficient, process-independent treatments of IR singularities at next-to-leading order (NLO) in QCD were developed many years ago [2–5], no fully general solution to this problem is available at next-to-next-to-leading order (NNLO) in spite of the significant progress achieved in the past twenty years. Indeed, during that time several NNLO subtraction and slicing schemes were proposed [1, 6–33], and applied to study many interesting processes at lepton and hadron colliders (see e.g. Refs [34–58] for a selection of phenomenological papers employing different theoretical methods). However, none solve the problem as comprehensively as at NLO. To illustrate this point, we emphasize that, as of today, none of the fully-local NNLO subtraction schemes has been used to demonstrate the cancellation of IR singularities and provide formulas for the finite remainders for processes with an arbitrary number of jets at a hadron collider.

In view of this, many of the NNLO subtraction schemes are being actively developed and refined. In Ref. [28], the *local analytic sector subtraction* scheme [25, 26] was used to demonstrate the cancellation of IR poles for an arbitrary final state produced in e^+e^- collisions. Similarly, important recent advances have been made in the context of *antenna subtractions* [27, 30–33, 59], which streamline the computation of antenna functions as well as broaden their applicability to include sub-leading color effects. Improved treatments of power corrections have been proposed for both the q_T slicing [60–64] and N -jettiness slicing [62, 65–69] methods. There has also been work towards extending local subtraction methods [70–73] and slicing methods [74–86] to N³LO.

One of the main obstacles in constructing a generalized NNLO subtraction scheme is the proliferation of terms required at different stages of the calculation and the need to combine multiple contributions to obtain the physically-transparent structure of the final result. In the recent paper [1], we employed the so-called *nested soft-collinear subtraction* scheme [17] to show how this problem can be overcome. The key element in the construction described in Ref. [1] is the iterative nature of IR subtraction terms that emerges at the level of color-correlated matrix elements. Using this feature as a guiding principle allows us to treat the problem in an almost process-independent way, and significantly simplifies the intermediate steps required to demonstrate the cancellation of IR poles. As a result, it became possible to show the cancellation of these poles for the process $q\bar{q} \rightarrow X + N_g g$, where the number of final-state gluons N_g is a parameter, and to produce relatively compact formulas for the finite remainder of the integrated subtraction terms.

Although the results reported in Ref. [1] are very general, we did take advantage of the symmetries of the final state and the simple structure of the collinear limits between final-state gluons and incoming partons. Hence, to generalize the calculation of Ref. [1] to an arbitrary process, we need to do two things: first, consider less symmetric final states and, second, account for flavor-changing initial-state collinear emissions. The goal

of this paper is to address the first issue and we do this by considering a process where an initial gq state produces an arbitrary number of gluons and a quark. Furthermore, we also consider corrections in which an additional quark-antiquark pair is radiated, and extract the contributions to the integrated subtraction terms that depend on the number of light quark flavors n_f . We show that the general approach presented in Ref. [1] can be applied to such processes with small modifications which mostly concern the interconnection of final-state collinear splittings as these may be affected by the choice of unresolved partons in different ways. We find that, if the subtraction terms described by these collinear limits are combined before integration, they lead to universal, physically-transparent structures. In fact, these features can already be seen at NLO, providing a guide as to how to solve this problem at NNLO.

In general, we find that the method introduced in Ref. [1] is sufficiently robust to provide the description of more complex final and initial states. Indeed, as we will see, the soft part is unchanged, while the collinear part can be treated leg-by-leg. Hence, in spite of the fact that we still work with a particular partonic process, we believe that the present paper is an important step towards providing integrated NNLO subtraction terms and finite remainders for *generic* multi-jet processes at hadron and lepton colliders.

The remainder of the paper is organized as follows. In Sec. 2, we introduce the process of interest and the notation. In Sec. 3, we present the treatment of NLO QCD corrections, and discuss our approach to the issues mentioned above. In Sec. 4, we consider the NNLO corrections, and show that the relevant soft and collinear limits can be manipulated to identify contributions with a well-defined number of resolved partons. We discuss the simplification of these contributions in Sec. 5. Once this is done, we demonstrate the cancellation of IR singularities in Sec. 6, and present the finite remainder in Sec. 7. We conclude in Sec. 8. We collect many of the constants, functions, and operators used throughout the paper in Appendix A.

2 The leading order process

We are interested in understanding higher-order QCD corrections to the inclusive production of N jets at a hadron collider in association with a color-neutral system X , $pp \rightarrow X + N$ jets. This process receives contributions from many partonic channels. In this paper, we will focus on the following one

$$\mathcal{A}_0 : a_g + b_q \rightarrow X + N_g g + q, \quad (2.1)$$

where $N_g = N - 1$, and a and b label initial state partons with momenta $p_{a,b}$. We write the cross section for process \mathcal{A}_0 as

$$\begin{aligned} 2s_{ab} d\hat{\sigma}_{\text{LO}}^{gq} &= \left\langle \frac{1}{N_g!} F_{\text{LM}}^{gq}[\{g\}_{N_g}, \{q\}_1] \right\rangle \\ &= \frac{N_{\text{av}}}{N_g!} \int d\Phi (2\pi)^4 \delta^{(4)}(P_{\text{fin}} - P_{\text{in}}) |\mathcal{M}_0(P_{\text{fin}}, P_{\text{in}})|^2 \mathcal{O}(P_{\text{fin}}), \end{aligned} \quad (2.2)$$

and we note that, in order to obtain its contribution to the hadronic process, we need to convolute the partonic cross section with parton distribution functions (PDF) f_g and f_q . In

Eq. (2.2), \mathcal{M}_0 is the matrix element of the process in Eq. (2.1), $d\Phi$ is the Lorentz-invariant phase space for the final-state particles, and \mathcal{O} is an IR-safe observable which ensures that the final state contains at least N resolved jets. The total momenta of the initial- and final-state particles are denoted as P_{in} and P_{fin} , respectively, and the delta function in Eq. (2.2) enforces energy-momentum conservation. Summation over spins and colors of final-state partons, and averaging over spins and colors of initial-state partons are assumed in Eq. (2.2). The corresponding spin and color averaging factors are described by the factor N_{av} .

Furthermore, $\{g\}_{N_g}$ and $\{q\}_1$ in Eq. (2.2) denote lists of N_g final-state gluons, and of a single final-state quark, respectively. This notation is redundant for the process that we consider, but we introduce it with an eye on future generalizations. The factor $1/N_g!$ corresponds to the symmetry factor arising from having N_g indistinguishable gluons in the final state. It is convenient to absorb this symmetry factor and omit the initial-state partons and lists of final-state partons by defining

$$F_{\text{LM}}^{\mathcal{A}_0} = \frac{1}{N_g!} F_{\text{LM}}^{gq}[\{g\}_{N_g}, \{q\}_1]. \quad (2.3)$$

Then, the differential cross section for the process \mathcal{A}_0 reads

$$2s_{ab} d\hat{\sigma}_{\text{LO}}^{gq} = \langle F_{\text{LM}}^{\mathcal{A}_0} \rangle, \quad (2.4)$$

and we will use this notation in what follows.

3 NLO QCD corrections

Having fixed the notation, we continue by discussing the NLO QCD corrections to the partonic process \mathcal{A}_0 in Eq. (2.1). As usual, one has to compute virtual corrections to \mathcal{A}_0 and account for real-emission contributions that are comprised of processes where the number of final-state partons is increased by one. For the complete calculation, many real-emission processes need to be considered. However, since we would like to discuss the cancellation of IR singularities in higher-order corrections to \mathcal{A}_0 , we can omit contributions that do not lead to this process when soft or collinear limits are taken. There are also processes which lead to \mathcal{A}_0 under collinear limits but which are convoluted with different PDFs; we do not consider such processes either.

With this clarification, it is easy to convince oneself that the following three processes should be considered

$$\begin{aligned} \mathcal{A}_1 : \quad & a_g + b_q \rightarrow X + (N_g + 1)g + q, \\ \mathcal{A}_2 : \quad & a_g + b_q \rightarrow X + (N_g - 1)g + q + q'\bar{q}', \quad q' \neq q, \\ \mathcal{A}_3 : \quad & a_g + b_q \rightarrow X + (N_g - 1)g + q + q\bar{q}. \end{aligned} \quad (3.1)$$

However, further simplifications are possible. Indeed, in order to provide a singular contribution to the process \mathcal{A}_0 , each of the three processes $\mathcal{A}_{1,2,3}$ should “lose” a parton in such a way that one gets both the initial and the final state of \mathcal{A}_0 . There are several ways in

which this can happen in the process \mathcal{A}_1 , but in processes \mathcal{A}_2 and \mathcal{A}_3 this can only occur if a quark-antiquark pair, $q'\bar{q}'$ or $q\bar{q}$, respectively, is clustered into a gluon. Since these collinear limits are identical for processes \mathcal{A}_2 and \mathcal{A}_3 , we do not need to consider the latter since we can retrieve all the required information from the former.

We begin with the discussion of process \mathcal{A}_1 in Eq. (3.1), and write the corresponding differential cross section as

$$2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_1} = \left\langle \frac{1}{(N_g + 1)!} F_{\text{LM}}^{gq}[\{g\}_{N_g+1}, \{q\}_1] \right\rangle. \quad (3.2)$$

Following Ref. [1], we introduce partitions of unity $\Delta^{(i)}$ and split the cross section into a sum of terms such that in each term only one of the partons can lead to a singular limit once it becomes unresolved. We find

$$2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_1} = \sum_i \left\langle \frac{\Delta^{(i)}}{(N_g + 1)!} F_{\text{LM}}^{gq}[\{g\}_{N_g+1}, \{q\}_1] \right\rangle. \quad (3.3)$$

The index i in the above formula runs over all final-state partons, and a possible choice of partitions $\Delta^{(i)}$ is described in Appendix B of Ref. [1].

The potentially-unresolved parton i can be either a (anti)quark or a gluon. Accounting for the symmetry properties of F_{LM} , we rename the final state momenta and call the unresolved parton \mathbf{m} . There are $N_g + 1$ ways to choose an unresolved gluon and a single choice for an unresolved quark. Hence, we can write

$$2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_1} = \langle \Delta^{(\mathbf{m})} F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle, \quad (3.4)$$

where

$$F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] = F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q], \quad (3.5)$$

and the functions F_{LM} are defined as

$$\begin{aligned} F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] &= \frac{1}{N_g!} F_{\text{LM}}^{ab}[\{g\}_{N_g}, \{q\}_1 | \mathbf{m}_g], \\ F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] &= \frac{1}{(N_g + 1)!} F_{\text{LM}}^{ab}[\{g\}_{N_g+1} | \mathbf{m}_q]. \end{aligned} \quad (3.6)$$

We note that in the last F_{LM} function the quark list has disappeared, and that the symmetry factors that multiply various functions F_{LM} are defined by the number of *resolved* gluons.

Following the discussion in Ref. [1], we extract soft and collinear divergences that affect the real emission contribution, and obtain

$$2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_1} = \langle S_{\mathbf{m}} F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle + \sum_{i \in \mathcal{H}} \langle \bar{S}_{\mathbf{m}} C_{i\mathbf{m}} \Delta^{(\mathbf{m})} F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle + \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \Delta^{(\mathbf{m})} F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle. \quad (3.7)$$

We denote the list of N final-state resolved partons contained in a given F_{LM} as \mathcal{H}_{f} , and the list that combines \mathcal{H}_{f} and the initial-state particles as $\mathcal{H} = \{a, b\} \cup \mathcal{H}_{\text{f}}$. The soft and collinear subtraction operators in Eq. (3.7) read

$$\bar{S}_{\mathbf{m}} = \mathbb{1} - S_{\mathbf{m}}, \quad \bar{C}_{i\mathbf{m}} = \mathbb{1} - C_{i\mathbf{m}}, \quad \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} = \sum_{i \in \mathcal{H}} \bar{S}_{\mathbf{m}} \bar{C}_{i\mathbf{m}} \omega^{\mathbf{m}i}. \quad (3.8)$$

The partition functions $\omega^{\mathbf{m}i}$ are defined in Ref. [1], and allow us to treat one collinear singularity at a time. We note that, in Eq. (3.7), the soft-limit operator $S_{\mathbf{m}}$ is only needed when \mathbf{m} is a gluon. Hence, to apply Eq. (3.7) for generic processes, we have to set the soft operator to zero when the unresolved parton is a quark or an antiquark, i.e.,

$$S_q = S_{\bar{q}} = 0. \quad (3.9)$$

Upon integrating over the phase space of the soft gluon, the first term on the right-hand side of Eq. (3.7) can be written as¹

$$\langle S_{\mathbf{m}} F_{\mathbf{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle \equiv \langle S_{\mathbf{m}} F_{\mathbf{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \rangle = [\alpha_s] \langle I_S(\epsilon) \cdot F_{\mathbf{LM}}^{\mathcal{A}_0} \rangle, \quad (3.10)$$

where I_S is the *soft operator* introduced in Ref. [1] and defined explicitly in Eq. (A.38).

Collinear limits require more attention. As stated above, we are interested in those limits that lead to singularities proportional to the matrix element of \mathcal{A}_0 , and we isolate such limits using the *projection operator* $\mathcal{P}_{\mathcal{A}_0}$. Terms that involve the soft operator $S_{\mathbf{m}}$ are unaffected by $\mathcal{P}_{\mathcal{A}_0}$, but some non-vanishing collinear limits are removed, e.g. the limit where the final state quark \mathbf{m}_q becomes collinear to the initial state gluon a_g ; thus

$$\mathcal{P}_{\mathcal{A}_0} \langle C_{am} \Delta^{(\mathbf{m})} F_{\mathbf{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \rangle = 0. \quad (3.11)$$

This procedure allows us to select a subset of limits of the NLO matrix element that do not change the Born level configuration and whose combination is free of $1/\epsilon$ singularities. This subset of limits does not remove *all* the singularities of the hard N(N)LO matrix element; in particular, those singularities whose limits include a different Born-level matrix element to that of \mathcal{A}_0 are not subtracted. We do this because our main aim is to study the structure of final-state collinear limits in the presence of quarks rather than provide complete formulas for the physical real correction to the cross section.

With these preliminary remarks out of the way, we can analyze collinear singularities related to the initial-state radiation. The only unresolved parton that can develop a collinear singularity with the incoming gluon a_g and still contribute to \mathcal{A}_0 is the final-state gluon \mathbf{m}_g . Then, following the steps detailed in Ref. [1], we obtain

$$\begin{aligned} \mathcal{P}_{\mathcal{A}_0} \langle \bar{S}_{\mathbf{m}} C_{am} \Delta^{(\mathbf{m})} F_{\mathbf{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle &= \langle \bar{S}_{\mathbf{m}} C_{am} \Delta^{(\mathbf{m})} F_{\mathbf{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \rangle \\ &= \frac{[\alpha_s]}{\epsilon} \left[\langle \Gamma_{a,g} F_{\mathbf{LM}}^{\mathcal{A}_0} \rangle + \langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\mathbf{LM}}^{\mathcal{A}_0} \rangle \right], \end{aligned} \quad (3.12)$$

where we have used the fact that $C_{am} \Delta^{(\mathbf{m})} = 1$, and introduced a particular notation for the left convolution

$$\mathcal{P}_{\alpha\beta}^{\text{gen}} \otimes F_{\mathbf{LM}} = \int_0^1 dz \mathcal{P}_{\alpha\beta}^{\text{gen}}(z) \frac{F_{\mathbf{LM}}[z \cdot p_a, p_b; \mathcal{H}_f]}{z}. \quad (3.13)$$

Definitions of the generalized initial-state anomalous dimensions $\Gamma_{a,fa}$ arising from the soft-collinear limits and of the (azimuthal averaged) collinear splitting functions $\mathcal{P}_{\alpha\beta}^{\text{gen}}$ are

¹We use dimensional regularization with $d = 4 - 2\epsilon$ throughout the paper.

reported in Eqs (A.17) and (A.18), respectively. The computation of the soft-subtracted collinear limit for the initial-state parton b_q is analogous. We find

$$\mathcal{P}_{A_0} \langle \bar{S}_m C_{bm} \Delta^{(m)} F_R^{\mathcal{A}_1}[\mathbf{m}] \rangle = \frac{[\alpha_s]}{\epsilon} \left[\langle \Gamma_{b,q} F_{LM}^{\mathcal{A}_0} \rangle + \langle F_{LM}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle \right], \quad (3.14)$$

where the right convolution is defined as

$$F_{LM} \otimes \mathcal{P}_{\alpha\beta}^{\text{gen}} = \int_0^1 dz \mathcal{P}_{\alpha\beta}^{\text{gen}}(z) \frac{F_{LM}[p_a, z \cdot p_b; \mathcal{H}_f]}{z}. \quad (3.15)$$

We continue with the discussion of the final-state collinear limits, which correspond to $i \in \mathcal{H}_f$ in Eq. (3.7). We note that, in this case, $C_{im} \Delta^{(m)} = E_i / (E_i + E_m) = z_{i,m}$, where $E_{i,m}$ are energies of the corresponding partons. The appearance of such factors makes the relation between the integrated collinear limits and conventional splitting functions and anomalous dimensions less transparent. This is a new complication with respect to the calculation of Ref. [1].

To address it, we will consider the two contributions to the function $F_R^{\mathcal{A}_1}[\mathbf{m}]$ in Eq. (3.5) separately. We start with $F_{LM}^{\mathcal{A}_1}[\mathbf{m}_g]$, and compute the hard-collinear limits for final-state partons

$$\sum_{i \in \mathcal{H}_f} \langle \bar{S}_m C_{im} \Delta^{(m)} F_{LM}^{\mathcal{A}_1}[\mathbf{m}_g] \rangle = \frac{[\alpha_s]}{\epsilon} \sum_{i \in \mathcal{H}_f} \langle \Gamma_{i,f_i \rightarrow f_i g} F_{LM}^{\mathcal{A}_0} \rangle, \quad (3.16)$$

where we have used the fact that clustering a collinear gluon with any parton does not change that parton's flavor. The *generalized final-state anomalous dimension* $\Gamma_{i,f_i \rightarrow f_i g}$ can be found in Eq. (A.19). We note that these quantities generalize the final-state anomalous dimensions $\gamma_{z,f_i \rightarrow f_i g}^{22}(L_i)$ defined in Eq. (A.20) and used in Ref. [1], and that the quantity $\Gamma_{i,g \rightarrow gg}$ appeared in this reference where it was denoted by $\Gamma_{i,g}$. Additionally, $\gamma_{z,f_i \rightarrow f_i g}^{22}(L_i)$ carries a subscript $z = z_{i,m}$ to indicate that it is obtained from an integral of a particular splitting function weighted by a factor $z_{i,m}$. We drop this subscript in the definition of $\Gamma_{i,f_i \rightarrow f_i g}$ to lighten the notation.

We emphasize that the generalized final-state anomalous dimensions differ from the conventional collinear anomalous dimensions that arise from integrals of the collinear splitting functions without additional factors $z_{i,m}$. To see how conventional collinear anomalous dimensions emerge, we need to consider the other contribution to the function $F_R^{\mathcal{A}_1}[\mathbf{m}]$, where a *quark* is potentially-unresolved. Since in $F_{LM}^{\mathcal{A}_1}[\mathbf{m}_q]$ the only resolved final-state partons are gluons, the unresolved quark \mathbf{m}_q has to be clustered with one of them. The clustered parton is a quark, so that the number of resolved gluons decreases by one and the number of resolved quarks increases by one. Working out the symmetry factors, we obtain

$$\sum_{i \in \mathcal{H}_f} \langle C_{im} \Delta^{(m)} F_{LM}^{\mathcal{A}_1}[\mathbf{m}_q] \rangle = \frac{[\alpha_s]}{\epsilon} \sum_{i \in \mathcal{H}_{f,q}} \langle \Gamma_{i,q \rightarrow gq} F_{LM}^{\mathcal{A}_0} \rangle, \quad (3.17)$$

where, on the right-hand side, $\mathcal{H}_{f,q}$ denotes the list of resolved final-state quarks. In the current case, this notation is redundant as this list consists of a single parton, but we note that Eq. (3.17) holds independently of the number of final-state quarks.

Combining Eqs (3.16) and (3.17), we obtain

$$\sum_{i \in \mathcal{H}_f} \langle \bar{S}_m C_{im} \Delta^{(m)} F_R^{\mathcal{A}_1}[\mathbf{m}] \rangle = \frac{[\alpha_s]}{\epsilon} \left\langle \left[\sum_{i \in \mathcal{H}_{f,g}} \Gamma_{i,g \rightarrow gg} + \sum_{i \in \mathcal{H}_{f,q}} \Gamma_{i,q} \right] F_{LM}^{\mathcal{A}_0} \right\rangle, \quad (3.18)$$

where we define

$$\Gamma_{i,q} = \Gamma_{i,q \rightarrow qq} + \Gamma_{i,q \rightarrow gq}. \quad (3.19)$$

We emphasize that the quantity $\Gamma_{i,q}$ in Eq. (3.19) is directly related to the conventional quark anomalous dimension, since the weight factors $z_{i,m}$ disappear when combining the two splittings. To show this, we make use of the fact that Γ is proportional to $\gamma^{22}(L_i)$ (cf. Eq. (A.19)) and write²

$$\begin{aligned} \gamma_{z,q \rightarrow qq}^{22}(L_i) + \gamma_{z,q \rightarrow gq}^{22} &= - \int_0^1 dz (\mathbb{1} - S_z) \frac{z P_{qq}(z) + z P_{qg}(z)}{[z(1-z)]^{2\epsilon}} + C_F \frac{1 - e^{-2\epsilon L_i}}{\epsilon} \\ &= - \int_0^1 dz (\mathbb{1} - S_z) \frac{P_{qq}(z)}{[z(1-z)]^{2\epsilon}} + C_F \frac{1 - e^{-2\epsilon L_i}}{\epsilon} \\ &= \gamma_{1,q \rightarrow qq}^{22}(L_i) = \gamma_q + 2\mathbf{T}_q^2 L_i + \mathcal{O}(\epsilon), \end{aligned} \quad (3.20)$$

where $\gamma_q = \frac{3}{2}C_F$ is the quark anomalous dimension, $L_i = \log(E_{\max}/E_i)$, and we have used the relation between splitting functions $P_{qg}(1-z) = P_{qg}(z)$ to remove the factors of z from the integrand.

Performing a similar calculation for $g^* \rightarrow gg$ splitting, we observe that the quantity $\Gamma_{i,g \rightarrow gg} = \frac{11}{6}C_A + 2C_A L_i + \mathcal{O}(\epsilon)$, i.e. to leading order in ϵ , gives exactly the C_A term of the gluon anomalous dimension, but the n_f term is missing. This is easily remedied by considering the process with an additional $q'\bar{q}'$ pair, which we referred to as \mathcal{A}_2 . Its cross section reads

$$2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_2} = \left\langle \frac{1}{(N_g - 1)!} F_{LM}^{gq}[\{g\}_{N_g-1}, \{q\}_1, \{q'\}_1, \{\bar{q}'\}_1] \right\rangle. \quad (3.21)$$

We can analyze this contribution following what we did for the process \mathcal{A}_1 . Since we are interested in contributions that contain singularities proportional to the matrix element for the process \mathcal{A}_0 , only some collinear limits of \mathcal{A}_2 need to be considered. In fact, the only way to arrive at \mathcal{A}_0 starting from \mathcal{A}_2 is to cluster q' and \bar{q}' into a gluon. For this to happen, either q' or \bar{q}' should be designated as potentially-unresolved, and then the only collinear limit we need to consider is $q' \parallel \bar{q}'$. Making use of the projection operator $\mathcal{P}_{\mathcal{A}_0}$ to select the relevant terms, we write

$$\mathcal{P}_{\mathcal{A}_0} [2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_2}] = \langle C_{\bar{q}'m_{q'}} \Delta^{(m)} F_{LM}^{\mathcal{A}_2}[\mathbf{m}_{q'}] \rangle + \langle C_{q'm_{\bar{q}'}} \Delta^{(m)} F_{LM}^{\mathcal{A}_2}[\mathbf{m}_{\bar{q}'}] \rangle, \quad (3.22)$$

where $F_{LM}^{\mathcal{A}_2}[\mathbf{m}_{q'}]$ and $F_{LM}^{\mathcal{A}_2}[\mathbf{m}_{\bar{q}'}]$ are defined analogously to Eq. (3.6), but using the list of partons in process \mathcal{A}_2 instead of \mathcal{A}_1 . The two terms on the right-hand side of Eq. (3.22) give identical contributions since the splitting $g \rightarrow q'\bar{q}'$ is symmetric under the swapping $q' \leftrightarrow \bar{q}'$. Furthermore, this splitting does not depend on the quark flavor. Thus, we obtain

²In the following equation, S_z is analogous to S_m but refers specifically to the variable z , i.e. $\lim_{z \rightarrow 1}$.

the contribution of $n_f - 1$ quark species from the process \mathcal{A}_2 , and add the contribution from the \mathcal{A}_3 process, where $q' = q$. This is equivalent to taking one term in the right-hand side of Eq. (3.22), computing its contribution, and multiplying the result by a factor of $2n_f$, obtaining

$$\sum_{n=2,3} \mathcal{P}_{\mathcal{A}_0} [2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_n}] = \frac{[\alpha_s]}{\epsilon} \sum_{i \in \mathcal{H}_{f,g}} \langle 2n_f \Gamma_{i,g \rightarrow q\bar{q}} F_{\text{LM}}^{\mathcal{A}_0} \rangle. \quad (3.23)$$

The anomalous dimension $\Gamma_{i,g \rightarrow q\bar{q}}$ is given in Eq. (A.19). Note that we have “distributed” the clustered gluon over the list of gluons in $F_{\text{LM}}^{\mathcal{A}_0}$ using its symmetry under gluon permutations.

Combining the contributions from Eqs (3.18) and (3.23), we find

$$\begin{aligned} & \sum_{i \in \mathcal{H}_f} \langle \bar{S}_m C_{im} \Delta^{(m)} F_R^{\mathcal{A}_1}[\mathbf{m}] \rangle + \sum_{n=2,3} \mathcal{P}_{\mathcal{A}_0} [2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_n}] \\ &= \frac{[\alpha_s]}{\epsilon} \left\langle \left[\sum_{i \in \mathcal{H}_{f,g}} \Gamma_{i,g} + \sum_{i \in \mathcal{H}_{f,q}} \Gamma_{i,q} \right] F_{\text{LM}}^{\mathcal{A}_0} \right\rangle, \end{aligned} \quad (3.24)$$

where, by analogy with Eq. (3.19), we introduced

$$\Gamma_{i,g} = \Gamma_{i,g \rightarrow gg} + 2n_f \Gamma_{i,g \rightarrow q\bar{q}}. \quad (3.25)$$

We note that, at $\mathcal{O}(\epsilon^0)$, $\Gamma_{i,g}$ corresponds to $\Gamma_{i,g} = \gamma_g + 2T_g^2 L_i + \mathcal{O}(\epsilon)$, where γ_g is the gluon anomalous dimension. To see this, one can repeat the manipulations performed in Eq. (3.20) for $\Gamma_{i,g \rightarrow q\bar{q}}$, using the fact that P_{gq} is symmetric under $z \leftrightarrow (1-z)$. As a result, the z factor in the integrand can be replaced by a factor $1/2$ (see the discussion in Sec. 5.1 in Ref. [1]).

We can now use Eq. (3.24) to extend the definition of the hard-collinear operator I_C , introduced in Ref. [1], to

$$I_C(\epsilon) = \sum_{i \in \mathcal{H}} \frac{\Gamma_{i,f_i}}{\epsilon}. \quad (3.26)$$

Therefore, the whole hard-collinear contribution can be written as

$$\begin{aligned} & \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} \langle \bar{S}_m C_{im} \Delta^{(m)} F_R^{\mathcal{A}_1}[\mathbf{m}] \rangle + \sum_{n=2,3} \mathcal{P}_{\mathcal{A}_0} [2s_{ab} d\hat{\sigma}_R^{\mathcal{A}_n}] \\ &= [\alpha_s] \langle I_C(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle + \frac{[\alpha_s]}{\epsilon} \left[\langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle \right]. \end{aligned} \quad (3.27)$$

The real-radiation contribution to the NLO cross section is obtained by combining the above expression with the soft contribution given in Eq. (3.10), and the fully-regulated \mathcal{O}_{NLO} term in Eq. (3.7).

To complete the computation, we need to account for the virtual corrections and perform the collinear renormalization of parton distribution functions. We use the operator I_V , introduced in Ref. [1], and defined in Eq. (A.36), to write the virtual corrections as

$$2s_{ab} d\hat{\sigma}_V^{\mathcal{A}_0} = [\alpha_s] \langle I_V(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle F_{\text{LV,fin}}^{\mathcal{A}_0} \rangle. \quad (3.28)$$

The function $F_{\text{LV,fin}}^{\mathcal{A}_0}$ in the above equation is analogous to F_{LM} in Eq. (2.2), but with $|\mathcal{M}_0|^2$ replaced by $2 \text{Re}[\langle \mathcal{M}_1^{\text{fin}} | \mathcal{M}_0 \rangle]$, where $\mathcal{M}_1^{\text{fin}}$ is the finite part of the one-loop virtual amplitude. The collinear renormalization contributions relevant for \mathcal{A}_0 , which come from the PDF redefinition, read

$$2s_{ab} d\hat{\sigma}_{\text{pdf}}^{\mathcal{A}_0} = \frac{\alpha_s(\mu)}{2\pi\epsilon} \left[\langle \hat{P}_{gg}^{(0)} \otimes F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle F_{\text{LM}}^{\mathcal{A}_0} \otimes \hat{P}_{qq}^{(0)} \rangle \right], \quad (3.29)$$

where the Altarelli-Parisi splitting functions $\hat{P}_{\alpha\beta}^{(0)}$ are defined in Eq. (A.13).

Using these results, we can finally write the NLO corrections to the process \mathcal{A}_0 . We combine Eqs (3.7), (3.10), (3.27), (3.28), and (3.29), and find an expression that does not contain ϵ poles

$$\begin{aligned} 2s_{ab} \left[d\hat{\sigma}_{\text{V}}^{\mathcal{A}_0} + \sum_{n=1}^3 \mathcal{P}_{\mathcal{A}_0} d\hat{\sigma}_{\text{R}}^{\mathcal{A}_n} + d\hat{\sigma}_{\text{pdf}}^{\mathcal{A}_0} \right] &= \mathcal{P}_{\mathcal{A}_0} \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \rangle + \langle F_{\text{LV,fin}}^{\mathcal{A}_0} \rangle \\ &+ [\alpha_s] \langle I_{\text{T}}^{(0)} \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle + [\alpha_s] \left[\langle \mathcal{P}_{gg}^{\text{NLO}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{NLO}} \rangle \right]. \end{aligned} \quad (3.30)$$

Here, $I_{\text{T}}^{(0)}$ is the $\mathcal{O}(\epsilon^0)$ coefficient in the expansion of the IR-finite operator

$$I_{\text{T}}(\epsilon) = I_{\text{V}}(\epsilon) + I_{\text{S}}(\epsilon) + I_{\text{C}}(\epsilon), \quad (3.31)$$

defined analogously to Ref. [1]. Its expression is reported in Eq. (A.45). Finally, for the boosted contribution, we used the relation $\alpha_s(\mu)/2\pi = \tilde{c}_\epsilon[\alpha_s]$, with $\tilde{c}_\epsilon = \Gamma(1-\epsilon)/e^{\epsilon\gamma_{\text{E}}}$, and the following relation between generalized splitting functions and Altarelli-Parisi splitting kernels

$$\mathcal{P}_{\alpha\beta}^{\text{gen}}(z, E_a) + \tilde{c}_\epsilon \hat{P}_{\alpha\beta}^{(0)}(z) = \epsilon \mathcal{P}_{\alpha\beta}^{\text{NLO}}(z, E_a) + \mathcal{O}(\epsilon^2), \quad (3.32)$$

valid for all α and β . The explicit expressions of functions $\mathcal{P}_{\alpha\beta}^{\text{NLO}}$ are reported in the ancillary file `FinalResult.m` provided with this paper, see Table 1.

Before concluding the NLO analysis, we briefly discuss the treatment of the singularities in \mathcal{A}_1 that *do not lead* to the partonic process \mathcal{A}_0 . Considering the function $F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q]$ for the sake of example, and taking the collinear limit $\mathbf{m}_q \parallel a_g$, we derive

$$(f_g \otimes f_q) \otimes \langle C_{am} \Delta^{(\text{m})} F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \rangle = \frac{[\alpha_s]}{\epsilon} (f_g \otimes f_q) \otimes \langle \mathcal{P}_{\bar{q}g}^{\text{gen}} \otimes F_{\text{LM}}^{\bar{q}q}[\{g\}_{N_g+1}] \rangle. \quad (3.33)$$

This singularity is cancelled by the collinear PDF renormalization in the Born process $\bar{q}q \rightarrow X + (N_g + 1)g$. We emphasize that the use of the projection operator $\mathcal{P}_{\mathcal{A}_0}$ allows us to discard such singularities, and focus on the main goal of this paper, which is an understanding of intertwined final-state collinear limits.

4 NNLO QCD corrections: general framework

In this section, we study the NNLO QCD corrections to the process \mathcal{A}_0 in Eq. (2.1). Similarly to the NLO case, we discard all singular contributions that lead to partonic

processes which differ from \mathcal{A}_0 . To compute the NNLO QCD corrections to \mathcal{A}_0 , we have to account for the two-loop virtual corrections to it, the one-loop corrections to processes \mathcal{A}_1 and \mathcal{A}_2 , and three double-real emission processes

$$\begin{aligned}\mathcal{A}_4 : a_g + b_q &\rightarrow X + (N_g + 2)g + q, \\ \mathcal{A}_5 : a_g + b_q &\rightarrow X + N_g g + q + q' \bar{q}', \\ \mathcal{A}_6 : a_g + b_q &\rightarrow X + (N_g - 2)g + q + q' \bar{q}' + q'' \bar{q}'',\end{aligned}\tag{4.1}$$

where we consider both the cases $q' \neq q$ and $q' = q$, and the same for q'' . In order to cancel the $1/\epsilon$ poles at NNLO, we write

$$d\hat{\sigma}_{\text{NNLO}}^{\mathcal{A}_0} = d\hat{\sigma}_{\text{VV}}^{\mathcal{A}_0} + \mathcal{P}_{\mathcal{A}_0} \left[d\hat{\sigma}_{\text{RV}}^{\mathcal{A}_1} + n_f d\hat{\sigma}_{\text{RV}}^{\mathcal{A}_2} + d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_4} + d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_5, \text{int}} + n_f d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_5} + n_f^2 d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_6} \right] + d\hat{\sigma}_{\text{pdf}}^{\mathcal{A}_0}.\tag{4.2}$$

We note that the only singular limits of the process \mathcal{A}_5 that will project onto the process \mathcal{A}_0 are those in which q' and \bar{q}' are either simultaneously soft, or are clustered together in the collinear regime (see the corresponding discussion after Eq. (3.22)). In either case, they produce a factor of n_f , which we write explicitly in Eq. (4.2). Similarly, the process \mathcal{A}_6 produces a factor n_f^2 when both q' and \bar{q}' as well as q'' and \bar{q}'' become collinear. There is also an interference term arising from process \mathcal{A}_5 if $q = q'$, which does not appear with a factor of n_f . This contribution is only singular in the triple-collinear limit and hence only produces simple poles in ϵ . No other singular limits need to be considered as these would result in processes different from \mathcal{A}_0 . In particular, we omit the discussion of IR divergences from flavor singlet configurations.

To compute the double-real contributions to Eq. (4.2), we need to analyze quantities of the following type

$$2s_{ab} d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_n} = \left\langle \frac{F_{\text{LM}}^{qq}[\{g\}_{N_g}, \{q\}_{N_q}, \{\bar{q}\}_{N_{\bar{q}}}, \dots]}{N_g! \times N_q! \times N_{\bar{q}}! \times \dots} \right\rangle, \quad n \in \{4, 5, 6\},\tag{4.3}$$

where integer numbers $N_g, N_q, N_{\bar{q}}$, etc., indicate lengths of the relevant lists; they depend on the process \mathcal{A}_n under consideration. The first step consists of employing damping factors to choose the potentially-unresolved partons in F_{LM} . Following Ref. [1], we will refer to such partons as \mathbf{m} and \mathbf{n} , and to the relevant damping factor as $\Delta^{(\mathbf{mn})}$.³ Writing the right-hand side in Eq. (4.3) as the sum of different contributions, each depending on a specific pair (\mathbf{mn}) of potentially unresolved partons, we obtain

$$2s_{ab} d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_n} = \sum_{(\mathbf{mn})} \langle \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad n \in \{4, 5, 6\}.\tag{4.4}$$

In the above equation, the sum runs over all unordered (\mathbf{mn}) pairs, except in the case of a $q_i \bar{q}_i$ pair, where we consider both the pair $(q_i \bar{q}_i)$ and $(\bar{q}_i q_i)$. The function $\Theta_{\mathbf{mn}}$ will be specified shortly. For each \mathcal{A}_n , labels \mathbf{m} and \mathbf{n} may refer to partons of different types, and one has to account for all the possible (\mathbf{mn}) pairings compatible with \mathcal{A}_n . In particular, for process \mathcal{A}_4 , (\mathbf{mn}) can be a pair of gluons (gg) or a quark-gluon pair ($q_i g$); for \mathcal{A}_5 , (\mathbf{mn}) can

³The construction of damping factors is discussed in detail in Appendix B of Ref. [1].

be a (anti)quark-gluon pair $(q_i g)$ or $(\bar{q}_i g)$, or a quark-antiquark pair of the same $(q_i \bar{q}_i)$ or different flavor $(q_i \bar{q}_j)$; and for \mathcal{A}_6 , (\mathbf{mn}) must be a fermion-fermion pair $(q_i q_j)$, $(\bar{q}_i \bar{q}_j)$, and $(q_i \bar{q}_j)$.

For concreteness, we explicitly write the function $F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}]$ required for the process \mathcal{A}_4 in Eq. (4.4). The expression reads

$$2s_{ab} d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_4} = \langle \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} (F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g]) \rangle. \quad (4.5)$$

We note that symmetry factors depend on the list of resolved final-state partons; they are included in the definition of $F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}]$

$$\begin{aligned} F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] &= \frac{1}{N_g!} F_{\text{LM}}^{gq}[\{g\}_{N_g}, \{q\}_1 | \mathbf{m}_g, \mathbf{n}_g], \\ F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] &= \frac{1}{(N_g + 1)!} F_{\text{LM}}^{gq}[\{g\}_{N_g+1} | \mathbf{m}_q, \mathbf{n}_g]. \end{aligned} \quad (4.6)$$

There are important differences in the singular limits of the F_{LM} functions that appear in Eq. (4.4), depending on which partons are resolved and which are unresolved. If the unresolved partons (\mathbf{mn}) are either two gluons or a $q_i \bar{q}_i$ pair, then the double-soft limit is singular, whereas if they are qg , $\bar{q}g$, $q_i q_k$ etc., it is not. We will refer to the former combination as DS and the latter as \mathcal{DS} , so that

$$\text{DS} = \{(gg), (q_i \bar{q}_i), (\bar{q}_i q_i)\}, \quad (4.7)$$

and

$$\mathcal{DS} = \{(q_i \bar{q}_j), (q_i q_k), (\bar{q}_i \bar{q}_k), (q_i g), (\bar{q}_i g), \text{ with } i \neq j\}. \quad (4.8)$$

Then, if $(\mathbf{mn}) \in \text{DS}$, we introduce an energy ordering by identifying $\Theta_{\mathbf{mn}}$ with the Heaviside function $\Theta_{\mathbf{mn}} = \Theta(E_{\mathbf{m}} - E_{\mathbf{n}})$, where $E_{\mathbf{m}, \mathbf{n}}$ are the energies of partons \mathbf{m}, \mathbf{n} . Note that the presence of the energy ordering is the reason why we need to consider both $(q_i \bar{q}_i)$ and $(\bar{q}_i q_i)$ pairs. If, on the other hand, $(\mathbf{mn}) \in \mathcal{DS}$, we do not introduce the energy ordering and take $\Theta_{\mathbf{mn}} = 1$. Thus, the proper reading of Eq. (4.5) requires associating $\Theta_{\mathbf{mn}}$ with the energy-ordering Heaviside function in the first term in brackets and with 1 in the second term. Furthermore, we extend the convention of Eq. (3.9) so that $S_{\mathbf{mn}} = 0$ for $(\mathbf{mn}) \in \mathcal{DS}$.

Another important difference between various unresolved partons concerns single-soft singular limits, which only exist if at least one parton in the list (\mathbf{mn}) is a gluon. To retain generality of the calculations, we keep the convention in Eq. (3.9) that the single-soft operator makes an expression vanish if it is applied to a quark or an antiquark. Moreover, if the unresolved partons are a quark and a gluon, we always label them as $(\mathbf{m}_q \mathbf{n}_g)$, see Eq. (4.6).

Finally, since we are interested in unresolved contributions that lead to the process \mathcal{A}_0 , we need to account for different collinear limits depending on the list of unresolved partons. For example, in case of the $q_i \bar{q}_i$ final state, we exclude cases where q_i and \bar{q}_i are collinear to different hard partons and only allow limits where they are collinear to each other, as discussed after Eq. (4.2). In addition, we note that (sequential) double-collinear limits may not be invariant under the exchange $\mathbf{m} \leftrightarrow \mathbf{n}$ for a generic unresolved final state

(\mathbf{mn}). This possibility was pointed out in Ref. [1], although there (for the gg unresolved final state) these limits did in fact commute.

In spite of the differences between the various cases, the computational strategy is always the same and follows closely the discussion in Ref. [1]. In particular, as explained in Ref. [1], when computing the different contributions to the cross section in Eq. (4.2), it is beneficial to combine terms that have *equal* number of *resolved* partons. In the double-virtual contribution, the number of resolved partons is the same as in the Born process, i.e. N , in the real-virtual contribution this number can be $N + 1$ or N , and in the double-real contribution it can be $N + 2$, $N + 1$ or N .

We proceed with the discussion of the double-real contributions, starting from Eq. (4.4), and aiming at identifying the singular limits that result in a well-defined number of resolved partons. We begin by regularizing the double-soft and single-soft divergences, writing

$$2s_{ab} d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_n} = \sum_{(\mathbf{mn})} \left[\langle S_{\mathbf{mn}} \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle + \langle \bar{S}_{\mathbf{mn}} S_{\mathbf{n}} \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle + \langle \bar{S}_{\mathbf{mn}} \bar{S}_{\mathbf{n}} \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \right], \quad (4.9)$$

with $n \in \{4, 5, 6\}$, and where $\bar{S}_{\mathbf{mn}} = 1 - S_{\mathbf{mn}}$. The above formula applies to all unresolved partons (\mathbf{mn}) and all processes \mathcal{A}_n due to the definitions of the double-soft operator $S_{\mathbf{mn}}$ and the single-soft operator $S_{\mathbf{m}}$, explained above. The second and the third term on the right-hand side of Eq. (4.9) still contain collinear divergences that must be regularized. The latter requires particular care since the single- and triple-collinear limits overlap. To overcome this, the nested soft-collinear scheme employs phase-space partitions and sectors [13, 14, 17]. However, these procedures lead to repeated computations of similar integrals, which result in an unwieldy number of counterterms and obscures physical structures. Therefore, it is convenient to recombine the various limits from the different sectors, and to do so *prior to* evaluating the effect of each limit on the F_{LM} quantities and integrating over the unresolved phase space.

The steps required to do so are somewhat tedious and are outlined in Sec. 4 of Ref. [1]. Extending them to make them applicable to a generic unresolved final state, we write Eq. (4.9) as the sum of three contributions,

$$2s_{ab} d\hat{\sigma}_{\text{RR}}^{\mathcal{A}_n} = \sum_{(\mathbf{mn})} \left[\Sigma_{\text{FR}}^{\mathcal{A}_n}[(\mathbf{mn})]_{\text{RR}} + \Sigma_{\text{SU}}^{\mathcal{A}_n}[(\mathbf{mn})]_{\text{RR}} + \Sigma_{\text{DU}}^{\mathcal{A}_n}[(\mathbf{mn})]_{\text{RR}} \right], \quad n \in \{4, 5, 6\}. \quad (4.10)$$

The three terms on the right-hand side of the above equation correspond to the double-real fully-resolved, single-unresolved, and double-unresolved contributions, respectively. The fully-resolved term reads

$$\Sigma_{\text{FR}}^{\mathcal{A}_n}[(\mathbf{mn})]_{\text{RR}} = \langle \bar{S}_{\mathbf{mn}} \bar{S}_{\mathbf{n}} \Omega_1 \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad (4.11)$$

where the collinear regularization operator Ω_1 is defined in Eq. (D.5) of Ref. [1]. The double-unresolved and single-unresolved contributions in Eq. (4.10) have slightly different expressions depending on whether $(\mathbf{mn}) \in \text{DS}$ or $(\mathbf{mn}) \in \mathcal{DS}$. For this reason, we present them separately.

Beginning with the double-unresolved terms, if $(\mathbf{mn}) \in \text{DS}$, we write⁴

$$\begin{aligned}
& \Sigma_{\text{DU}}^{\mathcal{A}_n}[(\mathbf{mn}) \in \text{DS}]_{\text{RR}} \\
&= \langle S_{\mathbf{mn}} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle + \sum_{i \in \mathcal{H}} \langle \bar{S}_{\mathbf{m}} C_{i\mathbf{m}} \Delta^{(\mathbf{m})} \langle S_{\mathbf{n}} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle \\
&+ \sum_{i \in \mathcal{H}} \langle S_{\mathbf{n}} \langle \bar{S}_{\mathbf{m}} C_{i\mathbf{m}} \Delta^{(\mathbf{mn})} \Theta_{\mathbf{nm}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle + \frac{1}{2} \sum_{i,j \in \mathcal{H}} \langle \bar{S}_{\mathbf{n}} \bar{S}_{\mathbf{m}} C_{j\mathbf{n}} C_{i\mathbf{m}} \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \quad (4.12) \\
&+ \sum_{i \in \mathcal{H}} \frac{N_{\mathbf{m}||\mathbf{n}}(\epsilon)}{2} \langle \bar{S}_{\mathbf{m}} C_{i\mathbf{m}} \Delta^{(\mathbf{m})} \sigma_{i\mathbf{m}}^{-\epsilon} \langle (\mathbb{1} - 2\Theta_{\mathbf{mn}} S_{\mathbf{n}}) C_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle \\
&+ [\alpha_s]^2 2^{1+2\epsilon} \delta_{\mathbf{m}}(\epsilon) \left(\frac{2E_{\text{max}}}{\mu} \right)^{-2\epsilon} \left[-\langle I_S(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle + \frac{(2E_{\text{max}}/\mu)^{-2\epsilon}}{2\epsilon^2} N_c(\epsilon) \sum_{i \in \mathcal{H}} \mathbf{T}_i^2 \langle F_{\text{LM}}^{\mathcal{A}_0} \rangle \right] \\
&+ \Sigma_{\text{DU}}^{\mathcal{A}_n, \text{rest}}[(\mathbf{mn}) \in \text{DS}]_{\text{RR}},
\end{aligned}$$

where the terms collectively denoted as $\Sigma_{\text{DU}}^{\mathcal{A}_n, \text{rest}}[(\mathbf{mn}) \in \text{DS}]_{\text{RR}}$ in the above equation can be found in Eqs (A.49 – A.57); they contain single $1/\epsilon$ poles at most. The quantities $N_{\mathbf{m}||\mathbf{n}}$ and $N_c(\epsilon)$ are defined in Eq. (A.7), the quantities $\delta_{\mathbf{m}}(\epsilon)$ in Eq. (A.22), and $\sigma_{ij} = \eta_{ij}/(1 - \eta_{ij})$, where $\eta_{ij} = (1 - \cos \theta_{ij})/2$. If $(\mathbf{mn}) \in \mathcal{DS}$, the double-unresolved contribution can be written as

$$\begin{aligned}
& \Sigma_{\text{DU}}^{\mathcal{A}_n}[(\mathbf{mn}) \in \mathcal{DS}]_{\text{RR}} \\
&= \sum_{i \in \mathcal{H}} \langle C_{i\mathbf{m}} \Delta^{(\mathbf{m})} \langle S_{\mathbf{n}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle + \frac{1}{2} \sum_{i,j \in \mathcal{H}} \langle \bar{S}_{\mathbf{n}} (C_{j\mathbf{n}} C_{i\mathbf{m}} + C_{i\mathbf{m}} C_{j\mathbf{n}}) \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \quad (4.13) \\
&+ \sum_{i \in \mathcal{H}} N_{\mathbf{m}||\mathbf{n}}(\epsilon) \langle C_{i\mathbf{m}} \sigma_{i\mathbf{m}}^{-\epsilon} \Delta^{(\mathbf{m})} \langle \bar{S}_{\mathbf{n}} C_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle + \Sigma_{\text{DU}}^{\mathcal{A}_n, \text{rest}}[(\mathbf{mn}) \in \mathcal{DS}]_{\text{RR}}.
\end{aligned}$$

Again, terms denoted as $\Sigma_{\text{DU}}^{\mathcal{A}_n, \text{rest}}[(\mathbf{mn}) \in \mathcal{DS}]_{\text{RR}}$ in the above equation have at most $1/\epsilon$ poles; they can be found in Eqs (A.59 – A.63). We emphasize once more that the derivation of the above formulas, as well as the notation that we employ, follows the discussion in Ref. [1] with minor modifications which account for the more general nature of the final state (\mathbf{mn}) .

Moving on to single-unresolved terms, if $(\mathbf{mn}) \in \text{DS}$, we write

$$\begin{aligned}
& \Sigma_{\text{SU}}^{\mathcal{A}_n}[(\mathbf{mn}) \in \text{DS}]_{\text{RR}} \\
&= \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \Delta^{(\mathbf{m})} \langle S_{\mathbf{n}} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle + \sum_{i \in \mathcal{H}} \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} (\mathbb{1} - S_{\mathbf{n}} \Theta_{\mathbf{mn}}) C_{i\mathbf{n}} \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \\
&+ \frac{1}{2} \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \Delta^{(\mathbf{m})} \langle (\mathbb{1} - 2\Theta_{\mathbf{mn}} S_{\mathbf{n}}) C_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle \\
&+ \sum_{i \in \mathcal{H}} \left\langle \mathcal{O}_{\text{NLO}}^{(i, \mathbf{m})} \omega_{i||\mathbf{n}}^{\mathbf{m}i, \mathbf{n}i} [(\eta_{i\mathbf{m}}/2)^{-\epsilon} - 1] (\mathbb{1} - S_{\mathbf{n}} \Theta_{\mathbf{mn}}) C_{i\mathbf{n}} \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \right\rangle \quad (4.14)
\end{aligned}$$

⁴With nested angular brackets we indicate that the operations appearing within the innermost brackets have to be performed first, and then we integrate over the corresponding unresolved degrees of freedom. This result is then acted upon by the outer operator.

$$\begin{aligned}
& + \sum_{i \in \mathcal{H}} \frac{1}{2} \left\langle \mathcal{O}_{\text{NLO}}^{(i, \mathbf{m})} \omega_{\mathbf{m} \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i} \Delta^{(\mathbf{m})} [N_{\mathbf{m} \parallel \mathbf{n}}(\epsilon) \sigma_{i\mathbf{m}}^{-\epsilon} - 1] \langle (\mathbb{1} - 2\Theta_{\mathbf{mn}} S_{\mathbf{n}}) C_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \right\rangle \\
& + \sum_{i \in \mathcal{H}} [\alpha_s] \frac{N_\epsilon^{(b, d)}}{2} \gamma_{\perp, g \rightarrow \mathbf{mn}}^{22} \langle \mathcal{O}_{\text{NLO}}^{(i, [\mathbf{mn}])} \omega_{\mathbf{m} \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i} \sigma_{i[\mathbf{mn}]}^{-\epsilon} (E_{[\mathbf{mn}]} / \mu)^{-2\epsilon} (r_i^\mu r_i^\nu + g^{\mu\nu}) \Delta^{([\mathbf{mn}])} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{mn}] \rangle \\
& + \sum_{i \in \mathcal{H}} [\alpha_s] \frac{N_\epsilon^{(b, d)}}{2} \gamma_{\perp, g \rightarrow \mathbf{mn}}^{22, r} \langle \mathcal{O}_{\text{NLO}}^{(i, [\mathbf{mn}])} \omega_{\mathbf{m} \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i} \sigma_{i[\mathbf{mn}]}^{-\epsilon} (E_{[\mathbf{mn}]} / \mu)^{-2\epsilon} \Delta^{([\mathbf{mn}])} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{mn}] \rangle,
\end{aligned}$$

whereas if $(\mathbf{mn}) \in \mathcal{DS}$, the expression reads

$$\begin{aligned}
& \Sigma_{\text{SU}}^{\mathcal{A}_n}[(\mathbf{mn}) \in \mathcal{DS}]_{\text{RR}} \\
& = \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \Delta^{(\mathbf{m})} \langle S_{\mathbf{n}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle + \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \Delta^{(\mathbf{m})} \langle C_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \rangle \\
& + \sum_{i \in \mathcal{H}} \left\langle \mathcal{O}_{\text{NLO}}^{(i, \mathbf{m})} \omega_{\mathbf{m} \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i} [N_{\mathbf{m} \parallel \mathbf{n}}(\epsilon) \sigma_{i\mathbf{m}}^{-\epsilon} - 1] \Delta^{(\mathbf{m})} \langle C_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \right\rangle \\
& + \sum_{i \in \mathcal{H}} \left\langle \left[\mathcal{O}_{\text{NLO}}^{(\mathbf{n})} C_{i\mathbf{m}} + \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \bar{S}_{\mathbf{n}} C_{i\mathbf{n}} \right] \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \right\rangle \tag{4.15} \\
& + \sum_{i \in \mathcal{H}} \left\langle \left[\mathcal{O}_{\text{NLO}}^{(i, \mathbf{n})} \omega_{i \parallel \mathbf{m}}^{\mathbf{m}i, \mathbf{n}i} [(\eta_{i\mathbf{n}}/2)^{-\epsilon} - 1] C_{i\mathbf{m}} + \mathcal{O}_{\text{NLO}}^{(i, \mathbf{m})} \omega_{i \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i} [(\eta_{i\mathbf{m}}/2)^{-\epsilon} - 1] \bar{S}_{\mathbf{n}} C_{i\mathbf{n}} \right] \right. \\
& \left. \times \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \right\rangle.
\end{aligned}$$

In Eqs (4.14) and (4.15), we have used the notation

$$\mathcal{O}_{\text{NLO}}^{(i, x)} = \bar{S}_x \bar{C}_{ix}, \quad \mathcal{O}_{\text{NLO}}^{(x)} = \sum_{i \in \mathcal{H}} \mathcal{O}_{\text{NLO}}^{(i, x)} \omega^{xi}, \tag{4.16}$$

where ω^{xi} are NLO partition functions. The functions $\omega_{\mathbf{m} \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i}$, $\omega_{i \parallel \mathbf{m}}^{\mathbf{m}i, \mathbf{n}i}$, $\omega_{i \parallel \mathbf{n}}^{\mathbf{m}i, \mathbf{n}i}$ originate from the NNLO partition functions $\omega^{\mathbf{m}i, \mathbf{n}i}$, evaluated in the limits $\mathbf{m} \parallel \mathbf{n}$, $i \parallel \mathbf{m}$, and $i \parallel \mathbf{n}$, respectively. The anomalous dimensions $\gamma_{\perp, g \rightarrow \mathbf{mn}}^{22}$ and $\gamma_{\perp, g \rightarrow \mathbf{mn}}^{22, r}$ are defined in Eq. (A.25), and the vector r_i^μ is described in Appendix E of Ref. [1].

Although we do not discuss the derivation of the above formulas for the double-unresolved and single-unresolved cases, we believe it is useful to present them to emphasize their proximity to similar formulas presented in Ref. [1] for the (gg) unresolved final state. It is this similarity and the appearance of universal structures that will be key for deriving formulas for integrated NNLO subtraction terms for processes with an *arbitrary number of jets* at hadron colliders, a problem that we would like to investigate in the future.

We continue with the discussion of the real-virtual and double-virtual corrections in Eq. (4.2), starting from the former. They are analyzed in the same way as the real-emission contribution at NLO, see Sec. 3. A generic real-virtual contribution reads

$$2s_{ab} d\hat{\sigma}_{\text{RV}}^{\mathcal{A}_n} = \left\langle \frac{F_{\text{RV}}^{gg}[\{g\}_{N_g}, \{q\}_{N_q}, \{\bar{q}\}_{N_{\bar{q}}}, \dots]}{N_g! \times N_q! \times N_{\bar{q}}! \times \dots} \right\rangle, \quad n \in \{1, 2, 3\}, \tag{4.17}$$

where F_{RV} is defined similarly to F_{LM} in Eq. (2.2). We use damping factors $\Delta^{(\mathbf{m})}$ and write

$$2s_{ab} d\hat{\sigma}_{\text{RV}}^{\mathcal{A}_n} = \sum_{\mathbf{m}} \langle \Delta^{(\mathbf{m})} F_{\text{RV}}^{\mathcal{A}_n}[\mathbf{m}] \rangle, \tag{4.18}$$

where each $F_{\text{RV}}^{\mathcal{A}_n}[\mathbf{m}]$ function depends on a particular potentially-unresolved parton \mathbf{m} . We extract soft and collinear singularities following steps discussed in the NLO calculation, and obtain

$$2s_{ab} d\hat{\sigma}_{\text{RV}}^{\mathcal{A}_n} = \sum_{\mathbf{m}} \left[\Sigma_{\text{SU}}^{\mathcal{A}_n}[\mathbf{m}]_{\text{RV}} + \Sigma_{\text{DU}}^{\mathcal{A}_n}[\mathbf{m}]_{\text{RV}} \right], \quad n \in \{1, 2, 3\}, \quad (4.19)$$

where

$$\begin{aligned} \Sigma_{\text{SU}}^{\mathcal{A}_n}[\mathbf{m}]_{\text{RV}} &= \langle \mathcal{O}_{\text{NLO}}^{(\mathbf{m})} \Delta^{(\mathbf{m})} F_{\text{RV}}^{\mathcal{A}_n}[\mathbf{m}] \rangle, \\ \Sigma_{\text{DU}}^{\mathcal{A}_n}[\mathbf{m}]_{\text{RV}} &= \langle S_{\mathbf{m}} F_{\text{RV}}^{\mathcal{A}_n}[\mathbf{m}] \rangle + \sum_{i \in \mathcal{H}} \langle \bar{S}_{\mathbf{m}} C_{i\mathbf{m}} \Delta^{(\mathbf{m})} F_{\text{RV}}^{\mathcal{A}_n}[\mathbf{m}] \rangle. \end{aligned} \quad (4.20)$$

We reiterate that, if parton \mathbf{m} is a quark or an antiquark, we drop the corresponding soft limit $S_{\mathbf{m}}$ from Eq. (4.20) and replace $\bar{S}_{\mathbf{m}}$ with the identity operator.

Finally, we consider the double-virtual corrections. Their IR singularities are universal [87–89], and the corrections to the Born process \mathcal{A}_0 are simply given by

$$2s_{ab} d\hat{\sigma}_{\mathcal{A}_0}^{\text{VV}} = \langle F_{\text{VV}}^{\mathcal{A}_0} \rangle \equiv \Sigma_{\text{DU}}^{\mathcal{A}_0}|_{\text{VV}}. \quad (4.21)$$

The function F_{VV} is defined in a way that is similar to F_{LM} in Eq. (2.2). The explicit expression for Eq. (4.21) can be found in Eq. (4.86) in Ref. [1].

At this point, we can combine the single-unresolved and double-unresolved terms in Eqs (4.19) and (4.21), originating from the real-virtual and double-virtual corrections, with the corresponding single-unresolved and double-unresolved contributions from the double-real corrections in Eq. (4.10). Together, they should provide a physically-transparent description of $d\hat{\sigma}_{\text{NNLO}}^{\mathcal{A}_0}$ in Eq. (4.2). In the remaining sections of this paper, we discuss how all these different contributions can be combined and simplified, demonstrate the cancellation of the infrared poles, and derive compact results for finite remainders.

5 Simplifications of the double-unresolved contributions

The formulas presented in the previous section express the contributions to the NNLO corrections in terms of soft and collinear operators acting on the functions F_{LM} , after the limits from different partitions and sectors have been combined. The action of these operators on the F_{LM} quantities leads to universal structures (eikonal functions, collinear splitting functions) and reduced matrix elements that do not depend on the momenta of the unresolved partons. Integrating these universal functions, we obtain $1/\epsilon$ poles that have to disappear when double-virtual, real-virtual, and double-real contributions as well as the PDF renormalization are combined. In this section, we simplify the formulas obtained in the previous sections, focusing on the double-unresolved contributions, which contain the strongest singularities. This preliminary work will enable us to dramatically simplify the demonstration of the cancellation of singularities, which we discuss in Sec. 6.

In our analysis below, we follow Ref. [1], where pure gluonic final states were considered. The terms involving virtual contributions and/or soft limits require only minor modifications, and we describe these in Sec. 5.1. However, as already seen in Sec. 3, the

situation is more complex when considering the collinear limits of final states consisting of quarks and gluons, because certain combinations of such limits have to work in unison to produce such physical quantities as splitting functions and collinear anomalous dimensions. We discuss this in Sec. 5.2. Understanding how this happens at NNLO is an important step required to extend the analysis of Ref. [1] to generic partonic processes.

5.1 Simplifying virtual and soft corrections

In this section, we discuss the soft and virtual contributions. As stated above, these are very similar to those discussed in Ref. [1]. There, the treatment of the singular triple-color correlated terms $\sim f_{abc}T_i^a T_j^b T_k^c$ was performed using generic representations of the color charges, and thus can be applied verbatim to the present case. We therefore focus on contributions containing products of the color-charge operators $\mathbf{T}_i \cdot \mathbf{T}_j$ or $\{\mathbf{T}_i \cdot \mathbf{T}_j, \mathbf{T}_k \cdot \mathbf{T}_l\}$. These appear in the double-virtual corrections, in the soft and collinear limits of the real-virtual corrections, as well as in the double-soft limit and a combination of the single-soft and hard-collinear limits in the double-real terms.

The expressions for the double-virtual and the soft limit of the real-virtual contributions can be borrowed from Eqs (4.86, 4.102) in Ref. [1] with obvious replacements of F_{LM} and F_{VV} functions. This is possible because only an unresolved gluon contributes to the soft limit in the real-virtual case and because the double-virtual contributions are described by universal formulas that only depend on the momenta of hard partons, their color charges, and their collinear anomalous dimensions. The double-soft limit can again be borrowed from Ref. [1], but we have to supplement that result with the double-soft contribution from the unresolved $q'\bar{q}'$ pair, which was calculated for the nested subtraction scheme in Ref. [90]. These additional contributions are proportional to n_f , so it is easy to identify and track them. The full result reads

$$\begin{aligned} & \langle S_{\text{mn}} \Theta_{\text{mn}} (F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q')}, \mathbf{n}_{(\bar{q}')}) \rangle \\ &= [\alpha_s]^2 \left\langle \left[\frac{1}{2} I_{\text{S}}^2(\epsilon) + \left(\frac{C_{\text{A}}}{\epsilon^2} c_1(\epsilon) + \frac{\beta_0}{\epsilon} c_2(\epsilon) + \beta_0 c_3(\epsilon) - \frac{2}{3} n_f T_{\text{R}} c_4(\epsilon) \right) \tilde{I}_{\text{S}}(2\epsilon) \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \quad (5.1) \\ &+ [\alpha_s]^2 \sum_{(ij)} \langle [(S_{gg,T^2}^{\text{fin}})_{ij} + n_f (S_{q\bar{q},T^2}^{\text{fin}})_{ij}] (\mathbf{T}_i \cdot \mathbf{T}_j) \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle, \end{aligned}$$

where we have introduced the shorthand notation

$$F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q')}, \mathbf{n}_{(\bar{q}')}] = F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{q'}, \mathbf{n}_{\bar{q}'}] + F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{\bar{q}'}, \mathbf{n}_{q'}], \quad (5.2)$$

to denote the $q'\bar{q}'$ final state. The constant factors $c_{1,2,3}$ in Eq. (5.1) have already been presented in Ref. [1] and are reported in Eq. (A.9). The new coefficient

$$c_4(\epsilon) = -\frac{13}{6} + \left(\frac{125}{18} - \frac{35}{3} \log 2 - 12 \log^2 2 \right) \epsilon, \quad (5.3)$$

has been extracted from Ref. [90]. Finally, the last line of Eq. (5.1) contains finite remainders arising from the double soft configurations, which are extracted from Ref. [90]. The explicit expression of $(S_{gg,T^2}^{\text{fin}})_{ij}$ and $(S_{q\bar{q},T^2}^{\text{fin}})_{ij}$ are given in the ancillary file `FinalResult.m`

provided with this paper (cf. Table 1). Furthermore, β_0 in Eq. (5.1) stands for the leading-order QCD β -function. In contrast, in Ref. [1], β_0 was always assumed to be evaluated with $n_f = 0$ since only final states with gluons were accounted for in that reference.

Color-correlated terms in the double-unresolved contributions also arise when soft-limit operators or virtual corrections appear together with the collinear operators. The second term on the right-hand side of $\Sigma_{\text{DU}}^{\mathcal{A}_n}[\mathbf{m}]_{\text{RV}}$ in Eq. (4.20) is the first example of the latter; we will refer to it as the hard-collinear limit of the real-virtual contribution. Similarly to the NLO case, we are only interested in contributions that are relevant for cancelling IR divergences proportional to the matrix element squared for \mathcal{A}_0 ; this restricts the number of collinear limits that we have to consider. Indeed, for the initial-state collinear limits, we only need to account for the splitting $a_g \rightarrow \mathbf{m}_g g^*$ and $b_q \rightarrow \mathbf{m}_g q^*$. For the final-state collinear limits, we need to consider the $g^* \rightarrow \mathbf{m}_g i_g$, $q^* \rightarrow \mathbf{m}_g i_q$ and $q^* \rightarrow \mathbf{m}_q i_q$ splittings, as well as the branching $g^* \rightarrow \mathbf{m}_{q'} i_{q'}$ and $g^* \rightarrow \mathbf{m}_{\bar{q}'} i_{q'}$. The latter splittings are relevant for the process \mathcal{A}_2 , which we will collectively indicate as

$$F_{\text{RV}}^{\mathcal{A}_2}[\mathbf{m}_{(q'\bar{q}')}] = F_{\text{RV}}^{\mathcal{A}_2}[\mathbf{m}_{q'}] + F_{\text{RV}}^{\mathcal{A}_2}[\mathbf{m}_{\bar{q}'}]. \quad (5.4)$$

In full analogy with Ref. [1], we write

$$\begin{aligned} & \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} \left\langle \bar{S}_{\mathbf{m}} C_{im} \Delta^{(\mathbf{m})} \left(F_{\text{RV}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{RV}}^{\mathcal{A}_1}[\mathbf{m}_q] + n_f F_{\text{RV}}^{\mathcal{A}_2}[\mathbf{m}_{(q'\bar{q}')}] \right) \right\rangle \\ &= [\alpha_s]^2 \left[\left\langle I_C(\epsilon) I_V(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle + \frac{1}{\epsilon} \left\langle \mathcal{P}_{gg}^{\text{gen}} \otimes [I_V(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0}] + [I_V(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0}] \otimes \mathcal{P}_{qq}^{\text{gen}} \right\rangle \right] \\ &+ [\alpha_s] \left[\left\langle I_C(\epsilon) \cdot F_{\text{LV,fin}}^{\mathcal{A}_0} \right\rangle + \frac{1}{\epsilon} \left\langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LV,fin}}^{\mathcal{A}_0} + F_{\text{LV,fin}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \right\rangle \right] \\ &- [\alpha_s]^2 \frac{\Gamma(1-\epsilon) \beta_0}{e^{\epsilon \gamma_E}} \frac{1}{\epsilon} \left[\left\langle I_C(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle + \frac{1}{\epsilon} \left\langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} + F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \right\rangle \right] \\ &- \frac{[\alpha_s]^2}{\epsilon^2} h_c(\epsilon) \left\langle \left[C_A \tilde{I}_C(2\epsilon) + 2(C_F - C_A) \tilde{I}_{C,\text{nf}}(2\epsilon) \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\ &- \frac{[\alpha_s]^2}{2\epsilon^3} C_A h_c(\epsilon) \left\langle \mathcal{P}_{gg}^{1\text{L,gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} + F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{1\text{L,gen}} \right\rangle, \end{aligned} \quad (5.5)$$

where $h_c(\epsilon)$ is reported in Eq. (A.10), and $\mathcal{P}_{\alpha\alpha}^{1\text{L,gen}}$ are the one-loop splitting functions defined in Eq. (A.31). The collinear operator $\tilde{I}_C(2\epsilon)$ is the generalization of the operator I_C to the real-virtual case. It is defined as follows

$$\tilde{I}_C(2\epsilon) = \sum_{i \in \mathcal{H}} \frac{\Gamma_{i,f_i}^{1\text{L}}(\epsilon)}{2\epsilon}, \quad (5.6)$$

where $\Gamma_{i,f_i}^{1\text{L}}$ represents the generalized initial- and final-state one-loop anomalous dimension, given in Eqs (A.30), (A.33) and (A.35). We note that if previously-introduced quantities are written with the subscript n_f , such as $\tilde{I}_{C,\text{nf}}$ in Eq. (5.5), only the n_f -dependent final-state contributions should be retained in the corresponding functions.

The last contribution that we need to discuss arises from the action of one single-soft and one hard-collinear operator on the double-real cross section (see the second and third

terms on the right-hand side of Eq. (4.12), and the first term on the right-hand side of Eq. (4.13)). We find

$$\begin{aligned}
& \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} \left[\left\langle \bar{S}_m C_{im} \Delta^{(m)} \left\langle S_n \left(\Theta_{mn} F_{LM}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + F_{LM}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] + n_f F_{LM}^{\mathcal{A}_5}[\mathbf{m}_{(q'\bar{q}'), \mathbf{n}_g} \right) \right\rangle \right. \right. \\
& \left. \left. + \left\langle S_n \left\langle \bar{S}_m C_{im} \Delta^{(mn)} \Theta_{nm} F_{LM}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] \right\rangle \right\rangle \right] \\
& = [\alpha_s]^2 \left[\left\langle I_S(\epsilon) I_C(\epsilon) \cdot F_{LM}^{\mathcal{A}_0} \right\rangle + \frac{1}{\epsilon} \left\langle \mathcal{P}_{gg}^{\text{gen}} \otimes [I_S(\epsilon) \cdot F_{LM}^{\mathcal{A}_0}] + [I_S(\epsilon) \cdot F_{LM}^{\mathcal{A}_0}] \otimes \mathcal{P}_{qq}^{\text{gen}} \right\rangle \right] \quad (5.7) \\
& + \frac{[\alpha_s]^2}{2\epsilon^3} C_A h_c(\epsilon) \left\langle \mathcal{P}_{gg}^{(4), \text{gen}} \otimes F_{LM}^{\mathcal{A}_0} + F_{LM}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{(4), \text{gen}} \right\rangle \\
& + \frac{[\alpha_s]^2}{\epsilon^2} h_c(\epsilon) \left\langle \left[C_A \left(I_C^{(4)}(\epsilon) + \sum_{i \in \mathcal{H}_{f,q}} \sigma_{i,q \rightarrow gq} \right) + (2C_F - C_A) \sum_{i \in \mathcal{H}_{f,g}} \sigma_{i,g \rightarrow q\bar{q}} \right) \right. \right. \\
& \left. \left. + 2(C_F - C_A) I_{C,n_f}^{(4)}(\epsilon) \right] \cdot F_{LM}^{\mathcal{A}_0} \right\rangle + \frac{[\alpha_s]^2}{2\epsilon^3} C_A h_c(\epsilon) \left\langle \mathcal{P}_{gg}^{(4), \text{gen}} \otimes F_{LM}^{\mathcal{A}_0} + F_{LM}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{(4), \text{gen}} \right\rangle,
\end{aligned}$$

where

$$\begin{aligned}
\sigma_{i,q \rightarrow gq} &= \frac{1}{2\epsilon} \left[e^{-2\epsilon L_{\max}} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \Gamma_{i,q \rightarrow gq} - \Gamma_{i,q \rightarrow gq}^{(4)} \right], \\
\sigma_{i,g \rightarrow q\bar{q}} &= 2n_f \frac{1}{2\epsilon} \left[e^{-2\epsilon L_{\max}} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \Gamma_{i,g \rightarrow q\bar{q}} - \Gamma_{i,g \rightarrow q\bar{q}}^{(4)} \right]. \quad (5.8)
\end{aligned}$$

As was pointed out in Ref. [1], one has to be careful with the order of soft and collinear operators when computing the hard-collinear limits of the double-real contributions. At the same time, the fact that in the present study we allow for more complex unresolved final states does not impact the computation significantly. The remaining double-unresolved terms in Eq. (4.12) and Eq. (4.13) that we have yet to analyze and which contribute to $\mathcal{O}(\epsilon^{-2})$ involve two collinear limits; we discuss them in the following section.

5.2 Soft-regulated double-collinear contributions

In this section, we discuss the double-unresolved terms that originate from collinear limits. Specifically, we consider the double-collinear limits present in the fourth and fifth terms on the right-hand side of Eq. (4.12), and in the second and third terms on the right-hand side of Eq. (4.13).

We begin with the terms that do not involve the C_{mn} limit. We write

$$\Sigma_{\text{DC}} = \Sigma_{\text{DC}}^{\text{dc}} + \Sigma_{\text{DC}}^{\text{tc}}, \quad (5.9)$$

where $\Sigma_{\text{DC}}^{\text{dc}}$ contains limits where the unresolved partons (mn) become collinear to two different partons, while in $\Sigma_{\text{DC}}^{\text{tc}}$ they become collinear to the same parton.⁵ The expressions

⁵The notation “dc” and “tc” refers to their origin in double-collinear and triple-collinear partitions.

for these two terms read

$$\begin{aligned}
\Sigma_{\text{DC}}^{\text{dc}} &= \mathcal{P}_{\mathcal{A}_0} \sum_{(ij)} \left\langle \bar{S}_n \bar{S}_m C_{jn} C_{im} \Delta^{(\text{mn})} \left(\frac{1}{2} F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] \right. \right. \\
&\quad \left. \left. + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q'\bar{q}')}], \mathbf{n}_g] + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_q] + n_f^2 F_{\text{LM}}^{\mathcal{A}_6}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_{(q''\bar{q}'')}] \right) \right\rangle, \\
\Sigma_{\text{DC}}^{\text{tc}} &= \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} \frac{1}{2} \left\langle \bar{S}_n \bar{S}_m C_{in} C_{im} \Delta^{(\text{mn})} \left(F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q')}, \mathbf{n}_{\bar{q}'}] \right) \right\rangle \quad (5.10) \\
&\quad + \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} \frac{1}{2} \left\langle \bar{S}_n (C_{in} C_{im} + C_{im} C_{in}) \Delta^{(\text{mn})} \left(F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_g] \right. \right. \\
&\quad \left. \left. + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_q] \right) \right\rangle,
\end{aligned}$$

where (ij) stands for $i, j \in \mathcal{H}$, with $i \neq j$. In Eq. (5.10), the function $F_{\text{LM}}^{\mathcal{A}_6}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_{(q''\bar{q}'')}]$ is defined as

$$F_{\text{LM}}^{\mathcal{A}_6}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_{(q''\bar{q}'')}] = F_{\text{LM}}^{\mathcal{A}_6}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_{q''}] + F_{\text{LM}}^{\mathcal{A}_6}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_{\bar{q}''}]. \quad (5.11)$$

We start with the discussion of $\Sigma_{\text{DC}}^{\text{dc}}$. Its calculation follows Ref. [1], and the final expression is equivalent to the product of two NLO-like expressions, since the two collinear limits C_{im} and C_{jn} (with $i \neq j$) are completely decoupled from each other. Combining the expressions that we obtain from the double-collinear limits of each F_{LM} function contributing to $\Sigma_{\text{DC}}^{\text{dc}}$ to reconstruct the quark and gluon anomalous dimensions, defined in Eqs (3.19) and (3.25), respectively, we obtain

$$\begin{aligned}
\Sigma_{\text{DC}}^{\text{dc}} &= \frac{[\alpha_s]^2}{\epsilon^2} \left[\frac{1}{2} \sum_{(ij)} \langle \Gamma_{i,fi} \Gamma_{j,fj} \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle \right. \\
&\quad \left. + \sum_{\substack{i \in \mathcal{H} \\ i \neq a}} \langle \mathcal{P}_{gg}^{\text{gen}} \otimes (\Gamma_{i,fi} \cdot F_{\text{LM}}^{\mathcal{A}_0}) \rangle + \sum_{\substack{i \in \mathcal{H} \\ i \neq b}} \langle (\Gamma_{i,fi} \cdot F_{\text{LM}}^{\mathcal{A}_0}) \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle \right]. \quad (5.12)
\end{aligned}$$

We note that the first term contains a factorized contribution of the anomalous dimensions of partons i and j . To identify the universal operator I_C and its square, we need to supplement Eq. (5.12) with $i = j$ double-collinear terms coming from $\Sigma_{\text{DC}}^{\text{tc}}$.

Although the analysis of $\Sigma_{\text{DC}}^{\text{tc}}$ is largely analogous to what was done in Ref. [1], it is useful to remind the reader that collinear limits for partons emitted off the same line are intertwined. However, in order to write $\Sigma_{\text{DC}}^{\text{tc}}$ in terms of two factorized anomalous dimensions or two splitting functions convoluted in exactly the same way as in the perturbative solution of the Altarelli-Parisi equation, it is useful to disentangle them.

We start with the initial-state radiation. In this case, only the terms in the first line on the right-hand side of $\Sigma_{\text{DC}}^{\text{tc}}$ in Eq. (5.10) give a nonzero contribution, since the remaining terms would involve an initial-state flavor change, and are thus projected out by $\mathcal{P}_{\mathcal{A}_0}$. Therefore, the computation of the double-collinear limit in the case $i = a$ or $i = b$ follows the procedure described in Ref. [1], except that here we also consider the case where the

initial-state parton is a gluon. As an example, we consider the function $F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g]$ in $\Sigma_{\text{DC}}^{\text{tc}}$. Computing the collinear limit explicitly for $i = a$, we obtain

$$\begin{aligned} \frac{1}{2} \langle \bar{\mathcal{S}}_{\mathbf{m}} \bar{\mathcal{S}}_{\mathbf{n}} C_{am} C_{an} \Delta^{(\text{mn})} F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] \rangle &= \frac{[\alpha_s]^2}{2\epsilon^2} \left[\langle \Gamma_{a,g}^2 \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle + 2 \langle \mathcal{P}_{gg}^{\text{gen}} \otimes (\Gamma_{a,g} \cdot F_{\text{LM}}^{\mathcal{A}_0}) \rangle \right. \\ &\quad \left. + \langle [\mathcal{P}_{gg}^{\text{gen}} \bar{\otimes} \mathcal{P}_{gg}^{\text{gen}}] \otimes F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle G_{gg} \otimes F_{\text{LM}}^{\mathcal{A}_0} \rangle \right]. \end{aligned} \quad (5.13)$$

We note that the first term on the right-hand side of Eq. (5.13) combines with the first term on the right-hand side of Eq. (5.12) and contributes to the construction of I_{C}^2 . The second contribution in Eq. (5.13) combines with the first term in the second line of Eq. (5.12), leading to the operator I_{C} convoluted with $\mathcal{P}_{gg}^{\text{gen}}$. Next, the term involving $[\mathcal{P}_{gg}^{\text{gen}} \bar{\otimes} \mathcal{P}_{gg}^{\text{gen}}]$, where the $\bar{\otimes}$ convolution is defined as

$$f_1(z_1, E) \bar{\otimes} f_2(z_2, E) = \int dz_1 dz_2 f_1(z_1, E) f_2(z_2, z_1 E) \delta(z - z_1 z_2), \quad (5.14)$$

is also expected, as it must combine with the contribution $[\hat{P}_{gg}^{(0)} \otimes \hat{P}_{gg}^{(0)}]$ arising from the PDF renormalization. Finally, the function G_{gg} in Eq. (5.13) is similar to the function G_i introduced in Ref. [1]. Here, we define it so that it can also be applied for an initial-state quark. We then write

$$G_{\alpha\alpha}(z, E_i) = (\Gamma_{i,\alpha} - \Gamma_{z \cdot i, \alpha}) \mathcal{P}_{\alpha\alpha}^{\text{gen}}(z, E_i), \quad \text{with } \Gamma_{z \cdot i, \alpha} = \Gamma_{i,\alpha} \big|_{E_i \mapsto z E_i}, \quad (5.15)$$

where $(i, \alpha) \in \{(a, g), (b, q)\}$. The function in the above equation incorporates the modifications required by the non-trivial dependence of the double-collinear phase space of two unresolved partons on their energies.

A similar calculation has to be performed for the final-state splittings but it is slightly more intricate because, in addition to the phase-space factors that appear in the collinear limits (discussed above), applying those limits to the damping factors $\Delta^{(\text{mn})}$ leads to energy fractions of final-state unresolved partons, and one needs to keep track of them to see simplified structures appearing in the final result. Although this complication was already present in the calculation reported in Ref. [1], it becomes more involved in the current case, as there are now different types of partons in the final state. To illustrate this, we consider the double-collinear limit

$$\langle \bar{\mathcal{S}}_{\mathbf{m}} \bar{\mathcal{S}}_{\mathbf{n}} C_{in} C_{im} \Delta^{(\text{mn})} F_{\text{LM}}^{ab}[\dots, i, \dots | \mathbf{m}, \mathbf{n}] \rangle, \quad i \in \mathcal{H}_{\text{f}}. \quad (5.16)$$

The action of the first collinear operator clusters partons i and \mathbf{m} into a parton $[i\mathbf{m}]$, while the second operator combines $[i\mathbf{m}]$ and \mathbf{n} to produce parton $[i\mathbf{mn}]$. We find

$$C_{in} C_{im} \Delta^{(\text{mn})} F_{\text{LM}}^{ab}[\dots, i, \dots | \mathbf{m}, \mathbf{n}] = \frac{g_{\text{s,b}}^4}{\rho_{im} \rho_{in}} \frac{P_{[imn][im]}(z_n) P_{[im]i}(z_m)}{E_m E_n E_{[im]} E_{[imn]}} F_{\text{LM}}^{ab}[\dots, [i\mathbf{mn}], \dots], \quad (5.17)$$

where $E_{[im]} = E_i + E_m$ and $E_{[imn]} = E_i + E_m + E_n$ are the energies of the clustered partons $[im]$ and $[imn]$, respectively, and we have used

$$z_{\mathbf{m}} = \frac{E_i}{E_{[im]}}, \quad z_{\mathbf{n}} = \frac{E_{[im]}}{E_{[imn]}}, \quad C_{in} C_{im} \Delta^{(\text{mn})} = z_n z_{\mathbf{m}} = \frac{E_i}{E_{[imn]}}, \quad (5.18)$$

in addition to the collinear limits described by the splitting functions $P_{[im]i}$ and $P_{[imn][im]}$.

The expression in Eq. (5.17) needs to be integrated over the collinear phase space of the unresolved partons \mathbf{m}, \mathbf{n} , and the resolved parton i , keeping the sum of the three energies fixed. We would like to write the result as the product of the splittings $\Gamma_{i,\alpha \rightarrow \rho\sigma}$ defined in Eq. (A.19), which will allow us to combine these expressions with those in Eq. (5.12). This does not happen automatically, because the integration over the variable z_m , which describes the *internal* splitting, introduces an *additional* factor $z_n^{-2\epsilon}$ into the integral over the *external* splitting variable z_n . Writing this additional factor as

$$z_n^{-2\epsilon} = 1 + (z_n^{-2\epsilon} - 1), \quad (5.19)$$

and integrating over z_n , we can obtain the desired product of generalized anomalous dimensions from the first term on the right-hand side in the above equation, and a new contribution from the second which has a stronger ϵ -suppression. Generalizing the discussion in Ref. [1], we write the double-collinear soft-subtracted limit as

$$\begin{aligned} & \langle \bar{S}_m \bar{S}_n C_{in} C_{im} \Delta^{(mn)} F_{LM}^{ab}[\dots, i, \dots | \mathbf{m}, \mathbf{n}] \rangle \\ &= [\alpha_s]^2 \left\langle \left[\Gamma_{[imn],[im] \rightarrow im} \Gamma_{[imn],[imn] \rightarrow [im]n} + G_{[imn]} \left| \frac{f(z), [im] \rightarrow im}{\tilde{f}(z), [imn] \rightarrow [im]n} \right| F_{LM}^{ab}[\dots, [imn], \dots] \right] \right\rangle, \end{aligned} \quad (5.20)$$

where

$$\begin{aligned} G_{[imn]} \left| \frac{f(z), [im] \rightarrow im}{\tilde{f}(z), [imn] \rightarrow [im]n} \right| &= \left[\left(\frac{2E_{[imn]}}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^2 \left[\gamma_{f(z), [im] \rightarrow im}^{22}(L_{[imn]}) \right. \\ &+ \delta_{[im]i} \frac{\mathbf{T}_{[im]}^2}{\epsilon} e^{-2\epsilon L_{[imn]}} \left. \left[\gamma_{\tilde{f}(z), [imn] \rightarrow [im]n}^{42}(L_{[imn]}) - \gamma_{\tilde{f}(z), [imn] \rightarrow [im]n}^{22}(L_{[imn]}) \right] \right]. \end{aligned} \quad (5.21)$$

Definitions relevant for the two equations above are collected in Appendix A.3.1. Note that we introduced two different functions $f(z)$ and $\tilde{f}(z)$ to describe the weights that appear in the integrands over energy fractions of the external and internal collinear splittings. Since these weights arise from the collinear limits of $\Delta^{(mn)}$ factors, they are always equal to the relevant energy fraction z , c.f. Eq. (5.18). However, as we pointed out several times, we benefit from combining various collinear limits before integrating over energy fractions since in some cases the weights may disappear. Because of this, we find it convenient to introduce two different weights $f(z), \tilde{f}(z)$ in the definition of G_α in Eq. (5.21). Additionally, we note that in the case where the collinear limits in Eq. (5.20) are taken in the order $C_{im}C_{in}$ instead of $C_{in}C_{im}$, Eq. (5.20) remains valid as long as we exchange $\mathbf{m} \leftrightarrow \mathbf{n}$.

To account for suitable combinations of the collinear limits, we introduce the shorthand notation

$$G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^g = G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^{z, g \rightarrow gg} + 2n_f G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^{z, g \rightarrow q\bar{q}}, \quad (5.22)$$

to describe the gluon splittings. Furthermore, using the properties of the generalized quark anomalous dimensions, we write

$$G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^q = G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^{z, q \rightarrow qq} + G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^{z, q \rightarrow gq} \equiv G_i \Big|_{z, \alpha \rightarrow \beta\gamma}^{1, q \rightarrow qq}, \quad (5.23)$$

where the second equality follows from Eq. (3.20).

Each contribution to $\Sigma_{\text{DC}}^{\text{tc}}$ in Eq. (5.10) can be described using Eq. (5.13) for initial-state radiation and Eq. (5.20) for final-state radiation. All these terms can then be combined with Eq. (5.12) to obtain the final result for the double-collinear contribution. It reads

$$\begin{aligned}
\Sigma_{\text{DC}} = & \frac{[\alpha_s]^2}{2} \left\langle \left\{ I_C^2 + \frac{1}{\epsilon^2} \left[\sum_{i \in \mathcal{H}_{f,q}} \Gamma_{i,q \rightarrow gq} (\Gamma_{i,g} - \Gamma_{i,q}) + \sum_{i \in \mathcal{H}_{f,g}} 2n_f \Gamma_{i,g \rightarrow q\bar{q}} (\Gamma_{i,q} - \Gamma_{i,g}) \right. \right. \right. \\
& + \sum_{i \in \mathcal{H}_{f,g}} (G_i |_{z,g \rightarrow gg}^g + 2n_f G_i |_{z,g \rightarrow q\bar{q}}^q) + \sum_{i \in \mathcal{H}_{f,q}} (G_i |_{z,q \rightarrow qg}^q + G_i |_{z,q \rightarrow gq}^g) \cdot F_{\text{LM}}^{\mathcal{A}_0} \left. \right\} \\
& + \frac{[\alpha_s]^2}{2\epsilon^2} \left[\langle [\mathcal{P}_{gg}^{\text{gen}} \otimes \mathcal{P}_{gg}^{\text{gen}}] \otimes F_{\text{LM}}^{\mathcal{A}_0} + F_{\text{LM}}^{\mathcal{A}_0} \otimes [\mathcal{P}_{qq}^{\text{gen}} \otimes \mathcal{P}_{qq}^{\text{gen}}] \rangle \right. \\
& + \frac{[\alpha_s]^2}{2\epsilon^2} \langle G_{gg} \otimes F_{\text{LM}}^{\mathcal{A}_0} + F_{\text{LM}}^{\mathcal{A}_0} \otimes G_{qq} \rangle \\
& \left. + \frac{[\alpha_s]^2}{\epsilon} \left[\langle \mathcal{P}_{gg}^{\text{gen}} \otimes [I_C(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0}] \rangle + \langle [I_C(\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0}] \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle + \langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle \right] \right. \end{aligned} \quad (5.24)$$

Finally, we consider those double-collinear terms that involve the operator C_{mn} , i.e. the limit where the two unresolved partons \mathbf{m} and \mathbf{n} become collinear to each other, forming a clustered parton $[\mathbf{mn}]$, which then goes collinear to a hard parton $i \in \mathcal{H}$. Such contributions are found in the fifth term on the right-hand side of Eq. (4.12), and the third term on the right-hand side of Eq. (4.13). Their computation closely follows the discussion in Ref. [1]. For this reason, we limit ourselves to reporting the final expressions. They read

$$\begin{aligned}
& \Sigma_{\mathbf{m}||\mathbf{n}}^{(1)} \\
& = \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} \frac{N_{\mathbf{m}||\mathbf{n}}(\epsilon)}{2} \left\langle \bar{S}_{\mathbf{m}} C_{i\mathbf{m}} \sigma_{i\mathbf{m}}^{-\epsilon} \left[(\mathbb{1} - 2\Theta_{\mathbf{mn}} S_{\mathbf{n}}) \Delta^{(\mathbf{m})} C_{\mathbf{mn}} (F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q')}, \mathbf{n}_{(\bar{q}')}] \right) \right] \right\rangle \\
& = \frac{[\alpha_s]^2}{\epsilon} N_{\text{sc}}^{(b,d)} \left\langle \gamma_g^{22}(0) \left(I_C^{(4)}(\epsilon) - \sum_{i \in \mathcal{H}_{f,q}} \frac{\Gamma_{i,q \rightarrow gq}^{(4)}}{2\epsilon} - \sum_{i \in \mathcal{H}_{f,g}} \frac{2n_f \Gamma_{i,g \rightarrow q\bar{q}}^{(4)}}{2\epsilon} \right) \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\
& + \frac{[\alpha_s]^2}{2\epsilon^2} N_{\text{sc}}^{(b,d)} \left\langle \gamma_g^{22}(0) \left[\mathcal{P}_{gg}^{(4),\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} + F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{(4),\text{gen}} \right] \right\rangle, \end{aligned} \quad (5.25)$$

and

$$\begin{aligned}
& \Sigma_{\mathbf{m}||\mathbf{n}}^{(2)} = \mathcal{P}_{\mathcal{A}_0} \sum_{i \in \mathcal{H}} N_{\mathbf{m}||\mathbf{n}}(\epsilon) \left\langle C_{i\mathbf{m}} \sigma_{i\mathbf{m}}^{-\epsilon} \Delta^{(\mathbf{m})} C_{\mathbf{mn}} (F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] + n_f F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q'\bar{q}')}, \mathbf{n}_g]) \right\rangle \\
& = \frac{[\alpha_s]^2}{\epsilon} N_{\text{sc}}^{(b,d)} \sum_{i \in \mathcal{H}_{f,g}} \left\langle \left[\gamma_q^{22}(L_i) \frac{2n_f \Gamma_{i,g \rightarrow q\bar{q}}^{(4)}}{2\epsilon} + \frac{1}{2\epsilon} H_{i,g \rightarrow q\bar{q}} \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\
& + \frac{[\alpha_s]^2}{\epsilon} N_{\text{sc}}^{(b,d)} \sum_{i \in \mathcal{H}_{f,q}} \left\langle \left[\gamma_q^{22}(L_i) \frac{\Gamma_{i,q \rightarrow gq}^{(4)}}{2\epsilon} + \frac{1}{2\epsilon} H_{i,q \rightarrow gq} \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle. \end{aligned} \quad (5.26)$$

In the above equations, we have used

$$\begin{aligned}
\gamma_g^{22}(L_i) &= \gamma_{z,g \rightarrow gg}^{22}(L_i) + 2n_f \gamma_{z,g \rightarrow q\bar{q}}^{22}, \\
\gamma_q^{22}(L_i) &= \gamma_{z,q \rightarrow qg}^{22}(L_i) + \gamma_{z,q \rightarrow gq}^{22} \equiv \gamma_{1,q \rightarrow qg}^{22}(L_i), \end{aligned} \quad \text{with } \gamma_{f_i}^{22}(0) \equiv \gamma_{f_i}^{22}(L_i = 0), \quad (5.27)$$

and

$$\begin{aligned}
H_{i,q \rightarrow gq} &= \frac{\mathbf{T}_q^2}{\epsilon} \left[e^{-2\epsilon L_i} \Gamma_{i,q \rightarrow gq}^{(4)} - e^{-2\epsilon L_{\max}} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \Gamma_{i,q \rightarrow gq} \right], \\
H_{i,g \rightarrow q\bar{q}} &= 2n_f \frac{\mathbf{T}_q^2}{\epsilon} \left[e^{-2\epsilon L_i} \Gamma_{i,g \rightarrow q\bar{q}}^{(4)} - e^{-2\epsilon L_{\max}} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \Gamma_{i,g \rightarrow q\bar{q}} \right].
\end{aligned} \tag{5.28}$$

We have now analyzed all the double-unresolved terms in Eq. (4.12) and Eq. (4.13) which produce singularities of $\mathcal{O}(\epsilon^{-2})$ or stronger, and have written the corresponding expressions in terms of the virtual, soft, and collinear operators $I_{V,S,C}$. In the following section, we will employ these results to discuss the cancellation of infrared singularities.

6 Cancellation of poles

In this section, we use the results obtained in the previous section to demonstrate the cancellation of infrared singularities in the NNLO QCD corrections to the process \mathcal{A}_0 . We have pointed out in Ref. [1] that this task is simplified significantly if one combines virtual, soft, and collinear operators I_V , I_S and I_C into a single color-correlated quantity I_T from Eq. (3.31), which is ϵ -finite. Indeed, identifying this operator in the NNLO contribution to cross sections allowed us to cancel the poles through $1/\epsilon^2$ “by eye”. Furthermore, we observed that combining the highest-order singularities into the I_T operator in two subsequent orders of QCD perturbation theory lead to an iterative structure consistent with the exponentiation

$$d\hat{\sigma} \sim \langle e^{[\alpha_s]I_T} \cdot F_{\text{LM}} \rangle. \tag{6.1}$$

This, together with the fact that we were able to identify the I_T operator for the more complex process that we consider in this paper (cf. Section 3), suggests that it would be beneficial to discuss the cancellation of $1/\epsilon$ poles by rewriting $d\hat{\sigma}^{\text{NNLO}}$ using the I_T^2 and I_T operators, as far as possible. We discuss this in the following sections, beginning with the single-unresolved contributions, and then turning to the double-unresolved ones.

6.1 Single-unresolved terms

In this section, we discuss the cancellation of singularities for the single-unresolved contribution, and highlight the need to identify the proper combinations of quark collinear splittings to simplify the calculation. The relevant expressions can be found in Eqs (4.14), (4.15), and (4.20). We begin by considering contributions which are proportional to the operator $\mathcal{O}_{\text{NLO}}^{(\text{m})}$, and first study those terms which do not involve collinear operators, namely the real-virtual corrections (see $\Sigma_{\text{SU}}^{\mathcal{A}_n}[\mathbf{m}]_{\text{RV}}$ in Eq. (4.20)) and the soft limit of the double-real ones (see the first terms in Eqs. (4.14) and (4.15)). As before, the unresolved parton in the former contribution can be either a quark or a gluon, while the pair of unresolved partons in the latter contribution can be either $(\mathbf{m}_g \mathbf{n}_g)$ or $(\mathbf{m}_q \mathbf{n}_g)$. Their sum evaluates to

$$\begin{aligned}
&\left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} \left[F_{\text{RV}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{RV}}^{\mathcal{A}_1}[\mathbf{m}_q] + S_n \left(\Theta_{\text{mn}} F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] + F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] \right) \right] \right\rangle \\
&= [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} [I_V + I_S(E_{\text{m}})] \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \rangle
\end{aligned} \tag{6.2}$$

$$+ [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} [I_V + I_S(E_{\text{max}})] \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \rangle + \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} (F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}_q]) \rangle,$$

where $F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}]$ is defined analogously to $F_{\text{LV,fin}}^{\mathcal{A}_1}$. We note that the I_V and I_S operators in Eq. (6.2) act on the $(N+1)$ hard partons of functions $F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g]$ and $F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q]$. Additionally, the energy dependencies of the two I_S operators appearing on the right-hand side of the above equation are different. This is due to the presence of the energy-ordering function Θ_{mn} in the case of the two-gluon unresolved final state, and its absence in the case of the quark-gluon one. We note that $I_S(E_{\text{m}})$ and $I_S(E_{\text{max}})$ are defined in Eq. (A.38), where in $I_S(E_{\text{m}})$ the substitution $E_{\text{max}} \mapsto E_{\text{m}}$ is applied.

In order to combine the terms in Eq. (6.2) into I_{T} operators, we require collinear operators I_{C} . These will arise from the second and the third term in Eq. (4.14) if $(\text{mn}) \in \text{DS}$, and from the second and the fourth term in Eq. (4.15) if $(\text{mn}) \in \mathcal{DS}$. Such terms originate from different configurations of unresolved partons, but when combined they produce anomalous dimensions that appear in the collinear operator I_{C} for a process \mathcal{A}_0 with an additional jet.

To show this, we first write the required contribution for the sum of the gg and $q'q'$ unresolved final states, i.e., for $(\text{mn}) \in \text{DS}$. The result, which can be easily obtained by generalizing Eq. (5.3) in Ref. [1], reads

$$\begin{aligned} \mathcal{P}_{\mathcal{A}_0} & \left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \left[\sum_{i \in \mathcal{H}} (\mathbb{1} - S_{\text{n}} \Theta_{\text{mn}}) C_{i\text{n}} \Delta^{(\text{mn})} + \frac{1}{2} \Delta^{(\text{m})} (\mathbb{1} - 2S_{\text{n}} \Theta_{\text{mn}}) C_{\text{mn}} \right] F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_g, \mathbf{n}_g] \right. \\ & \left. + \frac{n_{\text{f}}}{2} \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} C_{\text{mn}} F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q', \mathbf{n}_{\bar{q}'}]}] \right\rangle \\ & = [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} (\mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \otimes \mathcal{P}_{qq}^{\text{gen}}) \rangle \\ & + [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} \left(I_{\text{C}}(E_{\text{m}}) - \sum_{i \in \mathcal{H}_{\text{f},g}} \frac{2n_{\text{f}} \Gamma_{i,g \rightarrow q\bar{q}}}{\epsilon} - \sum_{i \in \mathcal{H}_{\text{f},q}} \frac{\Gamma_{i,q \rightarrow gq}}{\epsilon} \right) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \right\rangle. \end{aligned} \quad (6.3)$$

We note that the collinear operator $I_{\text{C}}(E_{\text{m}})$ in the above formula includes the contribution of the parton m and, similarly to $I_S(E_{\text{m}})$ discussed above, is computed with the replacement $E_{\text{max}} \mapsto E_{\text{m}}$.

For $(\text{mn}) \in \mathcal{DS}$, we find

$$\begin{aligned} \mathcal{P}_{\mathcal{A}_0} & \sum_{i \in \mathcal{H}} \left\{ \left\langle \left[\mathcal{O}_{\text{NLO}}^{(\text{n})} C_{i\text{m}} + \mathcal{O}_{\text{NLO}}^{(\text{m})} \bar{S}_{\text{n}} C_{i\text{n}} \right] \Delta^{(\text{mn})} F_{\text{LM}}^{\mathcal{A}_4}[\mathbf{m}_q, \mathbf{n}_g] \right\rangle + [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} \langle C_{\text{mn}} F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \rangle \right\rangle \\ & + \left\langle n_{\text{f}} \mathcal{O}_{\text{NLO}}^{(\text{n})} C_{i\text{m}} \Delta^{(\text{mn})} F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q' \bar{q}')}], \mathbf{n}_g \right\rangle + n_{\text{f}} \mathcal{O}_{\text{NLO}}^{(\text{m})} C_{i\text{n}} \Delta^{(\text{mn})} F_{\text{LM}}^{\mathcal{A}_5}[\mathbf{m}_{(q' \bar{q}')}], \mathbf{n}_q \right\rangle \Big\} \\ & = [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} \left[\mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] + F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \otimes \mathcal{P}_{qq}^{\text{gen}} \right] \right\rangle + [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} \right. \\ & \times \left[I_{\text{C}}(E_{\text{max}}) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] + \left(\sum_{i \in \mathcal{H}_{\text{f},g}} \frac{2n_{\text{f}} \Gamma_{i,g \rightarrow q\bar{q}}}{\epsilon} + \sum_{i \in \mathcal{H}_{\text{f},q}} \frac{\Gamma_{i,q \rightarrow gq}}{\epsilon} \right) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \right] \right\rangle. \end{aligned} \quad (6.4)$$

Upon comparing Eqs (6.3) and (6.4), it becomes apparent that *in their sum*, the anomalous dimensions that do not combine into an I_{C} operator cancel. Furthermore, the two collinear

operators that appear carry the appropriate energy dependence needed to reconstruct $I_T(E_{\max})$ and $I_T(E_m)$.

The remaining $1/\epsilon$ divergences that reside in the generalized splitting functions $\mathcal{P}_{gg}^{\text{gen}}$ and $\mathcal{P}_{qq}^{\text{gen}}$ in Eqs (6.3) and (6.4) are removed by \mathcal{O}_{NLO} -dependent terms that arise when PDF renormalization is applied to the finite remainder of the NLO cross section. Combining the two, we arrive at an ϵ -finite expression

$$[\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(m)} \Delta^{(m)} \left[I_T(E_m) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] + I_T(E_{\max}) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \right. \right. \\ \left. \left. + \mathcal{P}_{gg}^{\text{NLO}} \otimes F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] + F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \otimes \mathcal{P}_{qq}^{\text{NLO}} + F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}_q] \right] \right\rangle. \quad (6.5)$$

In the above equation, $F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}]$ is defined in Eq. (3.5), and the generalized collinear splitting functions $\mathcal{P}_{gg}^{\text{NLO}}$ and $\mathcal{P}_{qq}^{\text{NLO}}$ have already appeared in the discussion of the NLO corrections.

The remaining single-unresolved terms can easily be identified as being ϵ -finite. For example, the fourth and fifth term of Eq. (4.14), which are acted on by the subtraction operator $\mathcal{O}_{\text{NLO}}^{(i,m)}$, contain a hard-collinear limit that produces a $1/\epsilon$ pole, but also a prefactor of $\mathcal{O}(\epsilon)$. The same argument applies to the third, fifth, and sixth terms of Eq. (4.15), while the final two lines of Eq. (4.14) are manifestly ϵ -finite. The manipulations required to bring such terms to their final form are therefore minimal. The complete expression for the single-unresolved contribution to $d\hat{\sigma}_{\text{NLO}}^{\mathcal{A}_0}$ can be found in Eq. (7.3).

6.2 Double-boosted contribution

Having discussed the treatment of singularities arising in the single-unresolved contributions, we now turn to those terms in which two partons are unresolved. We find it convenient to organize them according to the number of initial-state partons which are boosted, and we begin with the double-boosted terms. These arise when soft-subtracted collinear operators act on *both* incoming partons, leading to double convolutions of generalized splitting functions with $F_{\text{LM}}^{\mathcal{A}_0}$. The relevant term appears as the last entry in Eq. (5.24). Further terms of this type arise from the collinear renormalization of parton distribution functions. Combining such contributions, we find

$$\Sigma_{\text{DU}}^{\text{db}} = \frac{[\alpha_s]^2}{\epsilon^2} \left[\langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle + \tilde{c}_\epsilon^2 \langle \hat{P}_{gg}^{(0)} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \hat{P}_{qq}^{(0)} \rangle \right. \\ \left. + \tilde{c}_\epsilon \langle \hat{P}_{gg}^{(0)} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{gen}} \rangle + \tilde{c}_\epsilon \langle \mathcal{P}_{gg}^{\text{gen}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \hat{P}_{qq}^{(0)} \rangle \right] \\ = [\alpha_s]^2 \langle \mathcal{P}_{gg}^{\text{NLO}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{NLO}} \rangle + \mathcal{O}(\epsilon), \quad (6.6)$$

where $\tilde{c}_\epsilon = \Gamma(1 - \epsilon)/e^{\epsilon\gamma_E}$. To derive the above result, we used the relation between $\mathcal{P}_{\alpha\beta}^{\text{gen}}$, $\hat{P}_{\alpha\beta}^{(0)}$, and $\mathcal{P}_{\alpha\beta}^{\text{NLO}}$, which we already presented in Eq. (3.32) in the context of NLO corrections.

6.3 Single-boosted contributions

We continue with the analysis of the single-boosted contributions. It is straightforward to identify terms proportional to a boost, convoluted with the operators I_V , I_S , and I_C in Eqs (5.5), (5.7), and (5.25), respectively, and to combine these into a boost acting on

the IR-finite operator I_T . Similarly, terms proportional to the finite remainder of the one-loop amplitude $F_{\text{LV,fin}}^{\mathcal{A}_0}$ in Eq. (5.5) can be combined with terms originating from the PDF renormalization to obtain an IR-finite result, as happens at NLO. We note that this renders finite *all* contributions with color correlations.

We therefore concentrate on the remaining terms proportional to $F_{\text{LM}}^{\mathcal{A}_0}$ that cannot be absorbed in I_T , and consider those that exhibit $\mathcal{O}(\epsilon^{-3})$ and $\mathcal{O}(\epsilon^{-2})$ singularities. We combine the relevant terms from Eqs (5.5), (5.7), (5.24), and (5.25), with terms coming from the renormalizations of the PDFs.⁶ For simplicity, we report below the contribution associated with the boost of the initial-state parton a_g , which reads

$$\begin{aligned} \Sigma_{\text{DU}, \mathcal{O}(\epsilon^{-2})}^{\text{sb}, a} &= \frac{[\alpha_s]^2}{2\epsilon^3} \left\langle \left[C_A h_c(\epsilon) (\mathcal{P}_{gg}^{(4), \text{gen}} - \mathcal{P}_{gg}^{1\text{L}, \text{gen}}) + \epsilon G_{gg} \right] \otimes F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\ &+ \frac{[\alpha_s]^2}{2\epsilon^2} \left\{ \left\langle \left[N_{\text{sc}}^{(b,d)} \gamma_g^{22}(0) \mathcal{P}_{gg}^{(4), \text{gen}} - 2\tilde{c}_\epsilon \beta_0 \mathcal{P}_{gg}^{\text{gen}} - \tilde{c}_\epsilon^2 \beta_0 \hat{P}_{gg}^{(0)} \right] \otimes F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \right. \\ &+ \left\langle \left[[\mathcal{P}_{gg}^{\text{gen}} \bar{\otimes} \mathcal{P}_{gg}^{\text{gen}}] + 2\tilde{c}_\epsilon [\hat{P}_{gg}^{(0)} \bar{\otimes} \mathcal{P}_{gg}^{\text{gen}}] + \tilde{c}_\epsilon^2 [\hat{P}_{gg}^{(0)} \otimes \hat{P}_{gg}^{(0)}] \right. \right. \\ &\left. \left. + 2n_f \left([\mathcal{P}_{qg}^{\text{gen}} \bar{\otimes} \mathcal{P}_{qg}^{\text{gen}}] + 2\tilde{c}_\epsilon [\hat{P}_{qg}^{(0)} \bar{\otimes} \mathcal{P}_{qg}^{\text{gen}}] + \tilde{c}_\epsilon^2 [\hat{P}_{qg}^{(0)} \otimes \hat{P}_{qg}^{(0)}] \right) \right] \otimes F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \left. \right\}. \end{aligned} \quad (6.7)$$

To discuss the $1/\epsilon$ -divergences in Eq. (6.7), we need to expand various quantities that appear there in powers of ϵ . Using the definitions of $\mathcal{P}_{\alpha\alpha}^{(4), \text{gen}}$ and $\mathcal{P}_{\alpha\alpha}^{1\text{L}, \text{gen}}$ in Eqs (A.18) and (A.31), respectively, we obtain

$$\begin{aligned} C_A h_c(\epsilon) \left(\mathcal{P}_{\alpha\alpha}^{(4), \text{gen}}(z, E) - \mathcal{P}_{\alpha\alpha}^{1\text{L}, \text{gen}}(z, E) \right) &= 2\mathbf{T}_\alpha^2 \epsilon \log(z) \hat{P}_{\alpha\alpha}^{(0)}(z) + \mathcal{O}(\epsilon^2), \\ \epsilon G_{\alpha\alpha}(z, E) &= -2\mathbf{T}_\alpha^2 \epsilon \log(z) \hat{P}_{\alpha\alpha}^{(0)}(z) + \mathcal{O}(\epsilon^2). \end{aligned} \quad (6.8)$$

This equation, which is valid for both $\alpha = q$ and g , implies that, in spite of its appearance, the first line in Eq. (6.7) is only $\mathcal{O}(\epsilon^{-1})$.

To show that the second line in Eq. (6.7) is at most $\mathcal{O}(\epsilon^{-1})$, it is sufficient to use the expansions

$$\gamma_g^{22}(0) = \beta_0 + \mathcal{O}(\epsilon), \quad \tilde{c}_\epsilon = 1 + \mathcal{O}(\epsilon^2), \quad N_{\text{sc}}^{(b,d)} = 1 + \mathcal{O}(\epsilon), \quad (6.9)$$

and also (cf. Eq. (A.18))

$$\mathcal{P}_{\alpha\beta}^{(k), \text{gen}} = -\hat{P}_{\alpha\beta}^{(0)} + \mathcal{O}(\epsilon), \quad k = 2, 4, \quad (6.10)$$

which holds for all indices $(\alpha\beta)$. Turning to the last two lines in Eq. (6.7), we can use Eqs (6.9) and (6.10) to write

$$\begin{aligned} &[\mathcal{P}_{\alpha\beta}^{\text{gen}} \bar{\otimes} \mathcal{P}_{\gamma\delta}^{\text{gen}}] + 2\tilde{c}_\epsilon [\hat{P}_{\alpha\beta}^{(0)} \bar{\otimes} \mathcal{P}_{\gamma\delta}^{\text{gen}}] + \tilde{c}_\epsilon^2 [\hat{P}_{\alpha\beta}^{(0)} \otimes \hat{P}_{\gamma\delta}^{(0)}] \\ &= [\hat{P}_{\alpha\beta}^{(0)} \bar{\otimes} \hat{P}_{\gamma\delta}^{(0)}] - 2[\hat{P}_{\alpha\beta}^{(0)} \bar{\otimes} \hat{P}_{\gamma\delta}^{(0)}] + [\hat{P}_{\alpha\beta}^{(0)} \otimes \hat{P}_{\gamma\delta}^{(0)}] + \mathcal{O}(\epsilon) \\ &= \mathcal{O}(\epsilon), \end{aligned} \quad (6.11)$$

⁶We point out that contributions proportional to one-loop splitting functions that appear in the PDF renormalization only affect the single $1/\epsilon$ pole, therefore we do not discuss them in this section.

for any flavour-pair $\alpha\beta$ and $\gamma\delta$. Note that the last step in Eq. (6.11) follows from the fact that $\hat{P}^{(0)}$ is energy-independent, which implies that the $\bar{\otimes}$ and \otimes convolutions are identical (cf. Eq. (5.14)). We have therefore demonstrated that the single-boosted term in Eq. (6.7) has at most poles of $\mathcal{O}(\epsilon^{-1})$, and we will return to the remaining poles of this order at the end of this section. Moreover, these simple poles do not contain color correlations; all singularities multiplying products of color charges have already been combined into I_T , as discussed at the beginning of this section.

The contribution with the boost applied to the initial-state parton b_q can be obtained from Eq. (6.7) by replacing $q \leftrightarrow g$ in the indices of the initial-state splitting functions, and by removing the final line. This second action is required because, as mentioned earlier, we do not consider flavor-singlet configurations in this paper. Since the relations in Eqs (6.8), (6.10), and (6.11) are applicable for all values of the indices α and β , the above discussion can be repeated verbatim to demonstrate the cancellation of the poles through $\mathcal{O}(\epsilon^{-2})$ for this piece too.

6.4 Unboosted terms

Finally, we consider the unboosted terms. Many of these originate from virtual corrections and/or a soft gluon emission, and such contributions include color-correlated matrix elements squared with quadratic $\mathbf{T}_i \cdot \mathbf{T}_j$, triple $T_i^a T_j^b T_k^c$ and quartic $\{\mathbf{T}_i \cdot \mathbf{T}_j, \mathbf{T}_k \cdot \mathbf{T}_l\}$ products of color charges.

We first comment on the triple-color correlators. In Ref. [1], we have shown that they originate from three distinct sources: i) the soft limit of the real-virtual contributions [91], ii) commutators of the soft I_S and virtual I_V operators, and iii) double-virtual corrections [87, 88, 92]. All divergent triple-color correlated contributions cancel when terms from the three sources are combined. Since the calculations in Ref. [1] employed generic representation of color charges, they can be applied verbatim to the discussion of such contributions in the NNLO corrections to \mathcal{A}_0 ; hence, we do not discuss them further. The remaining color-correlated terms arise from various virtual corrections and/or soft limits, and are captured in the operators I_S and I_V in Eqs (5.1), (5.5), and (5.7).⁷ The terms proportional to the finite remainder of the one-loop amplitude $F_{LV, \text{fin}}^{\mathcal{A}_0}$ can easily be combined into the finite operator I_T . We focus on the remaining terms proportional to $F_{LM}^{\mathcal{A}_0}$, and combine these with the collinear operators in Eqs (5.24), (5.25), and (5.26) to reconstruct, as far as possible, the IR finite operator I_T . Again, we discard the terms that lead only to $\mathcal{O}(\epsilon^{-1})$ poles for the time being, and write the unboosted contribution as a sum of two terms

$$\Sigma_{\text{DU}, \mathcal{O}(\epsilon^{-2})}^{\text{ub}} = \Sigma_{\text{DU}}^{\text{ub}, s} + \Sigma_{\text{DU}}^{\text{ub}, c}. \quad (6.12)$$

The first term on the right-hand side of Eq. (6.12) contains I_T as well as left-over I_S and I_V operators (i.e., all color-correlated contributions), while the second term contains left-over terms arising from collinear limits, and hence no color-correlated matrix elements are

⁷We also need to include the double-virtual contribution and the soft limit of the real-virtual correction, the latter of which we borrow from Ref. [1].

present there. The expression for the first term reads

$$\begin{aligned}
\Sigma_{\text{DU}}^{\text{ub},s} &= [\alpha_s]^2 \left\langle \left[\frac{1}{2} (I_{\text{T}}^{(0)})^2 + K I_{\text{T}}^{(0)} + \beta_0 I_{\text{T}}^{(1)} \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\
&+ [\alpha_s]^2 \frac{\beta_0}{\epsilon} \left\langle \left[(\tilde{c}_\epsilon - 1) I_V(2\epsilon) + c_2(\epsilon) \tilde{I}_S(2\epsilon) - I_S(2\epsilon) \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\
&+ 2^{1+2\epsilon} [\alpha_s]^2 \delta(\epsilon) \left\langle \left[(2E_{\text{max}}/\mu)^{-2\epsilon} \left(-I_S(\epsilon) + \frac{(2E_{\text{max}}/\mu)^{-2\epsilon}}{2\epsilon^2} N_c(\epsilon) \sum_{i \in \mathcal{H}} \mathbf{T}_i^2 \right) \right. \right. \\
&\left. \left. + 2I_S(2\epsilon) \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle + [\alpha_s] \left[C_A \left(\frac{c_1(\epsilon)}{\epsilon^2} - \frac{A_K(\epsilon)}{\epsilon^2} \right) - \tilde{c}_\epsilon K + \beta_0 c_3(\epsilon) - \frac{2}{3} n_f T_{\text{R}} c_4(\epsilon) \right. \\
&\left. - 2^{2+2\epsilon} \delta(\epsilon) \right] \left\langle I_S(2\epsilon) \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle, \tag{6.13}
\end{aligned}$$

where $\delta(\epsilon) = \delta_g(\epsilon) + 2n_f \delta_q(\epsilon)$ (cf. Eqs (A.22) and (A.23)), and the remaining coefficients are collected in Appendix A.1. The first line of Eq. (6.13) is manifestly finite, with the first term having the expected form for the exponentiation given in Eq. (6.1). In the second line, we can use the expansions of \tilde{c}_ϵ and c_2 , as well as the relation

$$I_S(\epsilon) - \tilde{I}_S(\epsilon) = \mathcal{O}(\epsilon), \tag{6.14}$$

to demonstrate that only $\mathcal{O}(\epsilon^{-1})$ poles are present, and that these are proportional to the residues of the double poles of I_S and I_V (i.e., sums over Casimir operators), and hence are not color-correlated. In the third line, the combination of $I_S(\epsilon)$, $I_S(2\epsilon)$, and the sum over the Casimirs cancels both the $1/\epsilon^2$ and the $1/\epsilon$ poles, leaving a finite result. Finally, it is simple to check that the terms in brackets in the final two lines are $\mathcal{O}(\epsilon)$, and this produces a color-uncorrelated simple pole when multiplied by the operator $I_S(2\epsilon)$. We therefore conclude that the expression in Eq. (6.13) has only $\mathcal{O}(\epsilon^{-1})$ singularities, with all color-correlated terms being ϵ -finite.

We now move on to the unboosted component arising from collinear limits. It reads

$$\begin{aligned}
\Sigma_{\text{DU}}^{\text{ub},c} &= \frac{[\alpha_s]^2}{\epsilon^2} h_c(\epsilon) C_A \left\langle \left[I_{\text{C}}^{(4)}(\epsilon) - \tilde{I}_{\text{C}}(2\epsilon) \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle + \frac{[\alpha_s]^2}{\epsilon^2} \sum_{i \in \mathcal{H}_{\text{f}}} \left\langle \left\{ h_c(\epsilon) \mathcal{X}_{i,f_i} \right. \right. \\
&+ \frac{1}{2} \left[G_i \Big|_{z,f_i \rightarrow f_i g}^{f_i} + c_{f_i} G_i \Big|_{z,f_i \rightarrow \tilde{f}_i q}^{\tilde{f}_i} + c_{f_i} \Gamma_{i,f_i \rightarrow \tilde{f}_i q}(\Gamma_{i,\tilde{f}_i} - \Gamma_{i,f_i}) \right] \\
&\left. \left. + \frac{N_{\text{sc}}^{(b,d)}}{2} \left[\gamma_q^{22}(L_i) \Gamma_{i,f_i \rightarrow \tilde{f}_i q}^{(4)} + H_{i,f_i \rightarrow \tilde{f}_i q} + \gamma_g^{22}(0) \Gamma_{i,f_i \rightarrow f_i g}^{(4)} \right] - \frac{\beta_0}{2} \Gamma_{i,f_i}(2\epsilon) \right\} \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle, \tag{6.15}
\end{aligned}$$

where the H -functions are defined in Eq. (5.28). Moreover, f_i labels the flavour of parton $i \in \mathcal{H}_{\text{f}}$, while \tilde{f}_i labels the “complementary flavor” of f_i . Therefore, if parton i is a gluon, we have $f_i = g$ and $\tilde{f}_i = q$, while if parton i is a quark, then $f_i = q$ and $\tilde{f}_i = g$. In Eq. (6.15), c_{f_i} and \mathcal{X}_{i,f_i} are defined as

$$\begin{aligned}
c_{f_i} &= \begin{cases} 2n_{\text{f}}, & f_i = g, \\ 1, & f_i = q, \end{cases} \\
\mathcal{X}_{i,f_i} &= \begin{cases} (2C_{\text{F}} - C_{\text{A}}) \sigma_{i,g \rightarrow q\bar{q}} + 4n_{\text{f}}(C_{\text{F}} - C_{\text{A}}) \left(\frac{\Gamma_{i,g \rightarrow q\bar{q}}^{(4)}}{2\epsilon} - \frac{\Gamma_{i,g \rightarrow q\bar{q}}^{1\text{L}}}{2\epsilon} \right), & f_i = g, \\ C_{\text{A}} \sigma_{i,q \rightarrow gq}, & f_i = q. \end{cases} \tag{6.16}
\end{aligned}$$

Furthermore, the σ -functions are defined in Eq. (5.8), $N_{\text{sc}}^{(b,d)}$ and $h_c(\epsilon)$ in Eqs (A.7), and (A.10), respectively, the G -functions in Eqs (5.21), (5.22) and (5.23), and the H -functions in Eq. (5.28).

Eq. (6.15) has two important features. The first one is that it is given by the sum of contributions over external final-state partons, which suggests a straightforward generalization for even more complex processes than considered here. The second is that the contributions from each leg can be written using the universal quantities c_{f_i} and \mathcal{X}_{i,f_i} . As we will show below, these features allow for a simple discussion of the cancellation of $1/\epsilon$ poles, at least through $\mathcal{O}(\epsilon^{-2})$, which is quite remarkable given the complexity of the process considered.

The cancellation of the $1/\epsilon^3$ pole takes place in a straightforward way. The only term of order $\mathcal{O}(\epsilon^{-3})$ is the first on the right-hand side of Eq. (6.15), since both collinear operators are $\mathcal{O}(\epsilon^{-1})$ (cf. Eq. (A.41)), while all the remaining quantities are $\mathcal{O}(\epsilon^0)$. However, $I_C^{(4)}$ and \tilde{I}_C are related by the equation

$$I_C^{(4)}(\epsilon) = \tilde{I}_C(2\epsilon) + \mathcal{O}(\epsilon^0), \quad (6.17)$$

which immediately ensures the cancellation of the $1/\epsilon^3$ pole.

We move to the $1/\epsilon^2$ pole, distinguishing the case of initial-state radiation from that of final-state radiation. We begin with the former, which is contained in the first term on the right-hand side of Eq. (6.15), since the second and third lines pertain to final-state radiation. The objects describing the contribution of the a and b legs in the operators $I_C^{(4)}$ and \tilde{I}_C correspond to the generalized initial-state anomalous dimensions $\Gamma_{i,f_i}^{(4)}$ and $\Gamma_{i,f_i}^{1\text{L}}$, defined in Eqs (A.17) and (A.30), respectively. Since we have

$$\frac{\Gamma_{i,f_i}^{(4)}}{2\epsilon} = \frac{\Gamma_{i,f_i}^{1\text{L}}}{2\epsilon} + \mathcal{O}(\epsilon), \quad i \in \{a, b\}, \quad (6.18)$$

and also $h_c(\epsilon) = 1 + \mathcal{O}(\epsilon^3)$, we conclude that the $1/\epsilon^2$ pole in Eq. (6.15) vanishes for $i \in \{a, b\}$.

Next, we consider the final-state radiation, that is, $i \in \mathcal{H}_f$, where the second and third lines in Eq. (6.15) also contribute. The cancellation mechanism for the final-state gluon and quark legs is entirely analogous. In this case, the cancellation of the $1/\epsilon^2$ pole does not occur immediately by inspection, but still follows from elementary algebraic manipulations. To see this, it is sufficient to expand the anomalous dimensions in Eq. (5.27) and Eq. (A.20) in powers of ϵ , obtaining

$$\gamma_x^{nk}(L_i) = \sum_{m=0}^{\infty} \gamma_x^{nk,(m)}(L_i) \epsilon^m, \quad L_i = \log\left(\frac{E_{\text{max}}}{E_i}\right). \quad (6.19)$$

Furthermore, a similar expansion for the one-loop anomalous dimension contained in $\Gamma_{i,f_i}^{1\text{L}}$ (cf. Eqs (A.33 – A.34)) reads

$$\gamma_x^{1\text{L}}(L_i) = -\frac{\epsilon^2 \cos(\pi\epsilon)}{C_x} \gamma_x^{3(k+1),1\text{L}}(L_i) = \sum_{m=0}^{\infty} \gamma_x^{1\text{L},(m)}(L_i) \epsilon^m, \quad (6.20)$$

where $C_x = 2C_F - C_A$ if $x = (g \rightarrow q\bar{q})$ and $C_x = C_A$ otherwise. We then observe that only the coefficients with $m = 0, 1$ contribute to the cancellation of the poles. For instance, in the case of the quark leg, we can use the expansions

$$\begin{aligned}\Gamma_{i,x}^{(k)} &= \gamma_x^{2k,(0)}(L_i) + \epsilon \left(\gamma_x^{2k,(1)}(L_i) - k \tilde{L}_i \gamma_x^{2k,(0)}(L_i) \right) + \mathcal{O}(\epsilon^2), \\ \Gamma_{i,f_i}^{1L} &= \gamma_{f_i}^{1L,(0)}(L_i) + \epsilon \left(\gamma_{f_i}^{1L,(1)}(L_i) - 4 \tilde{L}_i \gamma_{f_i}^{1L,(0)}(L_i) \right) + \mathcal{O}(\epsilon^2),\end{aligned}\tag{6.21}$$

and noticing that $\gamma_x^{2k,(0)}(L_i) \equiv \gamma_{f_i}^{1L,(0)}(L_i) = \gamma_i + 2\mathbf{T}_i^2 L_i$, obtain

$$h_c(\epsilon) C_A \left(\frac{\Gamma_{i,q}^{(4)}}{2\epsilon} - \frac{\Gamma_{i,q}^{1L}}{2\epsilon} \right) = \frac{C_A}{2} \left[\gamma_q^{24,(1)}(L_i) - \gamma_q^{1L,(1)}(L_i) \right] + \mathcal{O}(\epsilon),\tag{6.22}$$

where $\tilde{L}_i = \log(2E_i/\mu)$. Similarly, we expand the remaining objects on the right-hand side of Eq. (6.15), and we find

$$\begin{aligned}\mathcal{X}_{i,q} &= \frac{C_A}{2} \left[\gamma_{z,q \rightarrow gq}^{22,(1)} - \gamma_{z,q \rightarrow gq}^{24,(1)} - 2L_i \gamma_{z,q \rightarrow gq}^{22,(0)} \right] + \mathcal{O}(\epsilon), \\ G_i \Big|_{z,f_i \rightarrow f_i g}^{f_i} &= \mathbf{T}_{f_i}^2 \left[\gamma_{z,f_i \rightarrow f_i g}^{42,(1)}(L_i) - \gamma_{z,f_i \rightarrow f_i g}^{22,(1)}(L_i) \right] + \mathcal{O}(\epsilon), \\ H_{i,f_i \rightarrow \tilde{f}_i q} &= C_F \left[\gamma_{z,q \rightarrow gq}^{24,(1)} - \gamma_{z,q \rightarrow gq}^{22,(1)} \right] + \mathcal{O}(\epsilon),\end{aligned}\tag{6.23}$$

while all remaining quantities can be approximated as $\gamma_x^{2k,(0)}$. Thus, Eq. (6.15) reduces to a simple expression involving coefficients $\gamma_x^{2k,(m)}$, $\gamma_x^{42,(m)}$, and $\gamma_x^{1L,(m)}$ with $m = 0, 1$, which vanishes at $\mathcal{O}(\epsilon^{-2})$. We observe that the same argument can be repeated for the gluon leg.

Before concluding this section, we briefly discuss the simple poles present in the single-boosted and unboosted double-unresolved contributions, which have been omitted from the preceding discussions. Similar to the all-gluon case in Ref. [1], their cancellation is difficult to trace by eye, but is entirely straightforward to achieve using computer algebra. We stress that our ability to absorb the many $1/\epsilon$ poles into finite structures, such as I_T , simplifies the analysis of the simple poles dramatically, since it reduces the bulk of it to a discussion of *independent* pole cancellations for each of the external legs.

To cancel the simple poles, we require the integrated triple-collinear subtraction terms, which we take from Ref. [93]. However, we need to slightly modify the results of this reference for two reasons. The first is to take into account the damping factors, which in Ref. [1] were simplified by the symmetry of the final state. The second is that the calculation in Ref. [93] assumed that all potentially-unresolved final-state partons are energy-ordered, whereas we impose this ordering only for final states that admit a singular double-soft limit, and do not require it for e.g. $(\mathbf{m}_q \mathbf{n}_g)$. The changes are relatively minor, as they affect only the strongly-ordered collinear limits. We include updated expressions for the integrated triple-collinear subtraction terms in the ancillary file `UsefulFunctions.m` (see the last four rows of Table 2).

Upon making these changes and including all sources of simple poles, we are able to confirm the cancellation of all poles, and hence obtain a finite remainder which we present in the following section.

7 Finite remainder of the NNLO contribution to the cross section

The goal of this section is to present the expression for $d\hat{\sigma}_{\mathcal{A}_0}^{\text{NNLO}}$ in Eq. (4.2) which is free of infrared singularities and, therefore, can be computed numerically. The main motivations for showing such a result are its compactness and iterative structure, as well as its proximity to the all-gluon case presented in Ref. [1]. These features, together with the rather general nature of the discussion in the preceding sections, give us confidence that a similar expression for *arbitrary* processes at hadron colliders can be derived in the near future.

To present the result, we split $d\hat{\sigma}_{\text{NNLO}}$ into fully-resolved, single-unresolved, and double-unresolved contributions, and write

$$2s_{ab} d\hat{\sigma}_{\text{NNLO}}^{\mathcal{A}_0} = 2s_{ab} [d\hat{\sigma}_{\text{FR}}^{\mathcal{A}_0} + d\hat{\sigma}_{\text{SU}}^{\mathcal{A}_0} + d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0}] . \quad (7.1)$$

The first term on the right-hand side of (7.1) is the finite, fully-regulated contribution. It corresponds to

$$2s_{ab} d\hat{\sigma}_{\text{FR}}^{\mathcal{A}_0} = \sum_{n=4}^6 \sum_{(\text{mn})} \mathcal{P}_{\mathcal{A}_0} \Sigma_{\text{FR}}^{\mathcal{A}_n}[(\text{mn})]_{\text{RR}} , \quad (7.2)$$

where the quantity $\Sigma_{\text{FR}}^{\mathcal{A}_n}[(\text{mn})]_{\text{RR}}$ is defined in Eq. (4.11).

The second term on the right-hand side of Eq. (7.1) is the finite remainder of the single-unresolved contribution. We write it as the sum of four contributions

$$2s_{ab} d\hat{\sigma}_{\text{SU}}^{\mathcal{A}_0} = \sum_{i=1}^4 \mathcal{P}_{\mathcal{A}_0} \Sigma_{\text{SU}}^{\text{fin},(i)}|_{\mathcal{A}_0} , \quad (7.3)$$

where

$$\begin{aligned} \Sigma_{\text{SU}}^{\text{fin},(1)}|_{\mathcal{A}_0} &= [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(a,\text{m})} \omega_{a||n}^{\text{ma,na}} \log\left(\frac{\eta_{am}}{2}\right) \Delta^{(\text{m})} [\hat{P}_{gg}^{(0)} \otimes F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}]] \right\rangle \\ &+ [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(b,\text{m})} \omega_{b||n}^{\text{mb,nb}} \log\left(\frac{\eta_{bm}}{2}\right) \Delta^{(\text{m})} [F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \otimes \hat{P}_{qq}^{(0)}] \right\rangle - \sum_{i \in \mathcal{H}} [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(i,\text{m})} \omega_{i||n}^{\text{mi,ni}} \right. \\ &\times \log\left(\frac{\eta_{im}}{2}\right) \Delta^{(\text{m})} \left[(\gamma_i + 2\mathbf{T}_i^2 L_i(E_{\text{m}})) F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] + (\gamma_i + 2\mathbf{T}_i^2 L_i) F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \right] \left. \right\rangle , \end{aligned} \quad (7.4)$$

$$\begin{aligned} \Sigma_{\text{SU}}^{\text{fin},(2)}|_{\mathcal{A}_0} &= - \sum_{i \in \mathcal{H}} [\alpha_s] \left\langle \mathcal{O}_{\text{NLO}}^{(i,\text{m})} \omega_{\text{m}||n}^{\text{mi,ni}} \log\left(\frac{\eta_{im}}{4(1-\eta_{im})}\right) \Delta^{(\text{m})} \left[\gamma_g F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \right. \right. \\ &\left. \left. + (\gamma_q + 2L_{\text{m}} \mathbf{T}_q^2) F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q] \right] \right\rangle , \end{aligned} \quad (7.5)$$

$$\begin{aligned} \Sigma_{\text{SU}}^{\text{fin},(3)}|_{\mathcal{A}_0} &= \sum_{i \in \mathcal{H}} \frac{[\alpha_s]}{2} \gamma_{\perp,g}^{22} \left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \omega_{\text{m}||n}^{\text{mi,ni}} (r_i^\mu r_i^\nu + g^{\mu\nu}) \Delta^{(\text{m})} F_{\text{LM},\mu\nu}^{\mathcal{A}_1}[\mathbf{m}_g] \right\rangle \\ &+ \sum_{i \in \mathcal{H}} \frac{[\alpha_s]}{2} \gamma_{\perp,g}^{22,\text{r}} \left\langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \omega_{\text{m}||n}^{\text{mi,ni}} \Delta^{(\text{m})} F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] \right\rangle , \end{aligned} \quad (7.6)$$

$$\begin{aligned}
\Sigma_{\text{SU}}^{\text{fin},(4)}|_{\mathcal{A}_0} &= \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} (F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}_g] + F_{\text{RV,fin}}^{\mathcal{A}_1}[\mathbf{m}_q]) \rangle \\
&+ [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} [I_{\text{T}}^{(0)}(E_{\text{m}}) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_g] + I_{\text{T}}^{(0)}(E_{\text{max}}) \cdot F_{\text{LM}}^{\mathcal{A}_1}[\mathbf{m}_q]] \rangle \\
&+ [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} [\mathcal{P}_{gg}^{\text{NLO}} \otimes F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}]] \rangle + [\alpha_s] \langle \mathcal{O}_{\text{NLO}}^{(\text{m})} \Delta^{(\text{m})} [F_{\text{R}}^{\mathcal{A}_1}[\mathbf{m}] \otimes \mathcal{P}_{qq}^{\text{NLO}}] \rangle.
\end{aligned} \tag{7.7}$$

The definitions of quantities that appear in the above formulas can be found in Eqs (4.14) and (4.15), as well as in Sec. 6.1. Here, we note that the objects $\gamma_{\perp,g}^{22}$ and $\gamma_{\perp,g}^{22,r}$ are given in Eq. (A.27), and their expressions are reported in the ancillary file `FinalResult.m` provided with this paper, see Table 1. The quantity $I_{\text{T}}^{(0)}$ is the $\mathcal{O}(\epsilon^0)$ expansion coefficient of the IR-finite operator $I_{\text{T}}(\epsilon)$; it can be found in Eq. (A.45). The splitting functions $\hat{P}_{\alpha\alpha}^{(0)}$ are defined in Eq. (A.13), while the splitting functions $\mathcal{P}_{\alpha\alpha}^{\text{NLO}}$ are reported in `FinalResult.m`, see Table 1. Finally, we remind the reader that the argument E_{m} in $L_i(E_{\text{m}})$ and $I_{\text{T}}^{(0)}(E_{\text{m}})$ indicates that E_{max} must be replaced with E_{m} .

The third term on the right-hand side of Eq. (7.1) corresponds to the finite remainder of the double-unresolved contribution. It is convenient to write it as the sum of the following quantities

$$d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0} = d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{db}} + d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{sb},a} + d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{sb},b} + d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{el}}, \tag{7.8}$$

where $d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{db}}$ contains all the *double-boosted* contributions, $d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{sb},a}$ and $d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{sb},b}$ contain all the contributions that are *single-boosted* on leg a and leg b , respectively, and $d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{el}}$ contains all the *elastic* contributions with the kinematics of the Born process. We present each contribution separately, using several functions that we collect in `FinalResult.m`.

The double-boosted contribution is described by the very simple expression (see Eq. (6.6))

$$2s_{ab} d\hat{\sigma}_{\mathcal{A}_0,\text{db}}^{\text{DU}} = [\alpha_s]^2 \langle \mathcal{P}_{gg}^{\text{NLO}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{NLO}} \rangle. \tag{7.9}$$

Expressions for the single-boosted contributions, related to the emissions off the incoming partons a and b , are slightly more complex. We write them as

$$\begin{aligned}
2s_{ab} d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{sb},a} &= [\alpha_s]^2 \left[\langle \mathcal{P}_{gg}^{\text{NLO}} \otimes [I_{\text{T}}^{(0)} \cdot F_{\text{LM}}^{\mathcal{A}_0}] \rangle + \langle \mathcal{P}_{gg}^{\mathcal{W}} \otimes [\mathcal{W}_a^{a||\text{n,fin}} \cdot F_{\text{LM}}^{\mathcal{A}_0}] \rangle \right. \\
&\quad \left. + \langle \mathcal{P}_{gg}^{\text{NNLO}} \otimes F_{\text{LM}}^{\mathcal{A}_0} \rangle + \langle \mathcal{P}_{gg}^{\text{NLO}} \otimes F_{\text{LV,fin}}^{\mathcal{A}_0} \rangle \right],
\end{aligned} \tag{7.10}$$

and

$$\begin{aligned}
2s_{ab} d\hat{\sigma}_{\text{DU}}^{\mathcal{A}_0,\text{sb},b} &= [\alpha_s]^2 \left[\langle [I_{\text{T}}^{(0)} \cdot F_{\text{LM}}^{\mathcal{A}_0}] \otimes \mathcal{P}_{qq}^{\text{NLO}} \rangle + \langle [\mathcal{W}_b^{b||\text{n,fin}} \cdot F_{\text{LM}}^{\mathcal{A}_0}] \otimes \mathcal{P}_{qq}^{\mathcal{W}} \rangle \right. \\
&\quad \left. + \langle F_{\text{LM}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{NNLO}} \rangle + \langle F_{\text{LV,fin}}^{\mathcal{A}_0} \otimes \mathcal{P}_{qq}^{\text{NLO}} \rangle \right].
\end{aligned} \tag{7.11}$$

The partition-dependent operators $\mathcal{W}_i^{i||\text{n,fin}}$, with $i = a, b$, are given below in Eq. (7.15). Furthermore, $\mathcal{P}_{\alpha\alpha}^{\mathcal{W}}$ and the NNLO splitting functions $\mathcal{P}_{\alpha\alpha}^{\text{NNLO}}$ are also reported in the ancillary file `FinalResult.m` (cf. Table 1).

Table 1: List of functions collected in the ancillary file `FinalResult.m`. The first column provides the names of the functions; the second specifies the index range i, j ; the third indicates the equation where they appear; the fourth provides their names in `FinalResult.m`. For brevity, in the second block of the table, \mathcal{P}_{gg}^{\dots} and \mathcal{P}_{qq}^{\dots} are encoded in \mathcal{P}_{xx}^{\dots} . To extract the gg or qq splittings, call `Pgg...[z,En]` or `Pqq...[z,En]`.

Functions collected in the ancillary file <code>FinalResult.m</code>			
<i>Function</i>	<i>Interval of i, j</i>	<i>Ref.</i>	<i>Name in the ancillary file</i>
<i>Integrals in spin-correlations</i>			
$\gamma_{\perp,g}^{22}$	/	Eq. (A.27)	<code>ygPerp</code>
$\gamma_{\perp,g}^{22,r}$	/	Eq. (A.27)	<code>ygPerpR</code>
$\delta^{(0)}$	/	Eq. (7.12)	<code>deltazero</code>
$\delta^{\perp,(0)}$	/	Eq. (7.12)	<code>deltaPerpzero</code>
<i>Splitting functions</i>			
$\mathcal{P}_{xx}^{\text{NLO}}(z, E_i)$	$i \in \{a, b\}$	Eqs (7.7, 7.9 – 7.11)	<code>PxxNLO[z,En]</code>
$\mathcal{P}_{xx}^{\mathcal{W}}(z, E_i)$	$i \in \{a, b\}$	Eqs (7.10, 7.11)	<code>PxxW[z,En]</code>
$\mathcal{P}_{xx}^{\text{NNLO}}(z, E_i)$	$i \in \{a, b\}$	Eqs (7.10, 7.11)	<code>PxxNNLO[z,En]</code>
<i>Elastic functions</i>			
$\gamma_{z,g \rightarrow gg}^{\mathcal{W}}(L_i)$	$i \in \mathcal{H}_{f,g}$	Eq. (7.12)	<code>gammaWgTOgg[En]</code>
$\gamma_{z,q \rightarrow qq}^{\mathcal{W}}(L_i)$	$i \in \mathcal{H}_{f,g}$	Eq. (7.12)	<code>gammaWqTOqq[En]</code>
$D_g^{\text{ISR}}(E_i)$	$i \in \{a, b\}$	Eq. (7.14)	<code>DgISR[En]</code>
$D_q^{\text{ISR}}(E_i)$	$i \in \{a, b\}$	Eq. (7.14)	<code>DqISR[En]</code>
$D_g^{\text{FSR}}(E_i)$	$i \in \mathcal{H}_{f,g}$	Eq. (7.14)	<code>DgFSR[En]</code>
$D_q^{\text{FSR}}(E_i)$	$i \in \mathcal{H}_{f,q}$	Eq. (7.14)	<code>DqFSR[En]</code>
<i>Double-soft finite remainders</i>			
$(S_{gg,T^2}^{\text{fin}})_{ij}$	$i, j \in \mathcal{H}, i \neq j$	Eq. (7.12)	<code>SggT2fin[i,j]</code>
$(S_{q\bar{q},T^2}^{\text{fin}})_{ij}$	$i, j \in \mathcal{H}, i \neq j$	Eq. (7.12)	<code>SqqbT2fin[i,j]</code>

Finally, the elastic contribution reads

$$\begin{aligned}
2s_{ab} \, d\hat{\sigma}_{\mathcal{A}_0, \text{el}}^{\text{DU}} &= [\alpha_s]^2 \langle [I_{\text{cc}}^{\text{fin}} + I_{\text{tri}}^{\text{fin}} + I_{\text{unc}}^{\text{fin}}] \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle \\
&+ [\alpha_s]^2 \sum_{i \in \mathcal{H}} \left\langle \left[\theta_{\mathcal{H}_f} \gamma_{z, f_i \rightarrow f_i g}^{\mathcal{W}} \mathcal{W}_i^{i \| n, \text{fin}} + \delta^{(0)} \mathcal{W}_i^{\text{m} \| n, \text{fin}} + \delta^{\perp, (0)} \mathcal{W}_i^{(i)} \right] \cdot F_{\text{LM}}^{\mathcal{A}_0} \right\rangle \\
&+ [\alpha_s]^2 \sum_{(ij)} \langle [(S_{gg, T^2}^{\text{fin}})_{ij} + n_f (S_{q\bar{q}, T^2}^{\text{fin}})_{ij}] (\mathbf{T}_i \cdot \mathbf{T}_j) \cdot F_{\text{LM}}^{\mathcal{A}_0} \rangle \\
&+ [\alpha_s] \langle I_{\text{T}}^{(0)} \cdot F_{\text{LV}, \text{fin}}^{\mathcal{A}_0} \rangle + \langle F_{\text{LV}^2, \text{fin}}^{\mathcal{A}_0} \rangle + \langle F_{\text{VV}, \text{fin}}^{\mathcal{A}_0} \rangle.
\end{aligned} \tag{7.12}$$

In the first line of Eq. (7.12), we have the combination of quartic and double color-correlated contributions in $I_{\text{cc}}^{\text{fin}}$, a triple color-correlated component $I_{\text{tri}}^{\text{fin}}$, and a color-uncorrelated part $I_{\text{unc}}^{\text{fin}}$. The first of these is particularly simple, and reads

$$\begin{aligned}
I_{\text{cc}}^{\text{fin}} &= \frac{1}{2} (I_{\text{T}}^{(0)})^2 + K I_{\text{T}}^{(0)} + \beta_0 \left[I_{\text{T}}^{(1)} + \tilde{I}_{\text{S}}^{(1)} - 2I_{\text{S}}^{(1)} + \frac{\pi^2}{24} I_{\text{V}}^{(-1)} + 2I_{\text{S}}^{(0)} \log 2 \right. \\
&+ \left. I_{\text{S}}^{(-1)} \left(4L_{\text{max}} \log 2 - 6 \log^2 2 - \frac{\pi^2}{2} \right) \right] + C_{\text{A}} \left[I_{\text{S}}^{(-1)} \left(\frac{1975}{108} - \frac{17\zeta_3}{4} - \frac{2}{3} \pi^2 \log 2 \right) \right. \\
&+ \left. \left(I_{\text{S}}^{(0)} + 2L_{\text{max}} I_{\text{S}}^{(-1)} \right) \left(\frac{\pi^2}{3} - \frac{131}{36} \right) \right] + n_f T_{\text{R}} \left[\frac{23}{18} \left(I_{\text{S}}^{(0)} + 2L_{\text{max}} I_{\text{S}}^{(-1)} \right) - \frac{331}{54} I_{\text{S}}^{(-1)} \right],
\end{aligned} \tag{7.13}$$

where K is defined in Eq. (A.8), and $I_{\text{S}}^{(n)}$, $\tilde{I}_{\text{S}}^{(n)}$, $I_{\text{V}}^{(n)}$, $I_{\text{T}}^{(n)}$ are the coefficients of the n -th power in the ϵ -expansion of the corresponding operators (cf. Appendix A.4). We note that Eq. (7.13) generalizes Eq. (I.8) in Ref. [1], which was computed for the process $q\bar{q} \rightarrow X + N_g g$ with $n_f = 0$, to the more general case of the \mathcal{A}_0 process in Eq. (2.1), where terms proportional to n_f have also been included. The definition of the operator $I_{\text{tri}}^{\text{fin}}$, which contains triple color-correlated components, remains identical to the one in Eq. (I.9) of Ref. [1]. It contains the most complicated functions appearing in the final result, namely classical and generalized polylogarithms (GPLs) up to weight three, whose arguments can be algebraic functions of the kinematic variables. The operator $I_{\text{unc}}^{\text{fin}}$ collects color-uncorrelated contributions. We write it as the sum of two terms, one that only depends on the color charges of each external parton, and another, named $D_{f_i}(E_i)$, whose functional form depends on the type of parton and whether it is in the initial or the final state. We write

$$\begin{aligned}
I_{\text{unc}}^{\text{fin}} &= \sum_{i \in \mathcal{H}} \mathbf{T}_i^2 \left\{ C_{\text{A}} \left[\left(\frac{2\pi^2}{3} - \frac{131}{18} + \frac{22}{3} \log 2 \right) L_{\text{max}}^2 - \frac{935}{72} \zeta_3 + \frac{9607}{324} - \frac{11}{3} \log^3 2 \right. \right. \\
&+ \left. \left(\frac{1433}{108} - 8\zeta_3 - \frac{11\pi^2}{6} \right) \log 2 - \frac{199\pi^4}{1440} - \frac{21\pi^2}{32} + \left(\frac{143}{36} - \frac{\pi^2}{3} \right) \log^2 2 \right] \\
&+ n_f T_{\text{R}} \left[\left(\frac{23}{9} - \frac{8}{3} \log 2 \right) L_{\text{max}}^2 + \frac{85}{18} \zeta_3 - \frac{746}{81} + \frac{4}{3} \log^3 2 + \left(\frac{2\pi^2}{3} - \frac{67}{27} \right) \log 2 \right. \\
&- \left. \left. \frac{47}{18} \log^2 2 + \frac{5\pi^2}{72} \right] \right\} + \sum_{i \in \mathcal{H}} D_{f_i}(E_i),
\end{aligned} \tag{7.14}$$

where the functions D_{f_i} are reported in the ancillary file `FinalResult.m`, see Table 1.⁸ The function $I_{\text{unc}}^{\text{fin}}$ in Eq. (7.14) is a generalization of a similar quantity in Ref. [1] (see Eq. (I.12) therein).

In the second line of Eq. (7.12), we have $\theta_{\mathcal{H}_f} = 1$ if $i \in \mathcal{H}_f$ and $\theta_{\mathcal{H}_f} = 0$ otherwise. The quantities $\delta^{(0)}$, $\delta^{\perp,(0)}$, and $\gamma_{z,f_i \rightarrow f_i g}^{\mathcal{W}}$ reported in `FinalResult.m`, while the partition-dependent operators $\mathcal{W}_i^{i||n,\text{fin}}$, $\mathcal{W}_i^{\text{m}||n,\text{fin}}$, $\mathcal{W}_r^{(i)}$ are defined as⁹

$$\begin{aligned}\mathcal{W}_i^{i||n,\text{fin}} &= \sum_{(kl)} \int \frac{d\Omega_{\text{m}}^{(3)}}{2\pi} \bar{C}_{im} \left[\log(\eta_{im}/2) \frac{\rho_{kl}}{\rho_{km}\rho_{lm}} \omega_{i||n}^{\text{mi},ni} \right] (\mathbf{T}_k \cdot \mathbf{T}_l), \\ \mathcal{W}_i^{\text{m}||n,\text{fin}} &= \sum_{(kl)} \int \frac{d\Omega_{\text{m}}^{(3)}}{2\pi} \bar{C}_{im} \left[\log\left(\frac{\eta_{im}}{1-\eta_{im}}\right) \frac{\rho_{kl}}{\rho_{km}\rho_{lm}} \omega_{\text{m}||n}^{\text{mi},ni} \right] (\mathbf{T}_k \cdot \mathbf{T}_l), \\ \mathcal{W}_r^{(i)} &= \sum_{k,l \in \mathcal{H}} \int \frac{d\Omega_{\text{m}}^{(3)}}{2\pi} \omega_{\text{m}||n}^{\text{mi},ni} (r_i^\mu r_i^\nu + g^{\mu\nu}) \frac{n_{k,\mu} n_{l,\nu}}{(n_k \cdot n_{\text{m}})(n_l \cdot n_{\text{m}})} (\mathbf{T}_k \cdot \mathbf{T}_l),\end{aligned}\tag{7.15}$$

where a light-like vector n_i^μ is defined through the equation $p_i^\mu = E_i n_i^\mu$, and (kl) means $k, l \in \mathcal{H}$, with $k \neq l$. The term in the third line of Eq. (7.12) refers to the finite remainder of the double-soft integrated subtraction term (cf. Eq. (5.1)), and it can be extracted from Ref. [90]. The explicit expressions of $(S_{gg,T^2}^{\text{fin}})_{ij}$ and $(S_{q\bar{q},T^2}^{\text{fin}})_{ij}$, which are functions of η_{ij} , are given in `FinalResult.m`, see Table 1. Finally, $F_{\text{LV}^2,\text{fin}}^{\mathcal{A}_0}$ and $F_{\text{VV},\text{fin}}^{\mathcal{A}_0}$ are the process-dependent finite remainders of the one-loop squared and two-loop virtual amplitudes, respectively.

8 Conclusions

In this paper we discussed the calculation of the integrated NNLO QCD subtraction terms for the process $gq \rightarrow X + N_g g + q$, where the number of hard gluons N_g is a *parameter* and X is an arbitrary colorless final state. We worked in the context of the nested soft-collinear subtraction scheme [17] and followed closely a similar study of the all-gluon final state reported in Ref. [1].

We have found that the approach described in Ref. [1] is sufficiently robust, and can be applied to a complex final state that contains both quarks and gluons. In doing so, we have seen that physical quantities such as collinear anomalous dimensions arise when different singular configurations are properly combined before the subtraction terms are integrated over the unresolved phase space – a feature that is important for making computations not only more feasible but also physically transparent. Finally, since i) soft limits at NNLO depend only on the color charges of hard partons, ii) the collinear singularities factorize on the external parton legs and iii) an improved understanding of the interplay of the different collinear limits has been achieved in this paper, we believe that it should be straightforward to extend these results to processes with an arbitrary number of jets at a hadron collider.

⁸In Table 1, we label these functions as $D_{f_i}^{\text{ISR}}$ and $D_{f_i}^{\text{FSR}}$ if they refer to *initial-state radiation* or *final-state radiation*, respectively.

⁹We note that the cancellation of poles, discussed in the previous two sections, occurs independently of the exact expressions of the partition functions, provided they obey certain defining conditions, specified in e.g. Ref. [1].

Acknowledgments

We are grateful to Max Delto for discussions concerning integrated triple-collinear subtraction terms. The research of K.M. and D.M.T. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant no. 396021762 - TRR 257. The research of M.T. is supported by a grant from Deutscher Akademischer Austauschdienst (DAAD). F.D. and C.S.-S. wish to thank the CERN Theoretical Physics Department for hospitality while part of this work was carried out. R.R. is partially supported by the Italian Ministry of Universities and Research (MUR) through grant PRIN2022BCXSW9. The work of R.R. was performed in part at Aspen Center for Physics, which is supported by National Science Foundation grant PHY-2210452. This research was supported in part by grant NSF PHY-2309135 to the Kavli Institute for Theoretical Physics (KITP).

A Collection of relevant constant, functions and operators

In this Appendix, we collect and define quantities that are used in the main text of this paper. We supplement the material presented here with an ancillary file, `UsefulFunctions.m`, which contains the ϵ -series expansions of the functions listed in Table 2. To simplify various formulas, we use the following notation

$$\bar{z} = 1 - z, \quad \mathcal{D}_n(z) = \left[\frac{\log^n(1-z)}{1-z} \right]_+, \quad (A.1)$$

$$\tilde{L}_i = \log\left(\frac{2E_i}{\mu}\right), \quad L_i = \log\left(\frac{E_{\max}}{E_i}\right), \quad L_{\max} = \log\left(\frac{2E_{\max}}{\mu}\right).$$

Table 2: List of functions collected in the ancillary file `UsefulFunctions.m`. The first column provides the names of the functions; the second specifies the index range i ; the third indicates the equation where they appear; the fourth provides their names in `UsefulFunctions.m`. For brevity, \mathcal{P}_{qq} , \mathcal{P}_{qg} , \mathcal{P}_{gq} , and \mathcal{P}_{gg} are encoded in \mathcal{P}_{xy} . To extract a specific configuration – e.g., the fifth row – call `Pqq[z,k,En]`, `Pqg[z,k,En]`, etc. This convention applies to all splittings in the second and third blocks of the table. For final-state anomalous dimensions, replace `wTOxy` with `qTOqg`, `qTOgq`, `gTOqq`, or `gTOgg` to find the corresponding functions in `UsefulFunctions.m`. The last four functions refer to triple-collinear contributions; for further details, see the two paragraphs below Eq. (6.23).

Functions collected in the ancillary file <code>UsefulFunctions.m</code>			
<i>Function</i>	<i>Interval of i</i>	<i>Ref.</i>	<i>Name in the ancillary file</i>
<i>Integrals in spin-correlations</i>			
$\delta_g(\epsilon)$	/	Eq. (A.22)	<code>dg[ε]</code>
$\delta_q(\epsilon)$	/	Eq. (A.22)	<code>dq[ε]</code>
$\delta_g^\perp(\epsilon)$	/	Eq. (A.22)	<code>dgPerp[ε]</code>
$\delta_q^\perp(\epsilon)$	/	Eq. (A.22)	<code>dqPerp[ε]</code>
<i>Tree-level splitting functions</i>			
$\mathcal{P}_{xy}^{(k)}(z, E_i)$	$i \in \{a, b\}$	Eq. (A.15)	<code>Pxy[z,k,En]</code>
$\mathcal{P}_{xy}^{(k),\text{gen}}(z, E_i)$	$i \in \{a, b\}$	Eq. (A.18)	<code>PxyGEN[z,k,En]</code>
<i>One-loop splitting functions</i>			
$\hat{P}_{gg}^{(1)}(z)$	$i \in \{a, b\}$	/	<code>PAPggOneL[z]</code>
$\hat{P}_{qq,\tilde{\text{ns}}}^{(1)}(z)$	$i \in \{a, b\}$	/	<code>PAPqqOneLns[z]</code>
$P_{xy,i}^{\text{1L}}(z)$	$i \in \{a, b\}$	Eq. (A.28)	<code>PxyOneLISR[z]</code>

$P_{xy}^{1L}(z)$	$i \in \mathcal{H}_f$	Eq. (A.34)	PxyOneLFSR[z]
$\mathcal{P}_{xy}^{1L}(z, E_i)$	$i \in \{a, b\}$	Eq. (A.28)	PxyOneL[z, En]
$\mathcal{P}_{xy}^{1L, \text{gen}}(z, E_i)$	$i \in \{a, b\}$	Eq. (A.31)	PxyOneLGEN[z, En]
<i>Tree-level anomalous dimensions</i>			
$\Gamma_{i,g}^{(k)}$	$i \in \{a, b\}$	Eq. (A.17)	$\Gamma_{\text{gISR}}[k, \text{En}]$
$\Gamma_{i,q}^{(k)}$	$i \in \{a, b\}$	Eq. (A.17)	$\Gamma_{\text{qISR}}[k, \text{En}]$
$\gamma_{z,w \rightarrow xy}^{nk}(L_i)$	$i \in \mathcal{H}_f$	Eq. (A.20)	$\gamma_{\text{wT0xy}}[n, k, \text{En}]$
$\Gamma_{i,w \rightarrow xy}^{(k)}$	$i \in \mathcal{H}_f$	Eq. (A.19)	$\Gamma_{\text{wT0xy}}[k, \text{En}]$
$\Gamma_{i,g}^{(k)}$	$i \in \mathcal{H}_f$	Eq. (A.21)	$\Gamma_{\text{gFSR}}[k, \text{En}]$
$\Gamma_{i,q}^{(k)}$	$i \in \mathcal{H}_f$	Eq. (A.21)	$\Gamma_{\text{qFSR}}[k, \text{En}]$
<i>One-loop anomalous dimensions</i>			
$\Gamma_{i,g}^{1L}$	$i \in \{a, b\}$	Eq. (A.30)	$\Gamma_{\text{OneLgISR}}[\text{En}]$
$\Gamma_{i,q}^{1L}$	$i \in \{a, b\}$	Eq. (A.30)	$\Gamma_{\text{OneLqISR}}[\text{En}]$
$\gamma_{z,w \rightarrow xy}^{33,1L}(L_i)$	$i \in \mathcal{H}_f$	Eq. (A.34)	$\gamma_{\text{OneLwT0xy}}[\text{En}]$
$\Gamma_{i,w \rightarrow xy}^{1L}$	$i \in \mathcal{H}_f$	Eq. (A.33)	$\Gamma_{\text{OneLwT0xy}}[\text{En}]$
$\Gamma_{i,g}^{1L}$	$i \in \mathcal{H}_f$	Eq. (A.35)	$\Gamma_{\text{OneLgFSR}}[\text{En}]$
$\Gamma_{i,q}^{1L}$	$i \in \mathcal{H}_f$	Eq. (A.35)	$\Gamma_{\text{OneLqFSR}}[\text{En}]$
<i>Triple-collinear splitting functions</i>			
$\text{TC}_g^{\text{ISR}}(z, E_i)$	$i \in \{a, b\}$	/	$\text{TCgISR}[z, \text{En}]$
$\text{TC}_q^{\text{ISR}}(z, E_i)$	$i \in \{a, b\}$	/	$\text{TCqISR}[z, \text{En}]$
$\text{TC}_g^{\text{FSR}}(E_i)$	$i \in \mathcal{H}_{f,g}$	/	$\text{TCgFSR}[\text{En}]$
$\text{TC}_q^{\text{FSR}}(E_i)$	$i \in \mathcal{H}_{f,q}$	/	$\text{TCqFSR}[\text{En}]$

A.1 Useful constants

This section generalizes Appendix A.1 of Ref. [1] where many useful constants have been defined. We denote the color-charge operators with \mathbf{T}_i . Squares of these operators are the Casimir operators of the corresponding representations of the QCD gauge group SU(3), i.e.

$$\mathbf{T}_q^2 = \mathbf{T}_{\bar{q}}^2 = C_F = \frac{N_c^2 - 1}{2N_c}, \quad \mathbf{T}_g^2 = C_A = N_c, \quad (\text{A.2})$$

where N_c is the number of colors. Quark and gluon anomalous dimensions read

$$\gamma_q = \frac{3}{2}C_F, \quad \gamma_g = \frac{11}{6}C_A - \frac{2}{3}T_R n_f, \quad (\text{A.3})$$

where $T_R = 1/2$, and n_f is the number of massless quark flavors.

The strong coupling is renormalized in the $\overline{\text{MS}}$ scheme. The relation between the bare and renormalized coupling constants reads

$$g_{s,b}^2 = g_s^2 S_\epsilon \mu^{2\epsilon} \left[1 - \frac{\alpha_s(\mu)}{2\pi} \frac{\beta_0}{\epsilon} + \left(\frac{\alpha_s(\mu)}{2\pi} \right)^2 \left(\frac{\beta_0^2}{\epsilon^2} - \frac{\beta_1}{2\epsilon} \right) + \mathcal{O}(\alpha_s^3) \right], \quad (\text{A.4})$$

where $S_\epsilon = (4\pi)^{-\epsilon} e^{\epsilon\gamma_E}$, and

$$\beta_0 = \frac{11}{6} C_A - \frac{2}{3} T_R n_f, \quad \beta_1 = \frac{17}{6} C_A^2 - \frac{5}{3} C_A T_R n_f - C_F T_R n_f. \quad (\text{A.5})$$

We note that, at leading order, the gluon anomalous dimension and the coefficient of the QCD β -function coincide, $\gamma_g = \beta_0$. Furthermore, it is convenient to define the following quantity

$$[\alpha_s] \equiv \frac{\alpha_s(\mu)}{2\pi} \frac{e^{\epsilon\gamma_E}}{\Gamma(1-\epsilon)}, \quad (\text{A.6})$$

related to the coupling constant.

In the main body of this paper, we have used ϵ -dependent constants

$$\begin{aligned} N_\epsilon^{(b,d)} &= \frac{\Gamma(1-\epsilon)\Gamma(1+2\epsilon)}{\Gamma(1+\epsilon)} = 1 + \frac{\pi^2}{3}\epsilon^2 + \mathcal{O}(\epsilon^3), \\ N_{\text{m|n}}(\epsilon) &= 2^{2\epsilon} \frac{\Gamma(1+2\epsilon)\Gamma(1-2\epsilon)}{\Gamma(1+\epsilon)\Gamma(1-\epsilon)} = 1 + 2\epsilon \log 2 + \frac{1}{2}\epsilon^2(\pi^2 + 4\log^2 2) + \mathcal{O}(\epsilon^3), \\ N_c(\epsilon) &= -\frac{\Gamma(1-\epsilon)\Gamma(1-2\epsilon)}{\Gamma(1-3\epsilon)} + \frac{2\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} = 1 + \mathcal{O}(\epsilon^3), \\ N_{\text{sc}}^{(b,d)} &= 2^{2\epsilon} \frac{\Gamma(1+2\epsilon)\Gamma^3(1-2\epsilon)}{\Gamma(1-3\epsilon)\Gamma(1+\epsilon)\Gamma^2(1-\epsilon)} = 1 + 2\epsilon \log 2 + \frac{1}{3}\epsilon^2(\pi^2 + 6\log^2 2) + \mathcal{O}(\epsilon^3), \end{aligned} \quad (\text{A.7})$$

which originate from the definition of certain angular integrals.

Singular parts of double-virtual amplitudes, written in Ref. [87], require the following quantities

$$\begin{aligned} K &= \left(\frac{67}{18} - \frac{\pi^2}{6} \right) C_A - \frac{10}{9} T_R n_f, \\ c_\epsilon &= \frac{e^{-\epsilon\gamma_E} \Gamma(1-2\epsilon)}{\Gamma(1-\epsilon)} = 1 + \frac{\pi^2}{4}\epsilon^2 + \frac{7}{3}\zeta_3\epsilon^3 + \mathcal{O}(\epsilon^4), \\ \tilde{c}_\epsilon &= \frac{\Gamma(1-\epsilon)}{e^{\epsilon\gamma_E}} = 1 + \frac{\pi^2}{12}\epsilon^2 + \mathcal{O}(\epsilon^3). \end{aligned} \quad (\text{A.8})$$

To parametrize integrated double-soft contributions, we have introduced

$$\begin{aligned} c_1(\epsilon) &= 1 + \left(\frac{\pi^2}{6} - \frac{32}{9} \right) \epsilon^2 + \left(\frac{217}{27} - \frac{137}{9} \log 2 - 22 \log^2 2 + \frac{11\zeta_3}{2} \right) \epsilon^3, \\ c_2(\epsilon) &= 1 + \frac{\pi^2}{3} \epsilon^2, \\ c_3(\epsilon) &= 4 \log 2 + 8\epsilon \log^2 2, \\ c_4(\epsilon) &= -\frac{13}{6} + \left(\frac{125}{18} - \frac{35}{3} \log 2 - 12 \log^2 2 \right) \epsilon. \end{aligned} \quad (\text{A.9})$$

$$\begin{aligned}
\text{FSR : } C_{im} \Delta^{(m)} & \left| \begin{array}{c} \text{Diagram: A circle with multiple lines entering from the left. A red line labeled } p_m \sim \bar{z} p_{[im]} \text{ exits to the right. A red line labeled } p_i \text{ enters from the bottom. A red line labeled } p_{[im]} \text{ exits to the right.} \end{array} \right|^2 \xrightarrow{\theta_{im} \rightarrow 0} \frac{g_{s,b}^2 \mu_0^{2\epsilon}}{p_i \cdot p_m} z P_{f_{[im]} f_i}(z) \times \left| \begin{array}{c} \text{Diagram: A circle with multiple lines entering from the left. A red line labeled } p_{[im]} \text{ exits to the right.} \end{array} \right|^2 \\
\text{ISR : } C_{im} \Delta^{(m)} & \left| \begin{array}{c} \text{Diagram: A circle with multiple lines entering from the right. A red line labeled } p_m \text{ enters from the top. A red line labeled } p_i \text{ enters from the bottom. A red line labeled } p_{[im]} \text{ enters from the left.} \end{array} \right|^2 \xrightarrow{\theta_{im} \rightarrow 0} \frac{g_{s,b}^2 \mu_0^{2\epsilon}}{p_i \cdot p_m} P_{f_{[im]} f_i, i}(z) \times \frac{1}{z} \left| \begin{array}{c} \text{Diagram: A circle with multiple lines entering from the right. A red line labeled } p_{[im]} \text{ enters from the left.} \end{array} \right|^2
\end{aligned}$$

Figure 1: Graphical representation of the convention adopted for collinear factorization. The top figure describes the final-state splitting process $[im]^* \rightarrow i(z) + m(\bar{z})$, where $z = 1 - E_m/E_{[im]}$, and $\bar{z} = 1 - z$. The bottom figure corresponds to the initial-state splitting process $i \rightarrow [im]^* + m$, with $z = 1 - E_m/E_i$. We note that in the case of FSR, the action of C_{im} on $\Delta^{(m)}$ produces an extra factor of z , which does not occur in ISR.

We emphasize that the above expressions are *exact* and do not contain higher powers of ϵ . To describe soft and collinear limits of the real-virtual contributions, it is convenient to define the following quantities

$$\begin{aligned}
A_K(\epsilon) &= \frac{\Gamma^3(1+\epsilon) \Gamma^5(1-\epsilon)}{\Gamma(1+2\epsilon) \Gamma^2(1-2\epsilon)} = 1 - \frac{\pi^2}{3} \epsilon^2 + \mathcal{O}(\epsilon^3), \\
h_c(\epsilon) &= \frac{\Gamma^2(1-2\epsilon) \Gamma(1+\epsilon)}{\Gamma(1-3\epsilon)} = 1 + \mathcal{O}(\epsilon^3).
\end{aligned} \tag{A.10}$$

A.2 Tree-level, one-loop and Altarelli-Parisi splitting functions

Consider the final-state splitting process $[im]^* \rightarrow i(z) + m(1-z)$, where i and m are two partons with flavors f_i and f_m , respectively. The variable $z = 1 - E_m/E_{[im]}$ represents the energy fraction carried by parton i in the collinear splitting, while parton m takes the energy fraction $1 - z$. As for $[im]$, it is the clustered “mother” parton. We denote the spin-averaged final-state splitting functions as $P_{f_{[im]} f_i}(z)$ (cf. Figure 1, above). These functions read

$$\begin{aligned}
P_{qq}(z) &= C_F \left[\frac{1+z^2}{1-z} - \epsilon(1-z) \right], \\
P_{qg}(z) &= C_F \left[\frac{1+(1-z)^2}{z} - \epsilon z \right] \equiv P_{qg}(1-z), \\
P_{gq}(z) &= T_R \left[1 - \frac{2z(1-z)}{1-\epsilon} \right], \\
P_{gg}(z) &= 2C_A \left[\frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right].
\end{aligned} \tag{A.11}$$

Next, we consider the initial-state splitting process $i \rightarrow [im]^* + m$, where i is now the incoming parton, $[im]$ is the parton that enters the hard scattering, and m is an outgoing

parton. The variable z corresponds to $z = 1 - E_{\mathbf{m}}/E_i$. In this case, we denote the initial-state splitting function, averaged over initial-state color and polarizations, as $P_{[i\mathbf{m}]i,i}(z)$ (cf. Figure 1, below). These functions are related to the splittings in Eq. (A.11) by the following relations

$$\begin{aligned} P_{qq,i}(z) &= -zP_{qq}(1/z) \equiv P_{qq}(z), \\ P_{qg,i}(z) &= \left[\frac{2N_c}{2(1-\epsilon)(N_c^2-1)} \right] zP_{qg}(1/z) \equiv P_{qg}(z), \\ P_{gq,i}(z) &= \left[\frac{2(1-\epsilon)(N_c^2-1)}{2N_c} \right] zP_{gq}(1/z) \equiv P_{gq}(z), \\ P_{gg,i}(z) &= -zP_{gg}(1/z) \equiv P_{gg}(z). \end{aligned} \tag{A.12}$$

We observe that the notation just described for initial- and final-state splittings, and illustrated in Figure 1, applies to all tree-level and one-loop splittings used in this article. Therefore, it must be applied to all functions defined in the remainder of this appendix, as well as those collected in `FinalResult.m` (cf. Table 1) and `UsefulFunctions.m` (cf. Table 2).

Finally, we report the conventional leading-order Altarelli-Parisi splitting functions that we use throughout the paper. They read [94]

$$\begin{aligned} \hat{P}_{qq}^{(0)}(z) &= C_F \left[2\mathcal{D}_0(z) - (1+z) + \frac{3}{2}\delta(1-z) \right], \\ \hat{P}_{qg}^{(0)}(z) &= T_R \left[(1-z)^2 + z^2 \right], \\ \hat{P}_{gq}^{(0)}(z) &= C_F \left[\frac{1 + (1-z)^2}{z} \right], \\ \hat{P}_{gg}^{(0)}(z) &= 2C_A \left[\mathcal{D}_0(z) + z(1-z) + \frac{1}{z} - 2 \right] + \beta_0\delta(1-z). \end{aligned} \tag{A.13}$$

We also require the one-loop Altarelli-Parisi splitting functions; they can be found in Sec. 4.3 of Ref. [94]. They are also given in the ancillary file `UsefulFunctions.m`, see Table 2. We point out that, for the qq splitting, we need the *non-singlet* splitting function from which the analog of the interference contribution of identical quarks has been subtracted.

A.3 Generalized splitting functions and anomalous dimensions

A.3.1 Tree-level

Phase-space remnants and damping factors introduce additional factors to integrands over splitting functions that we require. For this reason, it is convenient to define generalized splitting functions and their integrals.

For the initial-state splitting where parton \mathbf{m} becomes collinear to an initial-state parton a of flavor f_a , the following integral appears

$$E_a^{-k\epsilon} \int_0^1 dz (1-z)^{-k\epsilon} P_{f_{[a\mathbf{m}]}f_a,i}(z) F(z). \tag{A.14}$$

The initial-state splitting functions $P_{\alpha\beta,i}$ are given in Eq. (A.12), and the function $F(z)$ typically contains the relevant matrix element squared. Such integrals may contain infrared divergences which need to be extracted. To facilitate this, we define a new splitting function

$$\mathcal{P}_{f_{[am]}fa}^{(k)}(z, E_a) = \bar{S}_z \left[(1-z)^{-k\epsilon} P_{f_{[am]}fa,i}(z) \right] - 2\delta_{f_{[am]}fa} \mathbf{T}_{f_{[am]}}^2 \frac{1 - e^{-k\epsilon L_a}}{k\epsilon} \delta(1-z). \quad (\text{A.15})$$

Here, $\bar{S}_z = 1 - S_z$, where S_z subtracts the soft singularity at $z = 1$. The relevant powers of phase-space factors $(1-z)^{-\epsilon}$ are $k = 2$ at NLO, and $k = 4$ at NNLO. Explicit expressions for the functions $\mathcal{P}_{\alpha\beta}^{(k)}$ in Eq. (A.15) are provided in the ancillary file `UsefulFunctions.m`, see Table 2. We note that we omit the superscript for $k = 2$ when writing the generalized splitting functions and their integrals to simplify the notation. Furthermore, we point out that functions $\mathcal{P}_{\alpha\beta}^{(k)}(z, E_a)$ in Eq. (A.15) are, in fact, energy-independent if $\alpha \neq \beta$.

For the manipulations described in the main body of the paper, it is convenient to rewrite the generalized splitting functions as follows

$$-\left[\left(\frac{2E_a}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^{\frac{k}{2}} \mathcal{P}_{\alpha\beta}^{(k)}(z, E_a) = \Gamma_{a,\alpha}^{(k)} \delta_{\alpha\beta} \delta(1-z) + \mathcal{P}_{\alpha\beta}^{(k),\text{gen}}(z, E_a), \quad (\text{A.16})$$

with

$$\Gamma_{a,\alpha}^{(k)} = \left[\left(\frac{2E_a}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^{\frac{k}{2}} \left[\gamma_\alpha + 2\mathbf{T}_\alpha^2 \frac{1 - e^{-k\epsilon L_a}}{k\epsilon} \right], \quad (\text{A.17})$$

$$\mathcal{P}_{\alpha\beta}^{(k),\text{gen}}(z, E_a) = \left[\left(\frac{2E_a}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^{\frac{k}{2}} \left[-\hat{P}_{\alpha\beta}^{(0)}(z) + \epsilon \mathcal{P}_{\alpha\beta}^{(k),\text{fin}}(z) \right]. \quad (\text{A.18})$$

Here, $\hat{P}_{\alpha\beta}^{(0)}$ are the Altarelli-Parisi splitting functions given in Eq. (A.13), while $\mathcal{P}_{\alpha\beta}^{(k),\text{fin}}(z)$ corresponds to the ϵ -expansion of $-\mathcal{P}_{\alpha\beta}^{(k)}(z, E_a = 0)$ starting with $\mathcal{O}(\epsilon)$. Explicit expressions for quantities that appear in Eqs (A.17) and (A.18) are provided in `UsefulFunctions.m`, see Table 2.

We continue with the discussion of the final-state splittings. Then, if the final-state parton \mathbf{m} becomes collinear to the final-state parton i of flavour f_i , the generalized final-state anomalous dimension reads

$$\Gamma_{i,f_{[im]}\rightarrow f_i f_{\mathbf{m}}}^{(k)} = \left[\left(\frac{2E_i}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^{\frac{k}{2}} \gamma_{z,f_{[im]}\rightarrow f_i f_{\mathbf{m}}}^{2k}(L_i), \quad (\text{A.19})$$

where

$$\begin{aligned} \gamma_{g(z),f_{[im]}\rightarrow f_i f_{\mathbf{m}}}^{nk}(L_i) = & - \int_0^1 dz \bar{S}_z \left[z^{-n\epsilon} (1-z)^{-k\epsilon} g(z) P_{f_{[im]}f_i}(z) \right] \\ & + 2\delta_{f_{[im]}f_i} \mathbf{T}_{f_{[im]}}^2 \frac{1 - e^{-k\epsilon L_i}}{k\epsilon} g(1). \end{aligned} \quad (\text{A.20})$$

The splitting functions $P_{\alpha\beta}$ can be found in Eq. (A.11), while the explicit expressions for the quantities defined in Eqs (A.19) and (A.20) are provided in `UsefulFunctions.m`, see Table 2.

Finally, we define “physical” combinations of quark and gluon collinear anomalous dimensions

$$\Gamma_{i,q}^{(k)} = \Gamma_{i,q \rightarrow qg}^{(k)} + \Gamma_{i,q \rightarrow gq}^{(k)}, \quad \Gamma_{i,g}^{(k)} = \Gamma_{i,g \rightarrow gg}^{(k)} + 2n_f \Gamma_{i,g \rightarrow q\bar{q}}^{(k)}, \quad (\text{A.21})$$

which appear in the operators I_C and I_T .

A.3.2 Spin-correlations

To describe the spin-correlated contributions arising from sectors $\theta^{(b)}$ and $\theta^{(d)}$, we require the following integrals

$$\begin{aligned} \delta_{\mathbf{m}}(\epsilon) &= \frac{N_\epsilon^{(b,d)} E_{\max}^{4\epsilon}}{2} \int_{E_{\max}}^{2E_{\max}} \frac{dE_{[\mathbf{mn}]}}{E_{[\mathbf{mn}]^{1+4\epsilon}}} \int_{1-\xi}^{\xi} dz [z(1-z)]^{-2\epsilon} [P_{g\mathbf{m}}(z) + \epsilon P_{g\mathbf{m}}^{\perp,r}(z)], \\ \delta_{\mathbf{m}}^{\perp}(\epsilon) &= \frac{N_\epsilon^{(b,d)} E_{\max}^{4\epsilon}}{2} \int_{E_{\max}}^{2E_{\max}} \frac{dE_{[\mathbf{mn}]}}{E_{[\mathbf{mn}]^{1+4\epsilon}}} \int_{1-\xi}^{\xi} dz [z(1-z)]^{-2\epsilon} P_{g\mathbf{m}}^{\perp}(z), \end{aligned} \quad (\text{A.22})$$

where $\xi = E_{\max}/E_{[\mathbf{mn}]}$, \mathbf{m} can be either a quark or a gluon, and the transversal splitting functions are defined as follows

$$\begin{aligned} P_{gg}^{\perp}(z) &= 4C_A(1-\epsilon)z(1-z), & P_{gg}^{\perp,r}(z) &= 2C_A z(1-z)(1-2\epsilon), \\ P_{gq}^{\perp}(z) &= -4T_R z(1-z), & P_{gq}^{\perp,r}(z) &= -2T_R z(1-z) \frac{1-2\epsilon}{1-\epsilon}. \end{aligned} \quad (\text{A.23})$$

The ϵ -expansions of the δ -integrals in Eq. (A.22) are reported in `UsefulFunctions.m`, see Table 2. Note that we combine the gluon and quark components as

$$\delta(\epsilon) = \delta_g(\epsilon) + 2n_f \delta_q(\epsilon), \quad \delta^{\perp}(\epsilon) = \delta_g^{\perp}(\epsilon) + 2n_f \delta_q^{\perp}(\epsilon). \quad (\text{A.24})$$

Additionally, we find it convenient to introduce the following notation for the integrals

$$\gamma_{\perp,g \rightarrow \mathbf{mn}}^{22} = - \int_0^1 dz \frac{P_{g\mathbf{m}}^{\perp}(z)}{[z(1-z)]^{2\epsilon}}, \quad \gamma_{\perp,g \rightarrow \mathbf{mn}}^{22,r} = - \int_0^1 dz \frac{P_{g\mathbf{m}}^{\perp,r}(z)}{[z(1-z)]^{2\epsilon}}, \quad (\text{A.25})$$

where $(\mathbf{mn}) = (gg)$ or $(q\bar{q})$. “Physical” combinations of anomalous dimensions are defined as follows:

$$\gamma_{\perp,g}^{22} = \gamma_{\perp,g \rightarrow gg}^{22} + 2n_f \gamma_{\perp,g \rightarrow q\bar{q}}^{22}, \quad \gamma_{\perp,g}^{22,r} = \gamma_{\perp,g \rightarrow gg}^{22,r} + 2n_f \gamma_{\perp,g \rightarrow q\bar{q}}^{22,r}. \quad (\text{A.26})$$

In the $\epsilon = 0$ limit, they read

$$\gamma_{\perp,g}^{22} = -\frac{2}{3}(C_A - 2n_f T_R), \quad \gamma_{\perp,g}^{22,r} = -\frac{1}{3}(C_A - 2n_f T_R), \quad (\text{A.27})$$

and are reported in `FinalResult.m`, see Table 1.

A.3.3 One-loop

Similar to the discussion of the generalized tree-level splitting functions and anomalous dimensions in Sec. A.3.1, we need to define generalized splitting functions for the one-loop case. For the initial state splitting, when a parton \mathbf{m} becomes collinear to an initial-state parton a with flavor f_a , we define

$$\begin{aligned} \mathcal{P}_{f_{[am]}f_a}^{(k),1L}(z, E_a) &= \overline{S}_z \left[(1-z)^{-k\epsilon} P_{f_{[am]}f_a, i}^{1L}(z) \right] \\ &+ 2\delta_{f_{[am]}f_a} C_A \mathbf{T}_{f_{[am]}}^2 \frac{1 - e^{-(2+k)\epsilon L_a}}{(2+k)\epsilon^2} \pi \cot(\pi\epsilon) \delta(1-z). \end{aligned} \quad (\text{A.28})$$

The splitting functions $P_{\alpha\beta, i}^{1L}$ can be obtained from Refs [95–97], and they are collected in `UsefulFunctions.m`, see Table 2. In analogy with Eq. (A.16), we find it convenient to rewrite the splitting function $\mathcal{P}_{\alpha\beta}^{(k),1L}$ as follows

$$\frac{\epsilon^2}{C_x} \left[\left(\frac{2E_a}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^k \mathcal{P}_{\alpha\beta}^{(k),1L}(z, E_a) = \Gamma_{a,\alpha}^{(k),1L} \delta_{\alpha\beta} \delta(1-z) + \mathcal{P}_{\alpha\beta}^{(k),1L, \text{gen}}(z, E_a), \quad (\text{A.29})$$

where $C_x = 2C_F - C_A$ if $\alpha\beta = gq$, and $C_x = C_A$ otherwise. Furthermore, we define

$$\Gamma_{a,\alpha}^{(k),1L} = \left[\left(\frac{2E_a}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^k \left[\gamma_\alpha + 2\mathbf{T}_\alpha^2 \frac{1 - e^{-(2+k)\epsilon L_a}}{(2+k)} \pi \frac{\cos(\pi\epsilon)}{\sin(\pi\epsilon)} \right], \quad (\text{A.30})$$

$$\mathcal{P}_{\alpha\beta}^{(k),1L, \text{gen}}(z, E_a) = \left[\left(\frac{2E_a}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^k \left[-\hat{P}_{\alpha\beta}^{(0)}(z) + \epsilon \mathcal{P}_{\alpha\beta}^{(k),1L, \text{fin}}(z) \right]. \quad (\text{A.31})$$

In Eq. (A.31), $\mathcal{P}_{\alpha\beta}^{(k),1L, \text{fin}}(z)$ corresponds to the ϵ -expansion of

$$\frac{\epsilon^2}{C_x} \mathcal{P}_{\alpha\beta}^{(k),1L}(z, E_a = 0), \quad (\text{A.32})$$

starting from $\mathcal{O}(\epsilon)$. Note that, although the above definitions are provided for any value of k , we set $k = 2$ in the real-virtual contributions. The explicit expression of the functions in Eqs (A.28), (A.30), (A.31), with $k = 2$, are provided in `UsefulFunctions.m`, see Table 2.

If the unresolved final-state parton \mathbf{m} goes collinear to a final-state parton, say i of flavor f_i , the generalized final-state anomalous dimensions read

$$\Gamma_{i, f_{[im]} \rightarrow f_i f_m}^{(k),1L} = - \left[\left(\frac{2E_i}{\mu} \right)^{-2\epsilon} \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \right]^k \frac{\epsilon^2 \cos(\pi\epsilon)}{C_x} \gamma_{z, f_{[im]} \rightarrow f_i f_m}^{3(k+1),1L}(L_i), \quad (\text{A.33})$$

where $C_x = 2C_F - C_A$ for the $g \rightarrow q\bar{q}$ splitting, and $C_x = C_A$ otherwise. The one-loop anomalous dimensions $\gamma_{g(z), f_{[im]} \rightarrow f_i f_m}^{n(k+1),1L}$ that appear in Eq. (A.33) are defined as

$$\begin{aligned} \gamma_{g(z), f_{[im]} \rightarrow f_i f_m}^{n(k+1),1L}(L_i) &= - \int_0^1 dz \overline{S}_z \left[z^{-n\epsilon} (1-z)^{-(k+1)\epsilon} g(z) P_{f_{[im]}f_i}^{1L}(z) \right] \\ &- 2\delta_{f_{[im]}f_i} C_A \mathbf{T}_{f_{[im]}}^2 \frac{1 - e^{-(2+k)\epsilon L_i}}{(2+k)} \frac{\pi}{\epsilon^2 \sin(\pi\epsilon)} g(1). \end{aligned} \quad (\text{A.34})$$

The quantities defined in Eqs (A.33) and (A.34), with $n = 3$ and $k = 2$, are provided in `UsefulFunctions.m`, together with the one-loop final-state splitting functions $P_{\alpha\beta}^{1L}$ (see Table 2). We note that, similar to the expressions in Eq. (A.21), we define the one-loop case as follows

$$\Gamma_{i,q}^{1L} = \Gamma_{i,q \rightarrow qg}^{1L} + \Gamma_{i,q \rightarrow gq}^{1L}, \quad \Gamma_{i,g}^{1L} = \Gamma_{i,g \rightarrow gg}^{1L} + 2n_f \Gamma_{i,g \rightarrow q\bar{q}}^{1L}. \quad (\text{A.35})$$

A.4 Definitions and relevant properties of I -operators

In Ref. [1], we have introduced virtual, soft, and collinear operators, and we employ similar operators in this paper as well. The virtual operator is defined as

$$I_V(\epsilon) = \bar{I}_1(\epsilon) + \bar{I}_1^\dagger(\epsilon), \quad (\text{A.36})$$

where [87]

$$\bar{I}_1(\epsilon) = \frac{1}{2} \sum_{(ij)} \frac{\mathcal{V}_i^{\text{sing}}(\epsilon)}{\mathbf{T}_i^2} \mathbf{T}_i \cdot \mathbf{T}_j \left(\frac{\mu^2}{2p_i \cdot p_j} \right)^\epsilon e^{i\pi\lambda_{ij}\epsilon}, \quad \mathcal{V}_i^{\text{sing}}(\epsilon) = \frac{\mathbf{T}_i^2}{\epsilon^2} + \frac{\gamma_i}{\epsilon}. \quad (\text{A.37})$$

In Eq. (A.37), γ_i are the anomalous dimensions that can be found in (A.3), the sum goes over the unordered pairs of external particles i, j with $i \neq j$, and λ_{ij} are constants which are equal to 1 if both i and j are incoming or outgoing, and to 0 otherwise. The soft operators which appear in the double-real and in the real-virtual corrections read

$$\begin{aligned} I_S(\epsilon) &= -\frac{(2E_{\text{max}}/\mu)^{-2\epsilon}}{\epsilon^2} \sum_{(ij)} \eta_{ij}^{-\epsilon} K_{ij}(\mathbf{T}_i \cdot \mathbf{T}_j), \\ \tilde{I}_S(2\epsilon) &= -\frac{(2E_{\text{max}}/\mu)^{-4\epsilon}}{(2\epsilon)^2} \sum_{(ij)} \eta_{ij}^{-2\epsilon} \tilde{K}_{ij}(\mathbf{T}_i \cdot \mathbf{T}_j), \end{aligned} \quad (\text{A.38})$$

where $\eta_{ij} = (1 - \cos \theta_{ij})/2$, and

$$\begin{aligned} K_{ij} &= \frac{\Gamma^2(1-\epsilon)}{\Gamma(1-2\epsilon)} \eta_{ij}^{1+\epsilon} {}_2F_1(1, 1, 1-\epsilon, 1-\eta_{ij}), \\ \tilde{K}_{ij} &= \frac{\Gamma^2(1-2\epsilon)}{\Gamma(1-4\epsilon)} \eta_{ij}^{1+3\epsilon} {}_2F_1(1+\epsilon, 1+\epsilon, 1-\epsilon, 1-\eta_{ij}). \end{aligned} \quad (\text{A.39})$$

A useful relation between I_S and \tilde{I}_S is

$$\tilde{I}_S(2\epsilon) = I_S(2\epsilon) + \mathcal{O}(\epsilon). \quad (\text{A.40})$$

The Laurent expansions of $I_V(\epsilon)$, $I_S(\epsilon)$, and $\tilde{I}_S(2\epsilon)$ can be found in Appendix A.5 of Ref. [1].

The collinear operators which appear in the double-real and in the real-virtual corrections read, respectively,

$$I_C^{(k)}(\epsilon) = \sum_{i \in \mathcal{H}} \frac{\Gamma_{i,f_i}^{(k)}}{\epsilon^{(k/2)}}, \quad \tilde{I}_C(2\epsilon) = \sum_{i \in \mathcal{H}} \frac{\Gamma_{i,f_i}^{1L}}{2\epsilon}, \quad (\text{A.41})$$

where generalized initial- and final-state anomalous dimensions are given in Eqs (A.17) and (A.19) for $I_C^{(k)}$, and in Eqs (A.30) and (A.33) for \tilde{I}_C . We note that the following relations hold

$$\tilde{I}_C(2\epsilon) = I_C(2\epsilon) + \mathcal{O}(\epsilon^0), \quad I_C^{(4)}(\epsilon) = I_C(2\epsilon) + \mathcal{O}(\epsilon^0). \quad (\text{A.42})$$

Finally, the ϵ -finite operator I_T is defined as follows

$$I_T(\epsilon) = I_V(\epsilon) + I_S(\epsilon) + I_C(\epsilon). \quad (\text{A.43})$$

Its expansion in powers of ϵ reads

$$I_T(\epsilon) = \sum_{n=0}^{\infty} \epsilon^n I_T^{(n)}, \quad (\text{A.44})$$

where the $n = 0, 1$ coefficients are given by

$$\begin{aligned} I_T^{(0)} = & - \sum_{(ij)} (\mathbf{T}_i \cdot \mathbf{T}_j) \left[\left(2L_{\max} + \frac{1}{2} \log \eta_{ij} \right) \log \eta_{ij} - \frac{1}{2} L_{ij} \left(L_{ij} + \frac{2\gamma_i}{\mathbf{T}_i^2} \right) \right. \\ & + \text{Li}_2(1 - \eta_{ij}) + \frac{\pi^2}{2} \lambda_{ij} \Big] + \sum_{i \in \mathcal{H}} \mathbf{T}_i^2 \left[2L_{\max}^2 - \frac{\pi^2}{6} - \left(2\tilde{L}_i \gamma_i^{22,(0)} - \gamma_i^{22,(1)} \right) \frac{\theta_{\mathcal{H}_f}}{\mathbf{T}_i^2} \right. \\ & \left. \left. - 2 \left(L_i^2 + 2L_i \tilde{L}_i + \tilde{L}_i \frac{\gamma_i}{\mathbf{T}_i^2} \right) \bar{\theta}_{\mathcal{H}_f} \right], \end{aligned} \quad (\text{A.45})$$

$$\begin{aligned} I_T^{(1)} = & \sum_{(ij)} (\mathbf{T}_i \cdot \mathbf{T}_j) \left[\frac{1}{6} (L_{ij}^3 + \log^3 \eta_{ij}) + 2L_{\max}^2 \log \eta_{ij} - \frac{\pi^2}{2} \lambda_{ij} L_{ij} \right. \\ & + \left(L_{\max} - \frac{1}{2} \log(1 - \eta_{ij}) \right) \log^2 \eta_{ij} + 2L_{\max} \text{Li}_2(1 - \eta_{ij}) - \text{Li}_3(\eta_{ij}) \\ & \left. - \text{Li}_3(1 - \eta_{ij}) + \frac{\gamma_i}{2\mathbf{T}_i^2} (L_{ij}^2 - \pi^2 \lambda_{ij}) \right] + \sum_{i \in \mathcal{H}} \mathbf{T}_i^2 \left\{ -\frac{4}{3} L_{\max}^3 + \frac{\pi^2}{3} L_{\max} - 3\zeta_3 \right. \\ & + \left[\left(2\tilde{L}_i^2 - \frac{\pi^2}{6} \right) \gamma_i^{22,(0)} - 2\tilde{L}_i \gamma_i^{22,(1)} + \gamma_i^{22,(2)} \right] \frac{\theta_{\mathcal{H}_f}}{\mathbf{T}_i^2} + \left[\frac{4}{3} L_i^3 + 4L_i^2 \tilde{L}_i + 4L_i \tilde{L}_i^2 \right. \\ & \left. \left. - \frac{\pi^2}{3} L_i + \frac{\gamma_i}{\mathbf{T}_i^2} \left(2\tilde{L}_i^2 - \frac{\pi^2}{6} \right) \right] \bar{\theta}_{\mathcal{H}_f} \right\}. \end{aligned} \quad (\text{A.46})$$

To write the above equations, we have used

$$\theta_{\mathcal{H}_f} = \begin{cases} 1 & \text{if } i \in \mathcal{H}_f \\ 0 & \text{otherwise,} \end{cases} \quad \bar{\theta}_{\mathcal{H}_f} = 1 - \theta_{\mathcal{H}_f}, \quad (\text{A.47})$$

together with the expansion coefficients of the collinear anomalous dimensions $\gamma_i^{22}(L_i)$, defined as

$$\gamma_i^{22}(L_i) = \sum_{n=0}^{\infty} \epsilon^n \gamma_i^{22,(n)}(L_i), \quad i = g, q. \quad (\text{A.48})$$

We note that in Eqs (A.45) and (A.46), the dependence of the coefficients $\gamma_i^{22,(n)}$ on the logarithms L_i is not shown explicitly.

A.5 Double-unresolved contributions

In this section, we define the double-unresolved terms in Eqs (4.12) and (4.13) which were left unspecified in the main body of the paper. These contributions are at most of $\mathcal{O}(\epsilon^{-1})$. We start from Eq. (4.12), which refers to the $(\mathbf{mn}) \in \text{DS}$ case (cf. Eq. (4.7)). We write¹⁰

$$\Sigma_{\text{DU}}^{\mathcal{A}_n, \text{rest}}[(\mathbf{mn}) \in \text{DS}]_{\text{RR}} = \Sigma_{\text{DU}}^{\text{rdc}}|_{\mathcal{A}_n} + \Sigma_{\text{DU}}^{(2)}|_{\mathcal{A}_n} + \Sigma_{\text{DU}}^{(8)}|_{\mathcal{A}_n} + \sum_{i=1}^5 \Sigma_{\text{DU}}^{\text{fin},(i)}|_{\mathcal{A}_n}, \quad (\text{A.49})$$

with

$$\Sigma_{\text{DU}}^{\text{rdc}}|_{\mathcal{A}_n} = \frac{1}{2} \sum_{i \in \mathcal{H}} \left\langle [2(\eta_{in}/2)^{-\epsilon} - 1] \mathcal{S}(\mathbf{m}, \mathbf{n}) C_{in} C_{im} \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \right\rangle, \quad (\text{A.50})$$

$$\Sigma_{\text{DU}}^{(2)}|_{\mathcal{A}_n} = \langle \bar{S}_{\mathbf{mn}} \bar{S}_{\mathbf{n}} \Omega_2 \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad (\text{A.51})$$

$$\begin{aligned} \Sigma_{\text{DU}}^{(8)}|_{\mathcal{A}_n} = \sum_{i \in \mathcal{H}} [\alpha_s] \frac{N_{\epsilon}^{(b,d)}}{2} \left\langle \bar{S}_{[\mathbf{mn}]} C_{i[\mathbf{mn}]} (E_{[\mathbf{mn}]} / \mu)^{-2\epsilon} \sigma_{i[\mathbf{mn}]}^{-\epsilon} \left[\gamma_{\perp, g \rightarrow \mathbf{mn}}^{22} (r_i^{\mu} r_i^{\nu} + g^{\mu\nu}) \right. \right. \\ \left. \left. - \gamma_{\perp, g \rightarrow \mathbf{mn}}^{22, \text{r}} g^{\mu\nu} \right] \Delta^{(\mathbf{m})} F_{\text{LM}, \mu\nu}^{\mathcal{A}_n}[[\mathbf{mn}]] \right\rangle, \end{aligned} \quad (\text{A.52})$$

$$\Sigma_{\text{DU}}^{\text{fin},(1)}|_{\mathcal{A}_n} = - \left[\left(\frac{\Gamma(1-2\epsilon)}{\Gamma^2(1-\epsilon)} \right)^2 - 1 \right] \sum_{(ij)} \langle \bar{S}_{\mathbf{n}} C_{jn} C_{im} \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad (\text{A.53})$$

$$\Sigma_{\text{DU}}^{\text{fin},(2)}|_{\mathcal{A}_n} = \sum_{i \in \mathcal{H}} \left\langle [(\eta_{im}/2)^{-\epsilon} - 1] \bar{C}_{im} S_{\mathbf{m}} \langle \bar{S}_{\mathbf{n}} C_{in} \omega^{\mathbf{mi}, \mathbf{ni}} \Delta^{(\mathbf{mn})} \Theta_{\mathbf{mn}} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle \right\rangle, \quad (\text{A.54})$$

$$\begin{aligned} \Sigma_{\text{DU}}^{\text{fin},(3)}|_{\mathcal{A}_n} = \left[\frac{\Gamma(1-2\epsilon)}{\Gamma^2(1-\epsilon)} - 1 \right] \left\langle \mathcal{S}(\mathbf{m}, \mathbf{n}) \left[\sum_{(ij)} C_{jn} C_{im} + \sum_{i \in \mathcal{H}} (\eta_{in}/2)^{-\epsilon} C_{in} C_{im} \right] \right. \\ \left. \times \Delta^{(\mathbf{mn})} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \right\rangle, \end{aligned} \quad (\text{A.55})$$

$$\Sigma_{\text{DU}}^{\text{fin},(4)}|_{\mathcal{A}_n} = [\alpha_s]^2 2^{2\epsilon} \delta_{\mathbf{m}}(\epsilon) \left(\frac{2E_{\text{max}}}{\mu} \right)^{-4\epsilon} \sum_{i \in \mathcal{H}} \langle \mathcal{W}_i^{\mathbf{m}||\mathbf{n}, \text{fin}} \cdot F_{\text{LM}}^{\mathcal{A}_n} \rangle, \quad (\text{A.56})$$

$$\Sigma_{\text{DU}}^{\text{fin},(5)}|_{\mathcal{A}_n} = [\alpha_s]^2 \delta_{\mathbf{m}}^{\perp}(\epsilon) \left(\frac{E_{\text{max}}}{\mu} \right)^{-4\epsilon} \sum_{i \in \mathcal{H}} \langle \mathcal{W}_i^{(i)} \cdot F_{\text{LM}}^{\mathcal{A}_n} \rangle. \quad (\text{A.57})$$

The terms in Eqs (A.50 – A.52) are $\mathcal{O}(\epsilon^{-1})$, whereas the terms in Eqs (A.53 – A.57) are ϵ -finite. In Eqs (A.50) and (A.55), the operator $\mathcal{S}(\mathbf{m}, \mathbf{n})$ is defined as

$$\mathcal{S}(\mathbf{m}, \mathbf{n}) = \begin{cases} \bar{S}_{\mathbf{n}}(\mathbb{1} - S_{\mathbf{m}} \Theta_{\mathbf{nm}}) + S_{\mathbf{n}} \bar{S}_{\mathbf{m}} \Theta_{\mathbf{nm}}, & \text{if } (\mathbf{mn}) = (gg), \\ \mathbb{1}, & \text{if } (\mathbf{mn}) = (q_i \bar{q}_i). \end{cases} \quad (\text{A.58})$$

In Eq. (A.51), the operator Ω_2 is the triple-collinear operator defined in Eq. (D.6) of Ref. [1]. The triple-collinear splittings are reported in `UsefulFunctions.m`, see Table 2.

¹⁰The numbering scheme for contributions to $\Sigma_{\text{DU}}|_{\mathcal{A}_n}$ and $\Sigma_{\text{DU}}^{\text{fin}}|_{\mathcal{A}_n}$ is kept consistent with the definitions introduced in Ref. [1].

In Eq. (A.52), we used $\sigma_{ij} = \eta_{ij}/(1 - \eta_{ij})$, the anomalous dimensions $\gamma_{\perp, g \rightarrow mn}^{22}$ and $\gamma_{\perp, g \rightarrow mn}^{22, r}$ defined in Eq. (A.25), and the vector r_i^μ specified in Appendix E of Ref. [1]. The quantities $\delta_m(\epsilon)$ and $\delta_m^\perp(\epsilon)$ appearing in Eqs (A.56 – A.57) are defined in Eq. (A.22), while the two partition-dependent operators $\mathcal{W}_i^{m||n, \text{fin}}$ and $\mathcal{W}_r^{(i)}$ are defined in Eq. (7.15).

For the double-unresolved terms appearing in Eq. (4.13), which refers to $(mn) \in \mathcal{DS}$ case, we write

$$\Sigma_{\text{DU}}^{\mathcal{A}_n, \text{rest}}[(mn) \in \mathcal{DS}]_{\text{RR}} = \Sigma_{\text{DU}}^{\text{rdc}}|_{\mathcal{A}_n} + \Sigma_{\text{DU}}^{(2)}|_{\mathcal{A}_n} + \sum_{i=1,3} \Sigma_{\text{DU}}^{\text{fin},(i)}|_{\mathcal{A}_n}, \quad (\text{A.59})$$

where

$$\Sigma_{\text{DU}}^{\text{rdc}}|_{\mathcal{A}_n} = \frac{1}{2} \sum_{i \in \mathcal{H}} \langle [2(\eta_{in}/2)^{-\epsilon} - 1] \bar{S}_n (C_{im} C_{in} + C_{in} C_{im}) \Delta^{(mn)} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad (\text{A.60})$$

$$\Sigma_{\text{DU}}^{(2)}|_{\mathcal{A}_n} = \langle \bar{S}_n \Omega_2 \Delta^{(mn)} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad (\text{A.61})$$

$$\Sigma_{\text{DU}}^{\text{fin},(1)}|_{\mathcal{A}_n} = - \left[\left(\frac{\Gamma(1-2\epsilon)}{\Gamma^2(1-\epsilon)} \right)^2 - 1 \right] \sum_{(ij)} \langle \bar{S}_n C_{jn} C_{im} \Delta^{(mn)} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \rangle, \quad (\text{A.62})$$

$$\begin{aligned} \Sigma_{\text{DU}}^{\text{fin},(3)}|_{\mathcal{A}_n} = & \left[\frac{\Gamma(1-2\epsilon)}{\Gamma^2(1-\epsilon)} - 1 \right] \left\langle \bar{S}_n \left[2 \sum_{(ij)} C_{jn} C_{im} + \sum_{i \in \mathcal{H}} (\eta_{in}/2)^{-\epsilon} (C_{in} C_{im} \right. \right. \\ & \left. \left. + C_{im} C_{in}) \right] \Delta^{(mn)} F_{\text{LM}}^{\mathcal{A}_n}[\mathbf{m}, \mathbf{n}] \right\rangle. \end{aligned} \quad (\text{A.63})$$

The terms in Eqs (A.60 – A.61) are $\mathcal{O}(\epsilon^{-1})$, and those in Eqs (A.62 – A.63) are ϵ -finite.

References

- [1] F. Devoto, K. Melnikov, R. Röntsch, C. Signorile-Signorile and D.M. Tagliabue, *A fresh look at the nested soft-collinear subtraction scheme: NNLO QCD corrections to N -gluon final states in $q\bar{q}$ annihilation*, *JHEP* **02** (2024) 016 [[2310.17598](#)].
- [2] S. Frixione, Z. Kunszt and A. Signer, *Three jet cross-sections to next-to-leading order*, *Nucl. Phys. B* **467** (1996) 399 [[hep-ph/9512328](#)].
- [3] S. Catani and M. Seymour, *A General algorithm for calculating jet cross-sections in NLO QCD*, *Nucl. Phys. B* **485** (1997) 291 [[hep-ph/9605323](#)].
- [4] Z. Nagy and D.E. Soper, *General subtraction method for numerical calculation of one loop QCD matrix elements*, *JHEP* **09** (2003) 055 [[hep-ph/0308127](#)].
- [5] G. Bevilacqua, M. Czakon, M. Kubocz and M. Worek, *Complete Nagy-Soper subtraction for next-to-leading order calculations in QCD*, *JHEP* **10** (2013) 204 [[1308.5605](#)].
- [6] S. Frixione and M. Grazzini, *Subtraction at NNLO*, *JHEP* **06** (2005) 010 [[hep-ph/0411399](#)].
- [7] A. Gehrmann-De Ridder, T. Gehrmann and E.W.N. Glover, *Antenna subtraction at NNLO*, *JHEP* **09** (2005) 056 [[hep-ph/0505111](#)].
- [8] J. Currie, E.W.N. Glover and S. Wells, *Infrared Structure at NNLO Using Antenna Subtraction*, *JHEP* **04** (2013) 066 [[1301.4693](#)].
- [9] G. Somogyi, Z. Trocsanyi and V. Del Duca, *Matching of singly- and doubly-unresolved limits of tree-level QCD squared matrix elements*, *JHEP* **06** (2005) 024 [[hep-ph/0502226](#)].
- [10] G. Somogyi and Z. Trocsanyi, *A Subtraction scheme for computing QCD jet cross sections at NNLO: Regularization of real-virtual emission*, *JHEP* **01** (2007) 052 [[hep-ph/0609043](#)].
- [11] V. Del Duca, C. Duhr, A. Kardos, G. Somogyi and Z. Trócsányi, *Three-Jet Production in Electron-Positron Collisions at Next-to-Next-to-Leading Order Accuracy*, *Phys. Rev. Lett.* **117** (2016) 152004 [[1603.08927](#)].
- [12] V. Del Duca, C. Duhr, A. Kardos, G. Somogyi, Z. Ször, Z. Trócsányi et al., *Jet production in the CoLoRFulNNLO method: event shapes in electron-positron collisions*, *Phys. Rev. D* **94** (2016) 074019 [[1606.03453](#)].
- [13] M. Czakon, *A novel subtraction scheme for double-real radiation at NNLO*, *Phys. Lett. B* **693** (2010) 259 [[1005.0274](#)].
- [14] M. Czakon, *Double-real radiation in hadronic top quark pair production as a proof of a certain concept*, *Nucl. Phys. B* **849** (2011) 250 [[1101.0642](#)].
- [15] M. Czakon and D. Heymes, *Four-dimensional formulation of the sector-improved residue subtraction scheme*, *Nucl. Phys. B* **890** (2014) 152 [[1408.2500](#)].
- [16] C. Anastasiou, K. Melnikov and F. Petriello, *A new method for real radiation at NNLO*, *Phys. Rev. D* **69** (2004) 076010 [[hep-ph/0311311](#)].
- [17] F. Caola, K. Melnikov and R. Röntsch, *Nested soft-collinear subtractions in NNLO QCD computations*, *Eur. Phys. J. C* **77** (2017) 248 [[1702.01352](#)].
- [18] S. Catani and M. Grazzini, *An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC*, *Phys. Rev. Lett.* **98** (2007) 222002 [[hep-ph/0703012](#)].

- [19] M. Grazzini, S. Kallweit and M. Wiesemann, *Fully differential NNLO computations with MATRIX*, *Eur. Phys. J. C* **78** (2018) 537 [[1711.06631](#)].
- [20] R. Boughezal, K. Melnikov and F. Petriello, *A subtraction scheme for NNLO computations*, *Phys. Rev. D* **85** (2012) 034025 [[1111.7041](#)].
- [21] J. Gaunt, M. Stahlhofen, F.J. Tackmann and J.R. Walsh, *N-jettiness Subtractions for NNLO QCD Calculations*, *JHEP* **09** (2015) 058 [[1505.04794](#)].
- [22] R. Boughezal, C. Focke, X. Liu and F. Petriello, *W-boson production in association with a jet at next-to-next-to-leading order in perturbative QCD*, *Phys. Rev. Lett.* **115** (2015) 062002 [[1504.02131](#)].
- [23] G.F.R. Sborlini, F. Driencourt-Mangin and G. Rodrigo, *Four-dimensional unsubtraction with massive particles*, *JHEP* **10** (2016) 162 [[1608.01584](#)].
- [24] F. Herzog, *Geometric IR subtraction for final state real radiation*, *JHEP* **08** (2018) 006 [[1804.07949](#)].
- [25] L. Magnea, E. Maina, G. Pelliccioli, C. Signorile-Signorile, P. Torrielli and S. Uccirati, *Local analytic sector subtraction at NNLO*, *JHEP* **12** (2018) 107 [[1806.09570](#)].
- [26] L. Magnea, G. Pelliccioli, C. Signorile-Signorile, P. Torrielli and S. Uccirati, *Analytic integration of soft and collinear radiation in factorised QCD cross sections at NNLO*, *JHEP* **02** (2021) 037 [[2010.14493](#)].
- [27] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss and M. Marcoli, *Automation of antenna subtraction in colour space: gluonic processes*, *JHEP* **10** (2022) 099 [[2203.13531](#)].
- [28] G. Bertolotti, L. Magnea, G. Pelliccioli, A. Ratti, C. Signorile-Signorile, P. Torrielli et al., *NNLO subtraction for any massless final state: a complete analytic expression*, *JHEP* **07** (2023) 140 [[2212.11190](#)].
- [29] Z. Capatti, V. Hirschi, D. Kermanschah and B. Ruijl, *Loop-Tree Duality for Multiloop Numerical Integration*, *Phys. Rev. Lett.* **123** (2019) 151602 [[1906.06138](#)].
- [30] O. Braun-White, N. Glover and C.T. Preuss, *A general algorithm to build real-radiation antenna functions for higher-order calculations*, *JHEP* **06** (2023) 065 [[2302.12787](#)].
- [31] O. Braun-White, N. Glover and C.T. Preuss, *A general algorithm to build mixed real and virtual antenna functions for higher-order calculations*, *JHEP* **11** (2023) 179 [[2307.14999](#)].
- [32] E. Fox and N. Glover, *Initial-final and initial-initial antenna functions for real radiation at next-to-leading order*, *JHEP* **12** (2023) 171 [[2308.10829](#)].
- [33] T. Gehrmann, E.W.N. Glover and M. Marcoli, *The colourful antenna subtraction method*, *JHEP* **03** (2024) 114 [[2310.19757](#)].
- [34] X. Chen, T. Gehrmann, E.W.N. Glover and M. Jaquier, *Precise QCD predictions for the production of Higgs + jet final states*, *Phys. Lett. B* **740** (2015) 147 [[1408.5325](#)].
- [35] R. Boughezal, F. Caola, K. Melnikov, F. Petriello and M. Schulze, *Higgs boson production in association with a jet at next-to-next-to-leading order*, *Phys. Rev. Lett.* **115** (2015) 082003 [[1504.07922](#)].
- [36] F. Caola, K. Melnikov and M. Schulze, *Fiducial cross sections for Higgs boson production in association with a jet at next-to-next-to-leading order in QCD*, *Phys. Rev. D* **92** (2015) 074032 [[1508.02684](#)].

- [37] X. Chen, J. Cruz-Martinez, T. Gehrmann, E.W.N. Glover and M. Jaquier, *NNLO QCD corrections to Higgs boson production at large transverse momentum*, *JHEP* **10** (2016) 066 [[1607.08817](#)].
- [38] J.M. Campbell, R.K. Ellis and S. Seth, *H + 1 jet production revisited*, *JHEP* **10** (2019) 136 [[1906.01020](#)].
- [39] M. Cacciari, F.A. Dreyer, A. Karlberg, G.P. Salam and G. Zanderighi, *Fully Differential Vector-Boson-Fusion Higgs Production at Next-to-Next-to-Leading Order*, *Phys. Rev. Lett.* **115** (2015) 082002 [[1506.02660](#)].
- [40] J. Cruz-Martinez, T. Gehrmann, E.W.N. Glover and A. Huss, *Second-order QCD effects in Higgs boson production through vector boson fusion*, *Phys. Lett. B* **781** (2018) 672 [[1802.02445](#)].
- [41] R. Gauld, A. Gehrmann-De Ridder, E.W.N. Glover, A. Huss and I. Majer, *VH + jet production in hadron-hadron collisions up to order α_s^3 in perturbative QCD*, *JHEP* **03** (2022) 008 [[2110.12992](#)].
- [42] S. Catani, S. Devoto, M. Grazzini, S. Kallweit, J. Mazzitelli and C. Savoini, *Higgs Boson Production in Association with a Top-Antitop Quark Pair in Next-to-Next-to-Leading Order QCD*, *Phys. Rev. Lett.* **130** (2023) 111902 [[2210.07846](#)].
- [43] H.A. Chawdhry, M.L. Czakon, A. Mitov and R. Poncelet, *NNLO QCD corrections to three-photon production at the LHC*, *JHEP* **02** (2020) 057 [[1911.00479](#)].
- [44] H.A. Chawdhry, M. Czakon, A. Mitov and R. Poncelet, *NNLO QCD corrections to diphoton production with an additional jet at the LHC*, *JHEP* **09** (2021) 093 [[2105.06940](#)].
- [45] M. Czakon, A. Mitov, M. Pellen and R. Poncelet, *NNLO QCD predictions for W+c-jet production at the LHC*, *JHEP* **06** (2021) 100 [[2011.01011](#)].
- [46] R. Gauld, A. Gehrmann-De Ridder, E.W.N. Glover, A. Huss, A.R. Garcia and G. Stagnitto, *NNLO QCD predictions for Z-boson production in association with a charm jet within the LHCb fiducial region*, *Eur. Phys. J. C* **83** (2023) 336 [[2302.12844](#)].
- [47] J. Currie, A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, A. Huss and J. Pires, *Precise predictions for dijet production at the LHC*, *Phys. Rev. Lett.* **119** (2017) 152001 [[1705.10271](#)].
- [48] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss and J. Mo, *NNLO QCD corrections in full colour for jet production observables at the LHC*, *JHEP* **09** (2022) 025 [[2204.10173](#)].
- [49] S. Badger, M. Czakon, H.B. Hartanto, R. Moodie, T. Peraro, R. Poncelet et al., *Isolated photon production in association with a jet pair through next-to-next-to-leading order in QCD*, *JHEP* **10** (2023) 071 [[2304.06682](#)].
- [50] M. Czakon, A. Mitov and R. Poncelet, *Next-to-Next-to-Leading Order Study of Three-Jet Production at the LHC*, *Phys. Rev. Lett.* **127** (2021) 152001 [[2106.05331](#)].
- [51] M. Czakon, D. Heymes and A. Mitov, *High-precision differential predictions for top-quark pairs at the LHC*, *Phys. Rev. Lett.* **116** (2016) 082003 [[1511.00549](#)].
- [52] S. Catani, S. Devoto, M. Grazzini, S. Kallweit and J. Mazzitelli, *Top-quark pair production at the LHC: Fully differential QCD predictions at NNLO*, *JHEP* **07** (2019) 100 [[1906.06535](#)].
- [53] L. Buonocore, S. Devoto, M. Grazzini, S. Kallweit, J. Mazzitelli, L. Rottoli et al., *Precise*

Predictions for the Associated Production of a W Boson with a Top-Antitop Quark Pair at the LHC, *Phys. Rev. Lett.* **131** (2023) 231901 [[2306.16311](#)].

- [54] M. Brucherseifer, F. Caola and K. Melnikov, *On the NNLO QCD corrections to single-top production at the LHC*, *Phys. Lett. B* **736** (2014) 58 [[1404.7116](#)].
- [55] E.L. Berger, J. Gao, C.P. Yuan and H.X. Zhu, *NNLO QCD Corrections to t-channel Single Top-Quark Production and Decay*, *Phys. Rev. D* **94** (2016) 071501 [[1606.08463](#)].
- [56] J. Campbell, T. Neumann and Z. Sullivan, *Single-top-quark production in the t-channel at NNLO*, *JHEP* **02** (2021) 040 [[2012.01574](#)].
- [57] C. Brønnum-Hansen, K. Melnikov, J. Quarrroz, C. Signorile-Signorile and C.-Y. Wang, *Non-factorisable contribution to t-channel single-top production*, *JHEP* **06** (2022) 061 [[2204.05770](#)].
- [58] M. Alvarez, J. Cantero, M. Czakon, J. Llorente, A. Mitov and R. Poncelet, *NNLO QCD corrections to event shapes at the LHC*, *JHEP* **03** (2023) 129 [[2301.01086](#)].
- [59] E. Fox, N. Glover and M. Marcoli, *Generalised antenna functions for higher-order calculations*, *JHEP* **12** (2024) 225 [[2410.12904](#)].
- [60] S. Catani, D. de Florian, G. Ferrera and M. Grazzini, *Vector boson production at hadron colliders: transverse-momentum resummation and leptonic decay*, *JHEP* **12** (2015) 047 [[1507.06937](#)].
- [61] M.A. Ebert, I. Moul, I.W. Stewart, F.J. Tackmann, G. Vita and H.X. Zhu, *Subleading power rapidity divergences and power corrections for q_T* , *JHEP* **04** (2019) 123 [[1812.08189](#)].
- [62] M.A. Ebert and F.J. Tackmann, *Impact of isolation and fiducial cuts on q_T and N-jettiness subtractions*, *JHEP* **03** (2020) 158 [[1911.08486](#)].
- [63] M.A. Ebert, J.K.L. Michel, I.W. Stewart and F.J. Tackmann, *Drell-Yan q_T resummation of fiducial power corrections at N^3LL* , *JHEP* **04** (2021) 102 [[2006.11382](#)].
- [64] G. Ferrera, W.-L. Ju and M. Schönherr, *Zero-bin subtraction and the q_T spectrum beyond leading power*, *JHEP* **04** (2024) 005 [[2312.14911](#)].
- [65] M.A. Ebert, I. Moul, I.W. Stewart, F.J. Tackmann, G. Vita and H.X. Zhu, *Power Corrections for N-Jettiness Subtractions at $\mathcal{O}(\alpha_s)$* , *JHEP* **12** (2018) 084 [[1807.10764](#)].
- [66] R. Boughezal, A. Isgrò and F. Petriello, *Next-to-leading-logarithmic power corrections for N-jettiness subtraction in color-singlet production*, *Phys. Rev. D* **97** (2018) 076006 [[1802.00456](#)].
- [67] R. Boughezal, A. Isgrò and F. Petriello, *Next-to-leading power corrections to $V + 1$ jet production in N-jettiness subtraction*, *Phys. Rev. D* **101** (2020) 016005 [[1907.12213](#)].
- [68] G. Vita, *N^3LO power corrections for 0-jettiness subtractions with fiducial cuts*, *JHEP* **07** (2024) 241 [[2401.03017](#)].
- [69] J. Campbell, T. Neumann and G. Vita, *Projection-to-Born-improved Subtractions at NNLO*, [2408.05265](#).
- [70] L. Magnea, C. Milloy, C. Signorile-Signorile and P. Torrielli, *Strongly-ordered infrared counterterms from factorisation*, *JHEP* **06** (2024) 021 [[2403.11975](#)].
- [71] P. Jakubčík, M. Marcoli and G. Stagnitto, *The parton-level structure of e^+e^- to 2 jets at N^3LO* , *JHEP* **01** (2023) 168 [[2211.08446](#)].

- [72] X. Chen, P. Jakubčík, M. Marcoli and G. Stagnitto, *The parton-level structure of Higgs decays to hadrons at N^3LO* , *JHEP* **06** (2023) 185 [[2304.11180](#)].
- [73] X. Chen, P. Jakubčík, M. Marcoli and G. Stagnitto, *Radiation from a gluon-gluino colour-singlet dipole at N^3LO* , *JHEP* **12** (2023) 198 [[2310.13062](#)].
- [74] K. Melnikov, R. Rietkerk, L. Tancredi and C. Wever, *Double-real contribution to the quark beam function at N^3LO QCD*, *JHEP* **02** (2019) 159 [[1809.06300](#)].
- [75] K. Melnikov, R. Rietkerk, L. Tancredi and C. Wever, *Triple-real contribution to the quark beam function in QCD at next-to-next-to-next-to-leading order*, *JHEP* **06** (2019) 033 [[1904.02433](#)].
- [76] A. Behring, K. Melnikov, R. Rietkerk, L. Tancredi and C. Wever, *Quark beam function at next-to-next-to-next-to-leading order in perturbative QCD in the generalized large- N_c approximation*, *Phys. Rev. D* **100** (2019) 114034 [[1910.10059](#)].
- [77] G. Billis, M.A. Ebert, J.K.L. Michel and F.J. Tackmann, *A toolbox for q_T and 0-jettiness subtractions at N^3LO* , *Eur. Phys. J. Plus* **136** (2021) 214 [[1909.00811](#)].
- [78] D. Baranowski, *NNLO zero-jettiness beam and soft functions to higher orders in the dimensional-regularization parameter ϵ* , *Eur. Phys. J. C* **80** (2020) 523 [[2004.03285](#)].
- [79] M.A. Ebert, B. Mistlberger and G. Vita, *N -jettiness beam functions at N^3LO* , *JHEP* **09** (2020) 143 [[2006.03056](#)].
- [80] D. Baranowski, M. Delto, K. Melnikov and C.-Y. Wang, *Same-hemisphere three-gluon-emission contribution to the zero-jettiness soft function at N^3LO QCD*, *Phys. Rev. D* **106** (2022) 014004 [[2204.09459](#)].
- [81] D. Baranowski, A. Behring, K. Melnikov, L. Tancredi and C. Wever, *Beam functions for N -jettiness at N^3LO in perturbative QCD*, *JHEP* **02** (2023) 073 [[2211.05722](#)].
- [82] G. Bell, B. Dehnadi, T. Mohrmann and R. Rahn, *The NNLO soft function for N -jettiness in hadronic collisions*, *JHEP* **07** (2024) 077 [[2312.11626](#)].
- [83] P. Agarwal, K. Melnikov and I. Pedron, *N -jettiness soft function at next-to-next-to-leading order in perturbative QCD*, *JHEP* **05** (2024) 005 [[2403.03078](#)].
- [84] D. Baranowski, M. Delto, K. Melnikov, A. Pikelner and C.-Y. Wang, *One-loop corrections to the double-real emission contribution to the zero-jettiness soft function at N^3LO in QCD*, *JHEP* **04** (2024) 114 [[2401.05245](#)].
- [85] D. Baranowski, M. Delto, K. Melnikov, A. Pikelner and C.-Y. Wang, *Zero-jettiness soft function to third order in perturbative QCD*, [2409.11042](#).
- [86] D. Baranowski, M. Delto, K. Melnikov, A. Pikelner and C.-Y. Wang, *Triple real-emission contribution to the zero-jettiness soft function at N^3LO in QCD*, [2412.14001](#).
- [87] S. Catani, *The Singular behavior of QCD amplitudes at two loop order*, *Phys. Lett. B* **427** (1998) 161 [[hep-ph/9802439](#)].
- [88] T. Becher and M. Neubert, *On the Structure of Infrared Singularities of Gauge-Theory Amplitudes*, *JHEP* **06** (2009) 081 [[0903.1126](#)].
- [89] T. Becher, G. Bell, C. Lorentzen and S. Marti, *Transverse-momentum spectra of electroweak bosons near threshold at NNLO*, *JHEP* **02** (2014) 004 [[1309.3245](#)].
- [90] F. Caola, M. Delto, H. Frellesvig and K. Melnikov, *The double-soft integral for an arbitrary angle between hard radiators*, *Eur. Phys. J. C* **78** (2018) 687 [[1807.05835](#)].

- [91] S. Catani and M. Grazzini, *Infrared factorization of tree level QCD amplitudes at the next-to-next-to-leading order and beyond*, *Nucl. Phys. B* **570** (2000) 287 [[hep-ph/9908523](#)].
- [92] E. Gardi and L. Magnea, *Factorization constraints for soft anomalous dimensions in QCD scattering amplitudes*, *JHEP* **03** (2009) 079 [[0901.1091](#)].
- [93] M. Delto and K. Melnikov, *Integrated triple-collinear counter-terms for the nested soft-collinear subtraction scheme*, *JHEP* **05** (2019) 148 [[1901.05213](#)].
- [94] R.K. Ellis, W.J. Stirling and B.R. Webber, *QCD and collider physics*, vol. 8, Cambridge University Press (2, 2011), [10.1017/CBO9780511628788](#).
- [95] Z. Bern, V. Del Duca, W.B. Kilgore and C.R. Schmidt, *The infrared behavior of one loop QCD amplitudes at next-to-next-to leading order*, *Phys. Rev. D* **60** (1999) 116001 [[hep-ph/9903516](#)].
- [96] D.A. Kosower and P. Uwer, *One loop splitting amplitudes in gauge theory*, *Nucl. Phys. B* **563** (1999) 477 [[hep-ph/9903515](#)].
- [97] J.M. Campbell and R. Ellis, *An Update on vector boson pair production at hadron colliders*, *Phys.Rev.* **D60** (1999) 113006 [[hep-ph/9905386](#)].