DESY 21–192 DO–TH 21/31 TTP 21–052 RISC Report Series 21–19 SAGEX–21–36 November 2021

## The three-loop polarized singlet anomalous dimensions from off-shell operator matrix elements

J. Blümlein<sup>a</sup>, P. Marquard<sup>a</sup>, C. Schneider<sup>b</sup> and K. Schönwald<sup>c</sup>

<sup>a</sup> Deutsches Elektronen–Synchrotron DESY, Platanenallee 6, 15738 Zeuthen, Germany

<sup>b</sup> Johannes Kepler University Linz, Research Institute for Symbolic Computation (RISC), Altenberger Straße 69, A-4040 Linz, Austria

> <sup>c</sup> Institut für Theoretische Teilchenphysik, Karlsruher Institut für Technologie (KIT) D-76128 Karlsruhe, Germany

#### Abstract

Future high luminosity polarized deep-inelastic scattering experiments will improve both the knowledge of the spin sub-structure of the nucleons and contribute further to the precision determination of the strong coupling constant, as well as, reveal currently yet unknown higher twist contributions in the polarized sector. For all these tasks to be performed, it is necessary to know the QCD leading twist scaling violations of the measured structure functions. Here an important ingredient consists in the polarized singlet anomalous dimensions and splitting functions in QCD. We recalculate these quantities to three-loop order in the M-scheme by using the traditional method of space-like off-shell massless operator matrix elements, being a gauge-dependent framework. Here one obtains the anomalous dimensions without referring to gravitational currents, needed when calculating them using the forward Compton amplitude. We also calculate the non-singlet splitting function  $\Delta P_{qq}^{(2),s,NS}$  and compare the final results to the literature, also including predictions for the region of small values of Bjorken x.

### 1 Introduction

Polarized deep-inelastic scattering allows to reveal the spin and angular momentum structure of nucleons [1]. At sufficiently high scales of the virtuality of the exchanged electro-weak current  $Q^2 \gg 1 \text{ GeV}^2$  and for not too small or too large values of Bjorken  $x = Q^2/(2p.q)$  [2] the twist-2 contributions dominate over the higher twist contributions [3] and target mass corrections [4]. Because of their strong variation, also the detailed control of the QED radiative corrections is required [5].

The scaling violations of the polarized deep-inelastic structure functions at the level of leading twist  $\tau = 2$  are determined by the anomalous dimensions of the composite quark and gluon operators [6,7] through the evolution of the polarized parton densities and the massless and massive Wilson coefficients. A major goal in measuring the polarized deep-inelastic scattering process to high precision, e.g. at the EIC in the future [8] consist in the detailed measurement of the polarized parton distribution functions at a reference scale  $Q_0^2$  together with the correlated measurement of the strong coupling constant  $a_s(M_Z^2) = \alpha_s(M_Z^2)/(4\pi)$  [9]. One of the important ingredients to this is the knowledge of the polarized singlet anomalous dimensions to highest possible order. The polarized non-singlet anomalous dimensions are known to 3-loop order [10, 11] and the polarized singlet anomalous dimensions were calculated to first [12-16], second [17-19]and third order in [20,21], using different computation techniques. In Ref. [20] the calculation has been performed using the forward Compton amplitude, which required reference to gravitational subsidiary currents for the gluonic sector. A direct calculation of the contributions  $\propto T_F^1$  of the three-loop anomalous dimensions is possible using massive on-shell operator matrix elements [21], without reference to gravitational currents. However, this method only allows to calculate the anomalous dimensions  $\Delta \gamma_{qq}^{(2),\text{PS}}$  and  $\Delta \gamma_{qg}^{(2)}$  in complete form.

In the present paper we use the traditional approach of massless off-shell operator matrix elements (OMEs), allowing the direct calculation, as before for the non-singlet case and the three-loop anomalous dimensions of transversity in Ref. [11]. However, for this method the computational framework is gauge dependent. Still the anomalous dimensions in minimal subtraction schemes remain gauge invariant. Specific physical projectors have to be used, cf. also [19]. In the polarized case, unlike the unpolarized case [22], there is no mixing with the so-called alien operators [19, 22–32]. The main goal of the present paper is the re–calculation of all three–loop polarized singlet anomalous dimensions by a different method, being the first re–calculation of the polarized anomalous dimensions  $\Delta \gamma_{gq}^{(2)}$  and  $\Delta \gamma_{gg}^{(2)}$ . To some extent different steps in the computation technology are the same as in the non–singlet case, cf. [11] which will be referred to this paper and we only will describe new additional elements in the present paper.<sup>2</sup> At the phenomenological side we mention that, unlike the singlet case, the flavor non–singlet description of the structure functions  $g_{1,2}^{NS}(x, Q^2)$  is well explored to three–loop order, also including the heavy flavor contributions [33–35].

The paper is organized as follows. In Section 2 we derive the structure of the physical part of the flavor singlet polarized unrenormalized off-shell OMEs to three-loop order. From their pole terms of  $O(1/\varepsilon)$  one can extract the singlet anomalous dimensions. We work in the Larin scheme [19,36] and perform finally the transformation to the M-scheme [19,20]. We also present the calculation details. Here a central method being applied is the method of arbitrary high moments [37]. In Section 3 we calculate also the polarized non-singlet anomalous dimension  $\Delta \gamma_{qq}^{(2),s,NS}$ , cf. also [38], and present the polarized singlet anomalous dimensions in Section 4.

<sup>&</sup>lt;sup>1</sup>The color factors are defined in [11], below Eq. (22).

<sup>&</sup>lt;sup>2</sup>This seems to be necessary to avoid critique by word processing codes, like e.g. iThenticate, even on the expense to not provide more comprehensive information to the reader in the same publication, which we regret.

Comparisons to the literature are given in Section 5, including a discussion of the small x limit. Section 6 contains the conclusions and a new Feynman rule is presented in Appendix A.

## 2 The unrenormalized polarized operator matrix elements and details of the calculation

The massless off-shell singlet OMEs are defined as expectation values of the local operators

$$O_{q;\mu_1...\mu_N}^{S,5} = i^{N-1} \mathbf{S} \left[ \overline{\psi} \gamma_5 \gamma_{\mu_1} D_{\mu_2} ... D_{\mu_N} \psi \right] - \text{ trace terms}, \tag{1}$$

$$O_{g;\mu_1\dots\mu_N}^{S,5} = 2i^{N-2} \mathbf{S} \operatorname{tr} \left[ \frac{1}{2} \varepsilon^{\mu_1 \alpha \beta \gamma} F^a_{\beta \gamma} D_{\mu_2} \dots D_{\mu_{N-1}} F^{\mu_N}_{\alpha,a} \right] - \text{ trace terms}$$
(2)

between quark (antiquark)  $\psi(\bar{\psi})$  and gluonic states  $F^a_{\mu_1\alpha}$  of space–like momentum  $p, p^2 < 0$ . The OMEs are given by

$$\Delta \hat{A}_{ij} = \langle q(p), j | O_i^{\mathrm{S},(5)} | q(p), j \rangle.$$
(3)

For the other definitions, here and in the following, we refer the reader to Ref. [11].

The Feynman rules of both QCD and of the local composite operators are given in [21,39,40] and have to be extended by that of the polarized local five–gluon vertex, given in Appendix A.

We use the Larin scheme [36]<sup>3</sup> to describe  $\gamma^5$  in  $D = 4 + \varepsilon$ -dimensions and express  $\gamma^5$  by

$$\gamma^5 = \frac{i}{24} \varepsilon_{\mu\nu\rho\sigma} \gamma^{\mu} \gamma^{\nu} \gamma^{\rho} \gamma^{\sigma}, \qquad (4)$$

$$\not\Delta \gamma^5 = \frac{i}{6} \varepsilon_{\mu\nu\rho\sigma} \Delta^{\mu} \gamma^{\nu} \gamma^{\rho} \gamma^{\sigma}.$$
(5)

The Levi-Civita symbols are now contracted in D dimensions,

$$\varepsilon_{\mu\nu\rho\sigma}\varepsilon^{\alpha\lambda\tau\gamma} = -\text{Det}[g^{\beta}_{\omega}], \quad \beta = \alpha, \lambda, \tau, \gamma; \quad \omega = \mu, \nu, \rho, \sigma.$$
 (6)

The Larin scheme is one of the consistent calculation schemes in the polarized case. We will later transform to another scheme, the M-scheme. It is needless to say that the calculation of any observable requires to calculate also the Wilson coefficients in the same scheme, which implies that the extracted twist-2 parton density functions are obtained in this scheme, despite being universal w.r.t. the scattering processes in which they are used. Any analytic continuation of  $\gamma_5$ or the Levi-Civita symbol to D dimensions violates Ward identities. One way to restore these consists in the calculation of scheme-invariant quantities.

The polarized operator matrix elements have the representation

$$\Delta \hat{A}_{qq}^{\text{PS}} = \left[ \gamma_5 \not\Delta \Delta \hat{A}_{qq}^{\text{PS,phys}} + \gamma_5 \not p \frac{\Delta \cdot p}{p^2} \Delta \hat{A}_{qq}^{\text{PS,EOM}} \right] (\Delta \cdot p)^{N-1}$$
(7)

$$\Delta \hat{A}_{qg} = \varepsilon_{\mu\nu\alpha\beta} \Delta^{\alpha} p^{\beta} \frac{1}{\Delta p} \Delta \hat{A}_{qg}^{\text{phys}} (\Delta p)^{N-1}$$
(8)

$$\Delta \hat{A}_{gq} = \left[ \gamma_5 \not\Delta \Delta \hat{A}_{gq}^{\text{phys}} + \gamma_5 \not p \frac{\Delta p}{p^2} \hat{A}_{gq}^{\text{EOM}} \right] (\Delta p)^{N-1}$$
(9)

<sup>3</sup>See Refs. [41] for other schemes and Refs. [42] concerning the finite renormalization in different approaches.

$$\Delta \hat{A}_{gg,\mu\nu} = \varepsilon_{\mu\nu\alpha\beta} \Delta^{\alpha} p^{\beta} \frac{1}{\Delta p} \Delta \hat{A}_{gq}^{\text{phys}} (\Delta p)^{N-1}, \qquad (10)$$

with  $\Delta$  obeying  $\Delta \Delta = 0$ . It is convenient to introduce the following projectors on the physical part (phys)

$$\Delta \hat{A}_{iq}^{\text{phys}} = -\frac{1}{4(D-2)(D-3)} \varepsilon_{\mu\nu\rho\sigma} p^{\rho} \Delta^{\sigma} \text{tr} \left[ p \gamma^{\mu} \gamma^{\nu} \Delta \hat{A}_{iq} \right] (\Delta . p)^{-N-1}$$
(11)

$$-\frac{p^2}{4(D-2)(D-3)}(\Delta .p)^{-N-2}\varepsilon_{\mu\nu\rho\sigma}p^{\rho}\Delta^{\sigma}\mathrm{tr}\left[\not{\Delta}\gamma^{\mu}\gamma^{\nu}\Delta\hat{A}_{iq}\right],\tag{12}$$

$$\Delta \hat{A}_{ig}^{\text{phys}} = \frac{1}{(D-2)(D-3)} \varepsilon_{\mu\nu\rho\sigma} \Delta^{\rho} p^{\sigma} (\Delta . p)^{-N-1} \Delta \hat{A}_{ig}^{\mu\nu}, \qquad (13)$$

with i = q, g. Only these contributions are necessary for the calculation of the singlet anomalous dimensions. In Mellin N space the following representation holds

$$\Delta \hat{A}_{ij} = \delta_{ij} + \sum_{k=1}^{\infty} \hat{a}^k S^k_{\varepsilon} \left(\frac{-p^2}{\mu^2}\right)^{\varepsilon k/2} \Delta \hat{A}^{(k)}_{ij}, \qquad (14)$$

with the spherical factor  $S_{\varepsilon}$  given in Eq. (13) of [11]. The bare coupling constant  $\hat{a}$  and the gauge parameter  $\xi$  are defined in Eqs. (15–22) of [11], working in the  $R_{\hat{\xi}}$  gauge, by using relations from [43–47].

In Mellin N space the Z-factor of a local singlet operator reads [40]

$$Z_{ij}^{S} = \delta_{ij} + a \frac{\Delta \gamma_{ij}^{(0)}}{\varepsilon} + a^{2} \left[ \frac{1}{\varepsilon^{2}} \left( \frac{1}{2} \Delta \gamma_{il}^{(0)} \Delta \gamma_{lj}^{(0)} + \beta_{0} \Delta \gamma_{ij}^{(0)} \right) + \frac{1}{2\varepsilon} \Delta \gamma_{ij}^{(1)} \right] \\ + a^{3} \left[ \frac{1}{\varepsilon^{3}} \left( \frac{1}{6} \Delta \gamma_{il}^{(0)} \Delta \gamma_{lk}^{(0)} \Delta \gamma_{kj}^{(0)} + \beta_{0} \Delta \gamma_{il}^{(0)} \Delta \gamma_{lj}^{(0)} + \frac{4}{3} \beta_{0}^{2} \Delta \gamma_{ij}^{(0)} \right) \right] \\ + \frac{1}{\varepsilon^{2}} \left( \frac{1}{6} \Delta \gamma_{il}^{(1)} \Delta \gamma_{lj}^{(0)} + \frac{1}{3} \Delta \gamma_{il}^{(0)} \Delta \gamma_{lj}^{(1)} + \frac{2}{3} \beta_{0} \Delta \gamma_{ij}^{(1)} + \frac{2}{3} \beta_{1} \Delta \gamma_{ij}^{(0)} \right) + \frac{1}{3\varepsilon} \Delta \gamma_{ij}^{(2)} \right].$$
(15)

In (15) the terms  $\Delta \gamma_{ij}^{(k)}$ , k = 0, 1, 2, ... denote the expansion coefficients of the anomalous dimension

$$\Delta \gamma_{ij} = \sum_{k=1}^{\infty} a^k \Delta \gamma_{ij}^{(k-1)}.$$
(16)

The partly renormalized polarized singlet OMEs,  $\Delta \tilde{A}_{ij}^{\text{phys}}$ , read

$$\Delta \tilde{A}_{ij}^{\text{phys}} = 1 + a \left[ \frac{a_{ij}^{(1,-1)}}{\varepsilon} + a_{ij}^{(1,0)} + a_{ij}^{(1,1)} \varepsilon \right] + a^2 \left[ \frac{a_{ij}^{(2,-2)}}{\varepsilon^2} + \frac{a_{ij}^{(2,-1)}}{\varepsilon} + a_{ij}^{(2,0)} \right] + a^3 \left[ \frac{a_{ij}^{(3,-3)}}{\varepsilon^3} + \frac{a_{ij}^{(3,-2)}}{\varepsilon^2} + \frac{a_{ij}^{(3,-1)}}{\varepsilon} \right].$$
(17)

The expansion coefficients  $a_{ij}^{(k,l)}$  are gauge dependent in general. The renormalized OMEs in the Larin scheme are given by

$$\Delta A_{ij}^{\text{phys}} = (\Delta Z_{ik}^S)^{-1} \Delta \tilde{A}_{kj}^{\text{phys}}, \qquad (18)$$

expanded to  $O(a^3)$  and setting  $S_{\varepsilon} = 1$ . The anomalous dimensions are iteratively extracted from the  $1/\varepsilon$  pole terms and the other expansion coefficients  $a_{ij}^{(k,l)}$  are given in Ref. [48]. The anomalous dimensions in the M–scheme  $[19, 20]^4$  are obtained by the following transfor-

mations

$$\Delta \gamma_{ij}^{(0),\mathrm{M}} = \Delta \gamma_{ij}^{(0),\mathrm{L}} \tag{19}$$

$$\Delta \gamma_{qq}^{(1),\text{NS,M}} = \Delta \gamma_{qq}^{(1),\text{NS,L}} + 2\beta_0 z_{qq}^{(1)},$$
(20)

$$\gamma_{qq}^{(1),\text{PS,M}} = \Delta \gamma_{qq}^{(1),\text{PS,L}}, \qquad (21)$$

$$\Delta \gamma_{qg}^{(1),M} = \Delta \gamma_{qg}^{(1),L} + \Delta \gamma_{qg}^{(0)} z_{qq}^{(1)}, \qquad (22)$$

$$\Delta \gamma_{gq}^{(1),M} = \Delta \gamma_{gq}^{(1),L} - \Delta \gamma_{gq}^{(0)} z_{qq}^{(1)},$$
(23)
$$\Delta z_{qq}^{(1),M} = \Delta z_{qq}^{(1),L}$$
(24)

$$\Delta \gamma_{gg}^{(2),\mathrm{NL}} = \Delta \gamma_{gg}^{(2),\mathrm{L}}. \tag{24}$$

$$\omega^{(2),\mathrm{NS,M}} = \Delta \gamma^{(2),\mathrm{NS,L}} = 2\beta \left( \left( z^{(1)} \right)^2 - 2z^{(2),\mathrm{NS}} \right) + 2\beta z^{(1)}. \tag{25}$$

$$\Delta \gamma_{qq}^{(2),\text{NS,M}} = \Delta \gamma_{qq}^{(2),\text{NS,L}} - 2\beta_0 \left( \left( z_{qq}^{(1)} \right)^2 - 2z_{qq}^{(2),\text{NS}} \right) + 2\beta_1 z_{qq}^{(1)}, \tag{25}$$
$$\Delta \gamma_{qq}^{(2),\text{PS,M}} = \Delta \gamma_{qq}^{(2),\text{PS,L}} + 4\beta_0 z_{qq}^{(2),\text{PS}}. \tag{26}$$

$$\Delta \gamma_{qq}^{(2),M} = \Delta \gamma_{qq}^{(2),L} + \Delta \gamma_{qq}^{(1),M} z_{qq}^{(1)} + \Delta \gamma_{qq}^{(0)} \left( z_{qq}^{(2)} - \left( z_{qq}^{(1)} \right)^2 \right), \tag{27}$$

$$\Delta \gamma_{gq}^{(2),M} = \Delta \gamma_{gq}^{(2),L} - \Delta \gamma_{gq}^{(1),M} z_{qq}^{(1)} - \Delta \gamma_{gq}^{(0)} z_{qq}^{(2)},$$
(28)

$$\Delta \gamma_{gg}^{(2),\mathrm{M}} = \Delta \gamma_{gg}^{(2),\mathrm{L}}, \tag{29}$$

with [19]

 $\Delta$ 

$$z_{qq}^{(1)} = -\frac{8C_F}{N(N+1)},$$
(30)  

$$z_{qq}^{(2),NS} = C_F T_F N_F \frac{16(-3-N+5N^2)}{9N^2(1+N)^2} + C_A C_F \left\{ -\frac{4R_1}{9N^3(1+N)^3} - \frac{16}{N(1+N)} S_{-2} \right\}$$

$$+ C_F^2 \left\{ \frac{8(2+5N+8N^2+N^3+2N^4)}{N^3(1+N)^3} + \frac{16(1+2N)}{N^2(1+N)^2} S_1 + \frac{16}{N(1+N)} S_2 + \frac{32}{N(1+N)} S_{-2} \right\},$$
(31)

$$z_{qq}^{(2),\text{PS}} = 8C_F T_F N_F \frac{(N+2)(1+N-N^2)}{N^3(N+1)^3},$$
(32)

$$z_{qq}^{(2)} = z_{qq}^{(2),\text{NS}} + z_{qq}^{(2),\text{PS}}$$
(33)

and

 $R_1 = 103N^4 + 140N^3 + 58N^2 + 21N + 36.$ (34)

Let us now turn now to the technical aspects of the present calculation, which has been widely automated and the corresponding chain of programs has been described to some extent in our previous paper [11]. A main point concerns the observation of the current crossing relations in the present case, cf. [6,49] and also Ref. [48], projecting onto the contributions of the odd integer moments

$$\sum_{N=0}^{\infty} (\Delta . k)^N \left( t^N - (-t)^N \right) \to \left[ \frac{1}{1 - \Delta . k \ t} - \frac{1}{1 + \Delta . k \ t} \right].$$

$$(35)$$

<sup>&</sup>lt;sup>4</sup>We reproduce these transformations here, because they are instrumental importance for the understanding of the present calculation, which is performed in the Larin scheme.

Through this one obtains a quadratic dependence on the resummation variable t. An expansion in t leads to the moments again.

Our chain of programs, from the generation of Feynman diagrams to the final result, the polarized anomalous dimensions at three-loop order includes the packages QGRAF, FORM, Color, EvaluateMultiSums,Crusher,SolveCoupledSystems,Guess,Sage,Sigma,HarmoncisSums [40, 50-71]. More details were given in Ref. [11]. We determine the contributions to the anomalous dimensions due to the different color and multiple zeta values [70] individually. In the unfolding process also higher order terms  $a_{ij}^{(k)}$  in the dimensional parameter  $\varepsilon$  are needed to be calculated, cf. Ref. [48]. We use the integration-by-parts relations [54, 72] to reduce the problem to master integrals. The method of arbitrary high moments [37] allows to find the needed difference equations, which are then solved using algorithms in difference ring theory [73]. To calculate initial values for the solution we use relations given in [74, 75].

We turn now to some statistical characteristics of the present calculation. In the polarized flavor singlet case 125 irreducible diagrams contribute for  $\Delta A_{qq}^{(3),\text{PS}}$ , 1101 for  $\Delta A_{qq}^{(3)}$ , 400 for  $\Delta A_{gq}^{(3)}$ , and 1598 for  $\Delta A_{gg}^{(3)}$ . For comparison the number of diagrams for  $\Delta A_{qq}^{(3),\text{NS}}$  amounts to 559. Finally, 24 diagrams contribute to the part of the forward Compton amplitude, from which  $\Delta \gamma_{qq}^{(2),\text{s,NS}}$  is extracted. The total number of irreducible diagrams in the polarized singlet case at three–loop order is larger by a factor of ~ 22 than in the two–loop order. The reducible diagrams are accounted for by wave–function renormalization [44, 46, 47], decorating the OMEs at lower order in the coupling constant [19, 48].

To calculate the anomalous dimensions  $\Delta \gamma_{ij}^{(2)}$  we generated 3000 odd moments. It turns out that the determination of the largest recurrence requires 462 moments for  $\Delta \gamma_{qq}^{(2),PS}$ , 989 moments for  $\Delta \gamma_{qg}^{(2)}$ , 1035 moments for  $\Delta \gamma_{gq}^{(2)}$ , 1568 moments for  $\Delta \gamma_{gg}^{(2)}$ .

In this way the anomalous dimensions are obtained. The largest difference equation contributing has order o = 16 and degree d = 304. These numbers are of the order obtained in the non-singlet case in Ref. [57], where the largest difference equation contributing had order o =16 and degree d = 192, and required 1079 moments. A moment based test-run for the polarized anomalous dimensions, like in the unpolarized case, has not been performed.

The overall computation time using the automated chain of codes described amounted to about 18 days of CPU time on Intel(R) Xeon(R) CPU E5-2643 v4 processors. Again we have only retained the first power of the gauge parameter to have a first check on the renormalization.

The anomalous dimensions,  $\Delta \gamma_{ij}$ , can all be expressed by harmonic sums [62, 63], defined in Eq. (34) of Ref. [11]. Their Mellin inversion to the splitting functions  $\Delta P_{ij}(z)$ 

$$\Delta \gamma_{ij}(N) = -\int_0^1 dz z^{N-1} \Delta P_{ij}(z)$$
(36)

is obtained using routines of the packages HarmonicSums and are expressed by harmonic polylogarithms [64], which are defined in Eqs. (36,37) of Ref. [11].

Because of the different requirements on the respective integrals, one distinguishes three contributions to the individual splitting functions in z-space

$$\Delta P(z) = \Delta P^{\delta}(z) + \Delta P^{\text{plu}}(z) + \Delta P^{\text{reg}}(z), \qquad (37)$$

which are defined in Eqs. (38,39) of Ref. [11].

Also here we reduce the expressions to the algebraic basis, cf. [69], which has the advantage that only the minimal set has to be calculated in numerical applications [76]. We will not present the splitting function in explicit form, since the expressions are rather lengthy. They are given in computer–readable form in an attachment to the present paper. The polarized singlet anomalous dimensions are given in Section 4.

# 3 The polarized non–singlet anomalous dimension $\Delta\gamma^{(2),s, ext{NS}}_{ ext{qq}}$

In our previous paper [11] we had not yet calculated the non-singlet anomalous dimension  $\Delta \gamma_{qq}^{(2),s,NS}$ , which emerges from three–loop order onward. It is best calculated using the vectoraxialvector interference term in the forward Compton amplitude, corresponding to the associated structure function  $g_5^-(x, Q^2)$ , Ref. [49], corresponding to the difference in case of  $W^-$  and  $W^+$ charged current scattering. Due to its crossing relations it has even moments. This anomalous dimension has been calculated previously in Ref. [38].

The corresponding gauge boson vertex is parameterized by

$$i\left[v\gamma_{\mu} + a\frac{i}{6}\varepsilon_{\mu\nu_{1}\nu_{2}\nu_{3}}\gamma^{\nu_{1}}\gamma^{\nu_{2}}\gamma^{\nu_{3}}\right]$$
(38)

and we consider the current interference term  $\propto a \cdot v$ . The projectors for the massless external quark lines of momentum p and boson lines corresponding to a tensor of rank two read

$$P^{q} = \frac{1}{4} \operatorname{tr} \left[ \frac{i}{6} \varepsilon_{p\rho_{1}\rho_{2}\rho_{3}} \gamma^{\rho_{1}} \gamma^{\rho_{2}} \gamma^{\rho_{3}} \dots \right]; \qquad P^{b}_{\mu\nu} = -\frac{g_{\mu\nu}}{D-1}.$$
(39)

The forward Compton amplitude depends on the invariants  $Q^2 = -q^2$  and  $p.q = Q^2/(2z) \equiv (Q^2/2)y$ . The diagrams can be represented as formal power series in y

$$F(y) = \sum_{N=0}^{\infty} f(N)y^N,$$
(40)

cf. e.g. [11], Sect. 2. The IBP-reduction in this case leads to three families and 101 master integrals in total. We consider the differential equations for the individual master integrals  $M_k(y,\varepsilon)$ , which are computed using the method described in [77]. After insertion of the master integrals into the amplitude and subsequent expansion in the dimensional parameter  $\varepsilon$ , the anomalous dimension can be determined by using the command GetMoment [F[y],y,N] of the package HarmonicSums from the pole term  $O(1/\varepsilon)$  of the forward Compton amplitude. We obtain

$$\Delta \gamma_{\rm NS}^{(2),\rm s} = -16 \frac{1 + (-1)^N}{2} N_F \frac{d^{abc} d_{abc}}{N_c} \left[ \frac{1}{N^2} + \frac{2Q_1}{N^4 (1+N)^4} S_1 + \frac{2(2+3N+3N^2)}{N^2 (1+N)^2} \right] \\ \times [S_3 - 2S_{-3} + 4S_{-2,1}] + \left( \frac{4(2+4N+4N^2+N^3+N^4)}{N^3 (1+N)^3} + \frac{8(-1+N)(2+N)}{N^2 (1+N)^2} S_1 \right) \\ \times S_{-2} \right]$$

$$(41)$$

with

$$Q_1 = 3N^6 + 8N^5 - N^4 - 14N^3 - 29N^2 - 21N - 6.$$
(42)

The corresponding splitting function reads

$$\Delta P_{\rm NS}^{(2),\rm s} = 16N_F \frac{d^{abc}d_{abc}}{N_c} \left\{ (1+x) \left[ -4H_0H_{-1} \left( -4+5H_{-1} \right) + 8H_{0,-1} \left( -2+5H_{-1} \right) + 18H_{-1}H_0^2 + \left( -2H_0^2 + 32H_{-1} \right)H_{0,1} + 8H_0H_{0,0,1} - 32H_{0,1,-1} - 32H_{0,-1,1} \right) \right\}$$

$$-40H_{0,-1,-1} - (52H_{-1} + 8H_{0,1})\zeta_{2} + (1-x) \left[ 24H_{1} - H_{0}^{2}H_{1} + 2H_{0}H_{0,1} + 4H_{0}^{2}H_{0,-1} + 8H_{0,-1}^{2} - 8H_{0}H_{0,0,-1} - 16H_{0}H_{0,-1,-1} - (20H_{1} + 8H_{0,-1})\zeta_{2} \right] - (1 + 24x)H_{0} - 4xH_{0}^{2} - 6xH_{0}^{3} + \frac{1}{3}xH_{0}^{4} + 2(5 + 4x)H_{0,1} - 28H_{0}H_{0,-1} + 4(1 - 9x)H_{0,0,1} + 4(5 - 9x)H_{0,0,-1} + (2(-5 + 4x) + 2(-3 + 37x)H_{0} - 2(3 + 5x)H_{0}^{2})\zeta_{2} + (2(3 + 44x) - 16xH_{0})\zeta_{3} + 2(5 + 3x)\zeta_{2}^{2} \right\}.$$
(43)

Here  $\zeta_k$  denotes the Riemann  $\zeta$ -function at integer argument  $k \geq 2$  and the color factor is normalized in the present case to  $d^{abc}d_{abc}/N_c = 5/18$  in QCD. The leading small z contribution of  $\Delta P_{\rm NS}^{(2),\rm s}$  is  $\propto \ln^2(z)$ . This behaviour is, however, not dominant in kinematic regions being accessible at present. The asymptotic behaviour reaches the complete function up to 10% below  $z \sim 10^{-11}$  only. One more logarithmic order allows a description below  $z \sim 10^{-3}$ .

#### 4 The polarized singlet anomalous dimensions

In the following we use the minimal representations in terms of the contributing harmonic sums and harmonic polylogarithms by applying the algebraic relations [69] between the harmonic sums and the harmonic polylogarithms. The polarized singlet anomalous dimensions can be represented by the following 23 harmonic sums up to weight w = 5 to three–loop order

$$\left\{ S_{-5}, S_{-4}, S_{-3}, S_{-2}, S_1, S_2, S_3, S_4, S_5, S_{-4,1}, S_{-3,1}, S_{-2,1}, S_{-2,2}, S_{-2,3}, S_{2,-3}, S_{2,1}, S_{3,1}, S_{-3,1,1}, S_{-2,1,1}, S_{-2,1,1}, S_{-2,2,1}, S_{2,1,-2}, S_{2,1,1}, S_{-2,1,1,1} \right\}.$$

$$(44)$$

This number is further reduced using also the structural relations [71,78] to at least 15 sums. In the present case the 10 sums

$$\left\{S_{1}, S_{2,1}, S_{-2,1}, S_{-3,1}, S_{-4,1}, S_{2,1,1}, S_{-2,1,1}, S_{2,1,-2}, S_{-3,1,1}, S_{-2,1,1,1}\right\}$$
(45)

suffice.

The splitting functions in z-space depend on the 26 harmonic polylogarithms

The harmonic sums are defined at the odd integers in the first place and the analytic continuation to  $N \in \mathbb{C}$  is performed from there, [71, 79].

We obtain the following expressions for the polarized singlet anomalous dimensions in Mellin-N space, using the shorthand notation  $S_{\vec{a}}(N) \equiv S_{\vec{a}}$  from one- to three-loop order. Here we dropped the prefactor  $\frac{1}{2}(1-(-1)^N)$ .

$$\Delta \gamma_{qq}^{(0)} = C_F \left[ -\frac{2(2+3N+3N^2)}{N(1+N)} + 8S_1 \right], \tag{47}$$

$$\Delta \gamma_{qg}^{(0)} = -T_F N_F \frac{8(N-1)}{N(1+N)}, \tag{48}$$

$$\Delta \gamma_{gq}^{(0)} = -C_F \frac{4(2+N)}{N(1+N)},\tag{49}$$

$$\Delta \gamma_{gg}^{(0)} = T_F N_F \frac{8}{3} + C_A \left[ -\frac{2(24 + 11N + 11N^2)}{3N(1+N)} + 8S_1 \right], \tag{50}$$

$$\Delta \gamma_{qq}^{(1),\text{PS}} = C_F T_F N_F \frac{16(2+N)(1+2N+N^3)}{N^3(1+N)^3}, \tag{51}$$

$$\Delta \gamma_{qg}^{(1)} = C_F T_F N_F \left[ -\frac{8(-1+N)\left(2-N+10N^3+5N^4\right)}{N^3(1+N)^3} + \frac{32(N-1)}{N^2(1+N)}S_1 - \frac{16(N-1)}{N(1+N)}[S_1^2 - S_2] \right] + C_A T_F N_F \left[ -\frac{16P_{20}}{N^3(1+N)^3} - \frac{64S_1}{N(1+N)^2} + \frac{16(N-1)}{N(1+N)}[S_1^2 + S_2 + 2S_{-2}] \right],$$
(52)

$$\Delta \gamma_{gq}^{(1)} = C_F \left[ T_F N_F \left( \frac{32(2+N)(2+5N)}{9N(1+N)^2} - \frac{32(2+N)}{3N(1+N)} S_1 \right) + C_A \left( -\frac{8P_{33}}{9N^3(1+N)^3} \right. \\ \left. + \frac{8(12+22N+11N^2)S_1}{3N^2(1+N)} - \frac{8(2+N)S_1^2}{N(1+N)} + \frac{8(2+N)S_2}{N(1+N)} + \frac{16(2+N)S_{-2}}{N(1+N)} \right) \right] \\ \left. + C_F^2 \left[ \frac{4(2+N)(1+3N)(-2-N+3N^2+3N^3)}{N^3(1+N)^3} - \frac{8(2+N)(1+3N)}{N(1+N)^2} S_1 \right. \\ \left. + \frac{8(2+N)}{N(1+N)} [S_1^2 + S_2] \right],$$
(53)

$$\begin{split} \Delta\gamma_{gg}^{(1)} &= C_F T_F N_F \frac{8P_{40}}{N^3(1+N)^3} + C_A^2 \left[ -\frac{4P_{46}}{9N^3(1+N)^3} + \left( \frac{8P_{17}}{9N^2(1+N)^2} - 32S_2 \right) S_1 \right. \\ &+ \frac{64}{N(1+N)} S_2 - 16S_3 + \left( \frac{64}{N(1+N)} - 32S_1 \right) S_{-2} - 16S_{-3} + 32S_{-2,1} \right] \\ &+ C_A T_F N_F \left[ \frac{32P_5}{9N^2(1+N)^2} - \frac{160}{9}S_1 \right], \end{split}$$
(54)  
$$\Delta\gamma_{qq}^{(2),\text{PS}} &= C_F \left[ T_F^2 N_F^2 \left[ -\frac{64(2+N)P_{30}}{27N^4(1+N)^4} + \frac{64(2+N)\left(6+10N-3N^2+11N^3\right)}{9N^3(1+N)^3} S_1 \right. \\ &\left. -\frac{32(N-1)(2+N)}{3N^2(1+N)^2} [S_1^2 + S_2] \right] + C_A T_F N_F \left[ \frac{8P_9}{3N^3(1+N)^3} S_1^2 + \frac{8P_{10}}{3N^3(1+N)^3} S_2 \right] \end{split}$$

$$\begin{split} &+ \frac{16P_{61}}{27N^5(1+N)^5} + \left(-\frac{16P_{51}}{9N^4(1+N)^4} + \frac{32(N-1)(2+N)}{N^2(1+N)^2}S_2\right)S_1 \\ &- \frac{32(N-1)(2+N)}{3N^2(1+N)^2}S_1^3 + \frac{16(-58+23N+23N^2)}{3N^2(1+N)^2}S_3 + \left(-\frac{32P_1}{N^3(1+N)^3} + \frac{64(N-1)(2+N)}{N^2(1+N)^2}S_1\right)S_2 + \frac{32(-10+7N+7N^2)}{N^2(1+N)^2}S_3 - \frac{64(N-1)(2+N)}{N^2(1+N)^2}S_{2,1} \\ &- \frac{64(-2+3N+3N^2)}{N^2(1+N)^2}S_{-2,1} - \frac{192(N-1)(2+N)}{N^2(1+N)^2}S_3\right] \\ &+ C_F^2 T_F N_F \left[-\frac{16(2+N)P_{54}}{N^5(1+N)^5} + \left(\frac{16(2+N)P_{57}}{N^4(1+N)^4} - \frac{32(N-1)(2+N)}{N^2(1+N)^2}S_2\right)S_1 \\ &- \frac{8(N-1)(2+N)(2+3N+3N^2)}{N^3(1+N)^3}S_1^2 + \frac{32(N-1)(2+N)}{3N^2(1+N)^2}S_3 \\ &- \frac{8(2+N)(14+23N+11N^3)}{N^3(1+N)^3}S_2 - \frac{224(N-1)(2+N)}{3N^2(1+N)^2}S_3 \\ &+ \frac{64(N-1)(2+N)}{N^3(1+N)^3}S_2 - \frac{224(N-1)(2+N)}{3N^2(1+N)^2}S_3 \\ &+ \frac{64(N-1)(2+N)}{N^2(1+N)^2}S_{2,1} + \frac{192(N-1)(2+N)}{N^2(1+N)^2}\zeta_3 \right], \quad (55) \\ \Delta\gamma_{qgg}^{(2)} &= C_F T_F^2 N_F^2 \left[\frac{4P_{64}}{27N^5(1+N)^5} + \left(-\frac{32(-24+4N+47N^2)}{27N^2(1+N)} - \frac{32(N-1)}{3N(1+N)}S_2\right)S_1 \\ &+ \frac{32(N-1)(3+10N)}{N^2(1+N)}S_1^2 - \frac{32(N-1)}{9N(1+N)}S_1^3 + \frac{32(5N-1)}{3N^2(1+N)}S_2 + \frac{320(N-1)}{9N(1+N)}S_3 \right] \\ &+ C_A C_F T_F N_F \left[\frac{8P_{29}}{3N^3(1+N)^3}S_2 + \frac{27N^5(1+N)^5(2+N)}{3N^2(1+N)^2} + \left(-\frac{384(N-1)}{N(1+N)}S_2\right) \\ &+ \frac{16P_{90}}{1N(1+N)}S_1 + \frac{16(75+14N+18N^2+N^3)S_2}{3N^2(1+N)^2} + \frac{400(N-1)}{3N(1+N)}S_3 \\ &- \frac{192(N-1)}{N(1+N)}S_3 \right)S_1 + \left(-\frac{8P_{25}}{9N^3(1+N)^3} + \frac{160(N-1)}{N(1+N)}S_2 \right)S_1^2 \\ &+ \frac{16(3-31N-18N^2+10N^3)}{9N^2(1+N)^2}S_1^3 + \frac{32(N-1)}{3N(1+N)}S_1^4 - \frac{64(N-1)}{N(1+N)}S_2 \\ &- \frac{16(N-1)(2+N)(-1+3N)}{9N^2(1+N)^2}S_1 + \frac{162(N-1)(4-N+N^2)}{N^2(1+N)^2}S_{2,1} \\ &- \frac{32(N-1)(2+N)(-1+3N)}{N^2(1+N)^2}S_3 + \frac{96(N-1)(4-N+N^2)}{N(1+N)}S_{2,2} \\ &+ \frac{32(N-1)(2+N)(-1+3N)}{N^2(1+N)^2}S_3 + \frac{96(N-1)}{N(1+N)}S_{2,2} \\ &+ \frac{32(N-1)(2+N)(-1+3N)}{N^2(1+N)^2}S_3 + \frac{96(N-1)}{N(1+N)}S_2 \\ &- \frac{32P_{24}}{N^3(1+N)^3}S_3 + \frac{96(N-1)(4+N+N^2)}{N^2(1+N)^2}S_{2,3} \\ &+ \frac{160(N-1)}{N(1+N)}S_4 + \frac{64(N-1)}{N(1+N)}S_{3,1} - \frac{128(N-1)}{N^2(1+N)^2}S_{2,2} \\ &+ \frac{32(N-1)(2+N)(-1+3N)}{N^2(1+N)^2}S_3 + \frac{96(N-1)}{N(1+N)}S_3 \\ &- \frac{32P_{2,2}}{N$$

$$\begin{split} &+\frac{192(N-1)}{N(1+N)}S_{2,1,1}-\frac{256(N-1)}{N(1+N)}S_{-2,1,1}-\frac{192(N-1)\left(-5+3N+3N^2\right)}{N^2(1+N)^2}\zeta_3 \\ &+C_4T_F^2N_F^2 \Bigg[\frac{16P_{55}}{27N^4(1+N)^4} + \left(\frac{64\left(23+50N+10N^2+19N^3\right)}{27N(1+N)^3}-\frac{32(N-1)}{3N(1+N)}\right) \\ &\times S_2\Bigg)S_1 - \frac{64\left(-2+5N^2\right)}{9N(1+N)^2}S_1^2 + \frac{32(N-1)}{9N(1+N)}S_1^3 - \frac{64\left(-2+6N+5N^2\right)}{9N(1+N)^2}S_2 \\ &+\frac{64(N-1)}{9N(1+N)}S_3 - \frac{128(-2+5N)}{9N(1+N)}S_{-2} + \frac{128(N-1)}{3N(1+N)} \Big[S_{-3} + S_{2,1}\Big]\Bigg] \\ &+C_A^2N_FT_F\Bigg[\frac{16P_{34}}{9N^3(1+N)^3}S_2 - \frac{8P_{70}}{27N^5(1+N)^5(2+N)} + \left(-\frac{8P_{56}}{27N^4(1+N)^4}\right) \\ &+\frac{8\left(-72+181N-48N^2+11N^3\right)}{3N^2(1+N)^2}S_2 - \frac{704(N-1)}{3N(1+N)}S_3 + \frac{128(N-1)}{N(1+N)}S_{2,1} \\ &+\frac{512(N-1)}{N(1+N)}S_{-2,1} + \frac{192(N-1)}{N(1+N)}\zeta_3\Bigg)S_1 + \left(\frac{16P_{31}}{9N^3(1+N)^3} - \frac{160(N-1)}{N(1+N)}S_2\Bigg)S_1^2 \\ &-\frac{8\left(-24-59N+11N^3\right)}{9N^2(1+N)^2}S_3^3 - \frac{32(N-1)}{N(1+N)}S_1^4 - \frac{16(N-1)}{N(1+N)}S_2^2 \\ &-\frac{16(345-428N+11N^3)}{9N^2(1+N)^2}S_3 - \frac{32(N-1)}{N(1+N)}S_1 + \left(\frac{32P_{49}}{9N^3(1+N)^3(2+N)}\right) \\ &-\frac{64(-5+N)(-1+2N)}{N^2(1+N)^2}S_3 - \frac{32(N-1)}{N(1+N)}S_1^2 - \frac{128(N-1)}{N(1+N)}S_1\Bigg)S_{-3} \\ &-\frac{352(N-1)}{N(1+N)}S_{-2} + \left(-\frac{32(69-92N+11N^3)}{3N^2(1+N)^2} - \frac{512(N-1)}{N(1+N)}S_1\Bigg)S_{-3} \\ &-\frac{352(N-1)}{N(1+N)}S_{-2,1} + \frac{448(-1+N)}{N(1+N)}S_{-2,2} + \frac{512(-1+N)}{N(1+N)}S_{-3,1} \\ &-\frac{66(-7+11N)}{N(1+N)}S_{-2,1} + \frac{448(-1+N)}{N(1+N)}S_{-2,2} + \frac{512(-1+N)}{N(1+N)}S_{-3,1} \\ &-\frac{768(-1+N)}{N(1+N)}S_{-2,1} + \frac{96(-1+N)(-8+3N+3N^2)}{N^2(1+N)^2}\zeta_3 \Bigg] \\ &+C_F^2T_FN_F\Bigg[-\frac{8P_{26}}{N^3(1+N)^3}S_1^2 + \frac{8P_{28}}{N^3(1+N)^3}S_2 - \frac{704(N-1)}{3N(1+N)}S_3 \\ &+\frac{256(N-1)}{N(1+N)}S_{2,1}\Bigg)S_1 - \frac{8(N-1)(-10-9N+3N^2)}{N^2(1+N)^2}S_3 - \frac{160(N-1)}{3N^2(1+N)^2}S_3 \\ &-\frac{160(N-1)}{3N^2(1+N)^2}S_3 - \frac{160(N-1)}{3N^2(1+N)^2}S_3 \\ &+\frac{256(N-1)}{N(1+N)}S_2^2 - \frac{16(N-1)(-22+27N+3N^2)}{3N^2(1+N)^2}S_3 - \frac{160(N-1)}{N(1+N)}S_4 \\ \end{aligned}$$

$$\begin{split} &+ \left(\frac{64P_{21}}{N^2(1+N)^3(2+N)} - \frac{256(N-1)}{N(1+N)^2}S_1 - \frac{128(N-1)}{N(1+N)}S_4\right)S_{-2} - \frac{64(N-1)}{N(1+N)}S_{-2}^2 \\ &+ \left(-\frac{128(N-1)^2}{N^2(1+N)^2} - \frac{256(N-1)}{N(1+N)}S_1\right)S_{-3} - \frac{320(N-1)}{N(1+N)}S_{-4} - \frac{128(N-1)}{N^2(1+N)^2}S_{2,1} \\ &+ \frac{64(N-1)}{N(1+N)}S_{3,1} + \frac{256(N-1)}{N(1+N)^2}S_{-2,1} + \frac{128(N-1)}{N(1+N)}S_{-2,2} + \frac{256(N-1)}{N(1+N)}S_{-3,1} \\ &- \frac{192(N-1)}{N(1+N)}S_{2,1,3} + \frac{96(N-1)(-2+3N+3N^2)}{N^2(1+N)^2}\zeta_3\right], \quad (56) \\ &\Delta\gamma_{gq}^{(2)} = C_F^2 \left[C_A \left[\frac{4P_{39}}{9N^3(1+N)^3}S_2 + \frac{P_{73}}{54(N-1)N^4(1+N)^5} + \left(-\frac{4P_{38}}{27N^4(1+N)^4} - \frac{8(30+203N+177N^2+49N^3)}{3N^2(1+N)^2}S_2 - \frac{640(2+N)}{3N(1+N)}S_3 + \frac{64(2+N)}{N(1+N)}S_{2,1} \right] \\ &+ \frac{128(2+N)}{N(1+N)}S_{-2,1} + \frac{288(2+N)}{88(2+N)}\zeta_3\right)S_1 + \left(\frac{4P_{36}}{9N^3(1+N)^3} - \frac{16(2+N)}{N(1+N)}S_2\right)S_1^2 \\ &- \frac{8(6+85N+132N^2+50N^3)}{9N^2(1+N)^2}S_1^3 + \frac{8(2+N)(102+65N+29N^2)}{9N^2(1+N)^2}S_3 \\ &+ \frac{16(2+N)}{3N(1+N)}S_1^4 - \frac{32(2+N)}{N(1+N)}S_4 + \left(-\frac{16P_{44}}{(N-1)N^3(1+N)^3} - \frac{64(7+3N)}{N(1+N)}S_1\right) \\ &- \frac{96(2+N)}{N(1+N)}S_1^2 - \frac{2(2+N)(4+3N)}{N(1+N)}S_4 + \frac{64(7+3N)}{N(1+N)}S_1 \\ &- \frac{96(2+N)}{N(1+N)}S_1^2 - \frac{32(2+N)}{N(1+N)}S_4 + \left(-\frac{16P_{44}}{3N^2(1+N)^2}S_{2,1} + \frac{224(2+N)}{N(1+N)}S_{3,1} \right) \\ &+ \frac{80(2+N)}{N(1+N)}S_1^2 - \frac{32(2+N)(1+N+N}{N(1+N)}S_2 - \frac{96(2+N)}{N(1+N)}S_{2,1} + \frac{224(2+N)}{N(1+N)}S_{3,1} \\ &- \frac{32(2+N)(5+3N)}{N(1+N)}S_{-2,1} - \frac{432(2+N)}{N(1+N)}S_2 - \frac{96(2+N)}{N(1+N)}S_{2,1} + \frac{224(2+N)}{N(1+N)}S_3 \\ &+ \frac{60(2+N)}{N(1+N)}S_1^3 - \frac{16(2+N)(9+46N+67N^2)}{9N^2(1+N)^2}S_2 + \frac{256(2+N)}{9N(1+N)}S_3 \\ &+ \frac{60(2+N)}{N(1+N)}S_1^3 - \frac{16(2+N)(9+46N+67N^2)}{9N^2(1+N)^2}S_2 + \frac{256(2+N)}{9N(1+N)}S_3 \\ &+ \frac{256}{(N-1)N^2(1+N)^2}S_2 - \frac{64(2+N)}{3N(1+N)}S_{2,1} - \frac{128(2+N)}{9N(1+N)}S_3 \\ &+ \frac{256}{(N-1)N^2(1+N)^2}S_2 - \frac{64(2+N)}{3N(1+N)}S_2 - \frac{162(2+N)}{9N(1+N)}S_3 \\ &+ \frac{256}{(N-1)N^2(1+N)^2}S_2 - \frac{64(2+N)}{3N(1+N)}S_2 - \frac{162(2+N)}{9N(1+N)^2}S_1 \\ &+ \frac{256}{(N-1)N^2(1+N)^2}S_1 - \frac{64(2+N)(2+5N)}{9N(1+N)^2}S_1 \\ &+ \frac{256}{(N-1)N^2(1+N)^2}S_1 + \frac{64(2+N)(2+5N)}{9N(1+N)^2}S_1 \\ &+ \frac{256}{(N-1$$

$$\begin{split} + & \left( -\frac{16P_{37}}{27N^3(1+N)^3} + \frac{80(2+N)S_2}{3N(1+N)} \right) S_1 + \frac{16(18+116N+129N^2+43N^3)}{9N^2(1+N)^2} S_1^2 \\ - & \frac{80(2+N)}{9N(1+N)} S_1^3 + \frac{16(-2+16N+9N^2+N^3)}{3N^2(1+N)^2} S_2 + \frac{512(2+N)}{9N(1+N)} S_3 \\ + & \left( -\frac{64P_7}{3(-1+N)N^2(1+N)^2} + \frac{256(2+N)S_1}{3N(1+N)} \right) S_2 + \frac{128(2+N)}{9N(1+N)} S_3 \\ + & \left( -\frac{64P_7}{3(-1+N)N^2(1+N)^2} + \frac{256(2+N)S_1}{3N^3(1+N)^3} \right) S_2 - \frac{128(2+N)}{3N(1+N)} S_3 \\ + & \left( \frac{4P_{62}}{27(N-1)N^4(1+N)^4} - \frac{4(120+158N+141N^2+55N^3)}{3N^2(1+N)^2} S_2 + \frac{128(2+N)}{3N(1+N)} S_3 \\ + & \left( \frac{4P_{62}}{27(N-1)N^4(1+N)^4} - \frac{4(120+158N+141N^2+55N^3)}{3N^2(1+N)^2} S_2 + \frac{128(2+N)}{3N(1+N)} S_3 \\ + & \frac{128(2+N)}{N(1+N)} S_{-2,1} - \frac{96(2+N)}{N(1+N)} \zeta_3 \right) S_1 + \left( -\frac{2P_{38}}{9N^3(1+N)^3} + \frac{48(2+N)}{N(1+N)} S_2 \right) S_1^2 \\ + & \frac{4(24+158N+165N^2+55N^3)}{9N^2(1+N)^2} S_1^3 - \frac{8(2+N)}{3N(1+N)} S_1^4 - \frac{40(2+N)}{N(1+N)} S_2^2 \\ - & \frac{8(-186+295N+528N^2+176N^3)}{9N^2(1+N)^2} S_3 - \frac{16(2+N)}{N(1+N)} S_4 \\ + & \left( -\frac{32P_{12}}{3(N-1)N^2(1+N)^2} S_1 + \frac{16P_{47}}{3(N-1)N^3(1+N)^3} + \frac{96(2+N)}{N(1+N)} S_1^2 \\ - & \frac{64(2+N)}{N(1+N)} S_2 \right) S_{-2} - \frac{16(2+N)}{N(1+N)} S_{-2} + \left( -\frac{16(-126-13N+66N^2+22N^3)}{3N^2(1+N)^2} \right) \\ - & \frac{64(2+N)}{N(1+N)} S_3 + \frac{224(2+N)}{N(1+N)} S_{-2,2} + \frac{256(2+N)}{N(1+N)} S_{-3,1} - \frac{384(2+N)}{N(1+N)} S_{-2,1,1} \\ + & \frac{144(2+N)}{N(1+N)} S_3 \right] + C_F^2 \left[ -\frac{2(2+N)P_{19}}{N^3(1+N)^3} S_2 + \frac{266(2+N)}{N(1+N)} S_{-3,1} - \frac{384(2+N)}{N(1+N)} S_{-2,1,1} \\ + & \frac{144(2+N)}{N(1+N)} S_3 \right] + C_F^2 \left[ -\frac{2(2+N)P_{19}}{N^3(1+N)^3} S_2 + \frac{266(2+N)}{N(1+N)} S_{-3,1} - \frac{384(2+N)}{N(1+N)} S_2 \right) S_1^2 \\ + & \left( -\frac{4(2+N)P_{43}}{N(1+N)} + \frac{4(2+N)(-2+19N+39N^2)}{N^3(1+N)^3} S_2 + \frac{128(2+N)}{N(1+N)} S_3 \right) S_1^2 \\ + & \left( -\frac{4(2+N)P_{43}}{N(1+N)} S_1 - \frac{38(2+N)}{N(1+N)} S_1 + \frac{24(2+N)}{N(1+N)} S_2 \right) S_1^2 \\ + & \frac{4(2+N)(-2+3N+15N^2)}{N^3(1+N)^3} S_1 - \frac{8(2+N)}{N(1+N)} S_1^4 - \frac{24(2+N)}{N(1+N)} S_2^2 \\ + & \frac{4(2+N)(-2+3N+15N^2)}{3N^2(1+N)^2} S_1 - \frac{8(2(2+N)}{N(1+N)} S_1 + \frac{28(2+N)}{N(1+N)} S_2 \right) S_1 \\ + & \frac{28(2+N)}{N(1+N)^2} S_1 - \frac{64(2+N)}{N(1$$

$$\begin{split} &-\frac{96(2+N)}{N(1+N)}S_{-2}^{2} + \left(-\frac{64(N-1)(2+N)}{N^{2}(1+N)^{2}} - \frac{128(2+N)}{N(1+N)}S_{1}\right)S_{-3} \\ &-\frac{160(2+N)}{N(1+N)}[S_{-4} + S_{3,1}] + \frac{128(2+N)}{N(1+N)^{2}}[S_{-2,1} + (1+N)S_{-3,1}] \\ &+\frac{64(2+N)}{N(1+N)}S_{-2,2} + \frac{96(2+N)}{N(1+N)}S_{1,1} + \frac{288(2+N)(\varsigma_{3})}{N(1+N)^{3}}\right], \quad (57) \\ &\Delta\gamma_{09}^{(2)} = C_{A}T_{F}^{2}N_{F}^{2}\left[-\frac{16F_{8}}{27N^{2}(1+N)^{2}}S_{1} - \frac{4P_{48}}{27N^{2}(1+N)^{2}}\right] + C_{F}\left[T_{F}^{2}N_{F}^{2}\left[-\frac{8F_{59}}{27N^{4}(1+N)^{4}} + \frac{64(N-1)(2+N)(-6-8N+N^{2})}{9N^{3}(1+N)^{3}}S_{1} + \frac{32(N-1)(2+N)}{3N^{2}(1+N)^{2}}S_{1}^{2} + \frac{32(N-1)(2+N)}{9N^{3}(1+N)^{2}}S_{1}^{2} + \frac{22F_{77}}{N^{2}(1+N)^{2}}S_{2}\right] + C_{A}T_{F}N_{F}\left[\frac{8P_{6}}{N^{3}(1+N)^{3}}S_{2} - \frac{8P_{9}}{3N^{3}(1+N)^{3}}S_{1}^{2} + \frac{22F_{77}}{27(N-1)N^{5}(1+N)^{5}(2+N)} + \left(-\frac{8F_{67}}{9(-1+N)N^{4}(1+N)^{4}(2+N)} - \frac{32(N-1)(2+N)}{N^{2}(1+N)^{2}}S_{2} + 128\zeta_{3}\right)S_{1} + \frac{32(N-1)(2+N)}{3N^{2}(1+N)^{2}}S_{1}^{3} - \frac{32(34+N+N^{2})}{3N^{2}(1+N)^{2}} \\ &\times S_{3} + \left(\frac{128P_{2}}{(N-1)N^{2}(1+N)^{2}}(2+N)S_{1} - \frac{32P_{33}}{3N^{2}(1+N)^{2}}S_{-2,1} - \frac{192(4-N-N^{2})}{N^{2}(1+N)^{2}}S_{-3,3} + \frac{64(N-1)(2+N)}{N^{2}(1+N)^{2}}S_{-2,1} - \frac{128(-8+N+N^{2})}{N^{2}(1+N)^{2}}S_{-2,1} \\ &-\frac{64(-3+N)(4+N)}{N^{2}(1+N)^{2}}\zeta_{3}\right]\right] + C_{4}^{3}\left[\frac{64P_{16}}{9N^{2}(1+N)^{2}}S_{-2,1} - \frac{32P_{18}}{N^{2}(1+N)^{2}}S_{-2,1} \\ &-\frac{64(-3+N)(4+N)}{N^{2}(1+N)^{2}}\zeta_{3}\right]\right] + C_{4}^{3}\left[\frac{64P_{16}}{9N^{2}(1+N)^{2}}S_{-2,1} - \frac{32P_{18}}{N^{2}(1+N)^{2}}S_{-2,1} \\ &-\frac{64P_{17}}{N^{2}(1+N)^{2}}S_{2} + 128S_{2}^{2} + \frac{16(-96+11N+11N^{2})}{3N(1+N)}S_{3} + 192S_{4} \\ &+\frac{1024}{N(1+N)^{3}}S_{-2} + 128S_{2}^{2} - 768S_{-3,1} + 1024S_{-2,1,1}\right)S_{1} \\ &+\left(-\frac{256(1+3N+3N^{2})}{N^{3}(1+N)^{3}} + 128S_{3} - 256S_{-2,1}\right)S_{1}^{2} + \left(-\frac{16P_{41}}{9N^{3}(1+N)^{3}} \\ &+64S_{3} + 640S_{-2,1}\right)S_{2} - \frac{256}{N(1+N)}S_{2}^{2} - \frac{384}{N(1+N)}S_{4} + 64S_{5} \\ &+\left(\frac{32P_{32}}{9(N-1)N^{3}(1+N)^{3}(2+N)} + \left(-\frac{64P_{32}}{9(N-1)N^{3}(1+N)^{3}} \\ &+64S_{3} - \frac{512}{N(1+N)^{3}}S_{2} + 128S_{3} - 768S_{2,1}\right)S_{-2} + \left(-\frac{16(24+11N+11N^{2})}{3N(1+N)}\right) \\ \end{aligned}$$

$$\begin{split} + 64S_{1} \bigg) S_{-2}^{2} + \bigg( -\frac{32P_{15}}{9N^{2}(1+N)^{2}} - \frac{1536}{N(1+N)}S_{1} + 384S_{1}^{2} - 320S_{2} \bigg) S_{-3} \\ + \bigg( -\frac{1024}{N(1+N)} + 512S_{1} \bigg) S_{-4} - 192S_{-5} - 384S_{2,-3} + \frac{1280}{N(1+N)}S_{-2,2} \\ + 384S_{-2,3} + \frac{1536}{N(1+N)}S_{-3,1} - 384S_{-4,1} + 768S_{2,1,-2} - \frac{2048}{N(1+N)}S_{-2,1,1} \\ + 768[S_{-2,2,1} + S_{-3,1,1}] - 1536S_{-2,1,1,1} \bigg] \\ + C_{F}^{2}T_{F}N_{F} \bigg[ -\frac{4P_{75}}{(N-1)N^{5}(1+N)^{5}(2+N)} + \bigg( \frac{32(N-1)(2+N)S_{2}}{N^{2}(1+N)^{2}} \\ - \frac{16P_{42}}{N^{4}(1+N)^{4}} \bigg) S_{1} + \frac{8(N-1)(2+N)(2+3N+3N^{2})}{N^{3}(1+N)^{3}} S_{1}^{2} - \frac{32(N-1)(2+N)}{3N^{2}(1+N)^{2}} \\ \times S_{1}^{3} - \frac{8(2+N)(2-11N-16N^{2}+9N^{3})}{N^{3}(1+N)^{3}} S_{2} + \frac{32(10+7N+7N^{2})}{3N^{2}(1+N)^{2}} S_{3} \\ + \bigg( -\frac{64(10+N+N^{2})}{N^{3}(1+N)^{2}} + \frac{512}{N^{2}(1+N)^{2}} S_{1} \bigg) S_{-2} + \frac{256}{N^{2}(1+N)^{2}} S_{-3} \\ - \frac{64(N-1)(2+N)}{N^{2}(1+N)^{2}} S_{2,1} - \frac{512}{N^{2}(1+N)^{2}} S_{-2,1} + \frac{192(-2-N-N^{2})}{N^{2}(1+N)^{2}} \zeta_{3} \bigg] \\ + C_{A}^{2}T_{F}N_{F} \bigg[ \frac{32P_{4}}{9N^{2}(1+N)^{2}} S_{2} + \frac{32P_{11}}{9N^{2}(1+N)^{2}} S_{-3} - \frac{64P_{11}}{9N^{2}(1+N)^{2}} S_{-2,1} \\ + \frac{16P_{13}}{9N^{2}(1+N)^{2}} S_{3} + \frac{2P_{76}}{27(N-1)N^{5}(1+N)^{5}(2+N)} + \bigg( \frac{1280}{9} S_{2} - \frac{64}{3} S_{3} \\ - \frac{8P_{68}}{27(-1+N)N^{4}(1+N)^{4}(2+N)} - 128\zeta_{3} \bigg) S_{1} + \frac{64}{3} S_{-2}^{2} \\ + \bigg( \frac{64P_{45}}{9(N-1)N^{2}(1+N)^{2}} \zeta_{3} \bigg], \qquad (58)$$

with the polynomials

$$P_{1} = N^{4} - 2N^{3} - 4N^{2} + 15N + 2,$$

$$P_{2} = N^{4} + 2N^{3} - 5N^{2} - 6N + 16$$
(59)
(60)

$$P_{2} = N^{4} + 2N^{3} - 5N^{2} - 6N + 16,$$

$$P_{3} = N^{4} + 44N^{3} + 45N^{2} + 38N + 12,$$
(60)
(61)

- $P_4 = 3N^4 + 6N^3 89N^2 92N + 12, (62)$
- $P_5 = 3N^4 + 6N^3 + 16N^2 + 13N 3, (63)$
- $P_6 = 3N^4 + 18N^3 + 17N^2 46N 28, \tag{64}$
- $P_7 = 5N^4 + 9N^3 4N^2 4N + 6, (65)$
- $P_8 = 8N^4 + 16N^3 19N^2 27N + 48, (66)$

$$\begin{array}{l} P_{1} = 11\Lambda^{4} + 22\Lambda^{3} + 13\Lambda^{2} + 2\Lambda - 12, \\ P_{10} = 11\Lambda^{4} + 34\Lambda^{3} + \Lambda^{2} - 70\Lambda - 12, \\ P_{11} = 20\Lambda^{4} + 40\Lambda^{3} + 11\Lambda^{2} - 9\Lambda + 54, \\ P_{12} = 22\Lambda^{4} + 50\Lambda^{3} + 5\Lambda^{2} - 47\Lambda + 6, \\ P_{12} = 22\Lambda^{4} + 50\Lambda^{3} + 5\Lambda^{2} - 47\Lambda + 6, \\ P_{13} = 40\Lambda^{4} + 80\Lambda^{3} + 73\Lambda^{2} + 33\Lambda + 54, \\ P_{15} = 67\Lambda^{4} + 13\Lambda^{3} + 49\Lambda^{2} + 54\Lambda - 360, \\ P_{16} = 67\Lambda^{4} + 13\Lambda^{3} + 49\Lambda^{2} + 54\Lambda - 72, \\ P_{16} = 67\Lambda^{4} + 13\Lambda^{3} + 67\Lambda^{2} + 14\Lambda + 72, \\ P_{17} = 67\Lambda^{4} + 13\Lambda^{3} + 67\Lambda^{2} + 14\Lambda + 72, \\ P_{17} = 67\Lambda^{4} + 13\Lambda^{3} + 67\Lambda^{2} + 14\Lambda + 72, \\ P_{17} = 67\Lambda^{4} + 13\Lambda^{3} + 67\Lambda^{2} + 14\Lambda + 72, \\ P_{17} = 67\Lambda^{4} + 13\Lambda^{3} + 3\Lambda^{2} - 38\Lambda - 12, \\ P_{20} = N^{5} + N^{4} - 4\Lambda^{3} + 3\Lambda^{2} - 38\Lambda - 12, \\ P_{21} = 2\Lambda^{5} + 6\Lambda^{4} - N^{3} - 8\Lambda^{2} + 15\Lambda + 10, \\ P_{22} = 2\Lambda^{5} + 6\Lambda^{4} + 5\Lambda^{3} + 4\Lambda^{2} + 9\Lambda - 2, \\ P_{21} = 2\Lambda^{5} + 6\Lambda^{4} + 5\Lambda^{3} + 4\Lambda^{2} + 9\Lambda - 2, \\ P_{22} = 2\Lambda^{5} + 6\Lambda^{4} + 5\Lambda^{3} + 4\Lambda^{2} + 9\Lambda - 2, \\ P_{23} = 3\Lambda^{5} + 5\Lambda^{4} - 33\Lambda^{3} - 45\Lambda^{2} + 6\Lambda - 16, \\ P_{24} = 3\Lambda^{5} + 1\Lambda^{4} + 21\Lambda^{3} + 20\Lambda^{2} - 22\Lambda - 10, \\ P_{25} = 8\Lambda^{5} - 31\Lambda^{4} + 20\Lambda^{5} - 35\Lambda^{2} - 25\Lambda - 20, \\ P_{26} = 14\Lambda^{5} + 15\Lambda^{4} - 19\Lambda^{3} - 13\Lambda^{2} - 25\Lambda - 20, \\ P_{28} = 26\Lambda^{5} + 41\Lambda^{4} - 21\Lambda^{3} + 21\Lambda^{2} - 21\Lambda - 108, \\ P_{29} = 36\Lambda^{5} - 55\Lambda^{4} - 23\Lambda^{3} + 16\Lambda^{2} - 90\Lambda - 72, \\ P_{30} = 58\Lambda^{5} + 7\Lambda^{4} + 59\Lambda^{3} + 60\Lambda^{2} + 3\Lambda - 108, \\ P_{31} = 67\Lambda^{5} + 201\Lambda^{4} - 67\Lambda^{3} - 57\Lambda^{2} - 103\Lambda - 108, \\ P_{31} = 67\Lambda^{5} + 201\Lambda^{4} - 67\Lambda^{3} - 57\Lambda^{2} - 103\Lambda - 108, \\ P_{31} = 67\Lambda^{5} + 201\Lambda^{4} + 67\Lambda^{3} - 57\Lambda^{2} + 109\Lambda - 189, \\ P_{31} = 65\Lambda^{5} + 152\Lambda^{4} + 37\Lambda^{3} + 3398\Lambda^{2} + 284\Lambda + 864, \\ (96) \\ P_{31} = 35\Lambda^{5} + 152\Lambda^{4} + 176\Lambda^{3} + 650\Lambda^{2} + 44\Lambda + 144, \\ (95) \\ P_{32} = 631\Lambda^{5} + 55\Lambda^{4} + 176\Lambda^{3} + 650\Lambda^{2} + 44\Lambda + 144, \\ (95) \\ P_{34} = 3\Lambda^{6} + 9\Lambda^{5} + 5\Lambda^{4} + 3R\Lambda^{3} + 2398\Lambda^{2} + 284\Lambda + 864, \\ (96) \\ P_{39} = 725\Lambda^{5} + 231\Lambda^{4} + 118\Lambda^{3} - 169N^{2} + 284\Lambda + 864, \\ (96) \\ P_{41} = 3\Lambda^{6} + 3\Lambda^{5} + 5\Lambda^{4} + 58\Lambda^{3} - 52\Lambda^{2} + 104\Lambda + 14\lambda, \\ (96) \\ P_{42} = 6\Lambda^{6} + 34\Lambda^{5} + 11\Lambda^{4} - 10\Lambda^{3} +$$

$$\begin{split} &P_{49} = 94\Lambda^6 + 315\Lambda^5 + 145\Lambda^4 - 63\Lambda^3 + 148\Lambda^2 - 441\Lambda^3 - 18, &(107) \\ P_{50} = 95\Lambda^6 + 285\Lambda^5 + 92\Lambda^4 - 291\Lambda^3 - 97\Lambda^2 + 96\Lambda - 36, &(108) \\ P_{51} = 160\Lambda^6 + 438\Lambda^5 + 364\Lambda^4 + 330\Lambda^3 + 529\Lambda^2 + 321\Lambda + 18, &(109) \\ P_{52} = 325\Lambda^6 + 975\Lambda^5 + 85\Lambda^4 - 879\Lambda^3 + 598\Lambda^2 - 240\Lambda - 72, &(110) \\ P_{53} = 17\Lambda^7 + 9\Lambda^6 - 95\Lambda^5 - 19\Lambda^4 + 76\Lambda^3 + 22\Lambda^2 + 26\Lambda + 28, &(111) \\ P_{54} = 21\Lambda^7 + 33\Lambda^6 + 13\Lambda^5 - 28\Lambda^4 - 31\Lambda^3 - 33\Lambda^2 - 26\Lambda - 8, &(112) \\ P_{55} = 165\Lambda^7 + 330\Lambda^6 + 191\Lambda^5 - 365\Lambda^4 - 136\Lambda^3 - 445\Lambda^2 - 18\Lambda + 144, &(113) \\ P_{56} = 475\Lambda^7 + 833\Lambda^6 + 1527\Lambda^5 + 2905\Lambda^4 - 1342\Lambda^3 + 5650\Lambda^2 + 3834\Lambda + 486, &(114) \\ P_{57} = 537\Lambda^7 + 1200\Lambda^6 - 1013\Lambda^5 - 2085\Lambda^4 + 1720\Lambda^3 - 855\Lambda^2 - 2468\Lambda - 492, &(115) \\ P_{58} = 1199\Lambda^7 + 3323\Lambda^6 + 161\Lambda^5 - 5053\Lambda^4 - 4214\Lambda^3 - 4800\Lambda^2 - 3168\Lambda - 864, &(116) \\ P_{59} = 33\Lambda^8 + 132\Lambda^7 + 70\Lambda^6 - 612\Lambda^5 - 839\Lambda^4 + 480\Lambda^3 + 712\Lambda^2 + 408\Lambda + 144, &(117) \\ P_{60} = 476\Lambda^8 + 2297\Lambda^7 + 2018\Lambda^6 - 4915\Lambda^5 - 7324\Lambda^4 + 242\Lambda^3 - 1218\Lambda^2 & -2700\Lambda - 864, &(118) \\ P_{51} = 914\Lambda^8 + 3005\Lambda^7 + 3368\Lambda^6 + 4349\Lambda^5 + 5183\Lambda^4 + 548\Lambda^3 + 1101\Lambda^2 & +936\Lambda + 324, &(129) \\ P_{62} = 2078\Lambda^8 + 8225\Lambda^7 + 10475\Lambda^6 - 1921\Lambda^5 - 10729\Lambda^4 - 2560\Lambda^3 - 5658\Lambda^2 - 7578\Lambda & -2700, &(120) \\ P_{63} = -5\Lambda^9 - 25\Lambda^8 + 228\Lambda^7 + 926\Lambda^6 - 201\Lambda^5 - 2377\Lambda^4 + 626\Lambda^3 + 2788\Lambda^2 + 2168\Lambda & +480, &(121) \\ P_{64} = 99\Lambda^9 + 297\Lambda^8 - 982\Lambda^7 - 662\Lambda^6 + 1035\Lambda^5 - 3079\Lambda^4 + 3448\Lambda^3 + 2868\Lambda^2 & (122) \\ P_{55} = -1251\Lambda^{10} - 655\Lambda^9 - 10972\Lambda^8 - 17422\Lambda^7 - 1423\Lambda^6 + 111905\Lambda^5 + 149894\Lambda^4 & \\ -2116\Lambda^3 - 37752\Lambda^2 + 1238\Lambda + 6012, &(123) \\ P_{56} = -115\Lambda^{10} + 825\Lambda^8 + 1102\Lambda^8 - 578\Lambda^7 - 1939\Lambda^6 - 239\Lambda^6 + 1184\Lambda^4 + 448\Lambda^3 & \\ -2456\Lambda^2 - 2256\Lambda - 864, &(125) \\ P_{69} = 735\Lambda^{10} + 3055\Lambda^9 + 6060\Lambda^8 + 6934\Lambda^7 + 11743\Lambda^6 - 41\Lambda^5 - 18290\Lambda^4 - 920\Lambda^3 & \\ + 2898\Lambda^2 - 11065\Lambda - 3744, &(127) \\ P_{70} = 741\Lambda^{10} + 3705\Lambda^9 + 2650\Lambda^8 + 1058\Lambda^7 - 1375\Lambda^6 - 11707\Lambda^5 + 446\Lambda^4 + 17123\Lambda^3 & \\ + 3432\Lambda^2 - 6624\Lambda - 3456, &(128) \\ P_{71} = 1657\Lambda^{10} + 8557\Lambda^9 + 5010\Lambda^8 + 10583\Lambda^7 - 3275\Lambda^6 - 11707\Lambda^5 + 446\Lambda^4 + 11723\Lambda^3 & \\ + 3432\Lambda^2 - 6624\Lambda - 3456, &(129) \\ P_{71} = 18$$

$$+31109N^{6} - 29734N^{5} + 318684N^{4} - 18512N^{3} + 82224N^{2} + 126720N + 48384,$$
(132)  

$$P_{75} = N^{12} + 6N^{11} - 27N^{10} - 186N^{9} - 197N^{8} + 310N^{7} + 899N^{6} + 1198N^{5} + 1020N^{4} + 112N^{3} - 192N^{2} + 64N + 64,$$
(133)  

$$P_{76} = 699N^{12} + 4194N^{11} + 16447N^{10} + 43214N^{9} + 42657N^{8} - 19098N^{7} - 36963N^{6} - 11670N^{5} - 45064N^{4} - 39392N^{3} + 7536N^{2} + 8064N + 1728,$$
(134)  

$$P_{77} = 723N^{12} + 4338N^{11} + 12623N^{10} + 17230N^{9} - 8583N^{8} - 30018N^{7} + 47709N^{6}$$

$$+75738N^{5} + 8776N^{4} + 67208N^{3} + 4416N^{2} - 41184N - 20736.$$
(135)

We mention the moment-relation

$$\Delta \gamma_{gg}^{(2)}(N=1) = -2\beta_2, \tag{136}$$

with [43, 44]

$$\beta_2 = \frac{2857}{54} C_A^3 + T_F N_F \left[ -\frac{1415}{27} C_A^2 - \frac{205}{9} C_A C_F + 2C_F^2 \right] + (T_F N_F)^2 \left[ \frac{158}{27} C_A + \frac{44}{9} C_F \right].$$
(137)

The so-called supersymmetric relation obtained by setting  $C_F = C_A = 2T_F N_F$  is broken using dimensional regularization from  $O(a_s^2)$  onward, while the relation

$$\Delta\gamma_{qq}^{(k)} + \Delta\gamma_{gq}^{(k)} - \Delta\gamma_{qg}^{(k)} - \Delta\gamma_{gg}^{(k)} = 0$$
(138)

also holds at k = 1 using dimensional reduction [17] in this limit.

Furthermore, one has

$$\Delta \gamma_{aa}^{(k)}(N=1) = 0, \text{ for } k = 0, 1, 2.$$
(139)

The expansions have been performed using the HarmonicSums command HarmonicSumsSeries since (56,58) contain evanescent poles at N = 1. More generally, also the first moment of the gluonic Wilson coefficient, related to (139), for the structure function  $g_1(x, Q^2)$  both for the massless [80, 81] and the massive case in the asymptotic representation to two-loop order [82] vanishes. This is known at one-loop order even for general kinematics [83].

The splitting functions in z-space are obtained by a Mellin inversion, cf. (36), and are given in computer readable form in an attachment to the paper.

#### 5 Comparison to the literature

We confirm the results for the singlet anomalous dimensions calculated in [20], where the on-shell forward Compton amplitude has been used for the computation. The contributions  $\propto T_F$  have already been calculated independently as a by-product of the massive on-shell operator matrix elements in Ref. [21], to which we also agree. The comparison to the large  $N_F$  expansions of Refs. [84,85] has been given in our previous paper [21] already, where all these terms are covered.

Let us finally consider the small  $z \text{ limit}^5$  of the present results and compare to the predictions given in [87, 88]. In Mellin–N space these terms are given by the most singular contribution expanding around N = 0. One obtains at one– and two–loop order

$$\Delta \gamma^{(0),N\to0} = -\frac{4}{N} \begin{pmatrix} C_F & -2T_F N_F \\ 2C_F & 4C_A \end{pmatrix}$$
(140)

<sup>&</sup>lt;sup>5</sup>For a review on the small z predictions of the different evolution kernels see [86].

$$\Delta \gamma^{(1),N \to 0} = -\frac{8}{N^3} \begin{pmatrix} C_F(2C_A - 3C_F - 4T_F N_F) & 2C_F(2C_A + C_F) \\ -2C_F(2C_A + C_F) & 4(2C_A^2 - C_F T_F N_F) \end{pmatrix}$$
(141)

which agree with the evolution kernels given in Refs. [87,88].

At three-loop order we have in the M–scheme

$$\Delta \gamma^{(2),N \to 0} = -\frac{16}{N^5} \times \begin{pmatrix} C_F (3C_A^2 - 12C_A C_F + 10C_F^2 + T_F N_F (12C_A + 16C_F)) & 2T_F N_F (15C_A^2 + 4C_A C_F - 8C_F T_F N_F) \\ -2C_F (15C_A^2 + 8C_A C_F - 4C_F^2 - 8C_F T_F N_F) & -4(14C_A^3 + T_F N_F (C_A^2 - 12C_A C_F - 2C_F^2)) \end{pmatrix}$$
(142)

and find a deviation both in case of  $\Delta \gamma_{qg}^{(2)}$  and  $\Delta \gamma_{gq}^{(2)}$ , already noticed in [89]. The relative deviation amounts to ~ ±3.2%, with the difference of the expression in the M–scheme and the result obtained for the infrared evolution equation (IEE),  $\delta \Delta \gamma_{ij}^{(2),N\to0} = \Delta \gamma_{ij}^{(2),N\to0,M} - \Delta \gamma_{ij}^{(2),N\to0,IEE}$ ,

$$\delta \Delta \gamma_{qg}^{(2),N \to 0} = \frac{48}{N^5} C_F T_F N_F (C_A - C_F), \qquad (143)$$

$$\delta \Delta \gamma_{gq}^{(2),N \to 0} = \frac{64}{N^5} C_F^2 (C_A - C_F).$$
(144)

One may consider a different theory but QCD by setting  $C_A = C_F$ . In this case the prediction [87,88] for the evolution kernels agrees with the perturbative calculation of the anomalous dimensions.

It has been the group of J. Kodaira [90], who also considered the effective Wilson coefficient in the non–singlet case Ref. [91]<sup>6</sup>, finding that these contributions are suppressed by a further power in N. This also applies to the effective Wilson coefficients in the non–singlet case and therefore the evolution kernels of [91, 92] agree<sup>7</sup>. By considering the expansion of the function to be interpreted as a matrix formulation for the effective Wilson coefficient in [87],

$$\frac{N}{\mathbf{1}N - \frac{1}{8\pi^2}\mathbf{F}_0} = \mathbf{1} + \sum_{k=1}^{\infty} \left(\frac{a_s}{N^2}\right)^k \mathbf{F}_{0,k}.$$
(145)

The expansion coefficients  $\mathbf{F}_{0,k}$  are 2 × 2 matrices only depending on color factors. Therefore, in the representation of [87], the Wilson coefficients are suppressed by one power in N, like in the non-singlet case, cf. [90,91]. To perform a full comparison one has to form two scheme invariant quantities. This is not really possible in pure polarized QCD, since there is only one structure function  $g_1(x, Q^2)$  and also considering the Wilson coefficients [80,81].<sup>8</sup> In [20] scheme invariant polarized evolution kernels for the contributions to the structure function  $g_1$  and additional fictitious gravitational contributions in the gluonic channels have been formed for which the prediction in [87,88] hold.

In data analyses or the phenomenological description of the polarized structure functions the consideration of only the leading small z terms in the evolution kernels is numerically not sufficient. Subleading terms dominate over the leading order effects, cf. [86,88,93].

<sup>&</sup>lt;sup>6</sup>In using infrared evolution equations the authors of Refs. [87,91,92] do not specify the factorization scheme, which complicates comparisons with calculations performed e.g. in the M–scheme.

<sup>&</sup>lt;sup>7</sup>After correcting Ref. [92] in [93].

<sup>&</sup>lt;sup>8</sup>At the level of twist-2 the structure function  $g_2(x, Q^2)$  is not an independent quantity, but related by the Wandzura–Wilczek relation [94] to the structure function  $g_1(x, Q^2)$ . One might consider the physical pair  $\{g_1(x, Q^2), \partial g_1(x, Q^2)/\partial \ln(Q^2)\}$ , cf. [95]. However, the so–called leading powers in N are here not of the same order.

### 6 Conclusions

We have calculated the polarized three–loop singlet anomalous dimensions  $\Delta \gamma_{qq}^{(2),\text{PS}}$ ,  $\Delta \gamma_{qg}^{(2)}$ ,  $\Delta \gamma_{qg}^{(2)}$ ,  $\Delta \gamma_{qg}^{(2)}$  $\Delta \gamma_{gg}^{(2)}$ , and the non-singlet anomalous dimension  $\Delta \gamma_{qg}^{(2),s,NS}$  in Quantum Chromodynamics and agree with the results of Refs. [20,21,38]. The singlet anomalous dimensions have been calculated by using the method of massless off-shell OMEs, which has been applied for this purpose for the first time. The calculation has been fully automated referring to the Larin scheme, performing a finite transformation to the M-scheme for the final results. Both schemes are valid to describe the scaling violations of the polarized structure functions, however, with a different outcome for the polarized parton distribution functions, which are scheme-dependent quantities. Comparing to predictions of the leading small z behaviour one finds deviations for the off-diagonal elements at three–loop order, from which one concludes that the calculation in Ref. [87] is not in the M-scheme starting with three-loop order. To obtain the complete picture, one has to consider also the associated behaviour of the Wilson coefficients and to form scheme-invariant quantities. We also mention that partial checks on the polarized anomalous dimensions are possible from the pole terms of the single- and two-mass massive OMEs to three loop order, cf. [96]. Both the anomalous dimension and splitting functions are given in computer readable form in the file ANOM3pol.m attached to this paper.

Acknowledgment. We would like to thank J. Ablinger and A. Vogt for a discussion. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska–Curie grant agreement No. 764850, SAGEX and from the Austrian Science Fund (FWF) grants SFB F50 (F5009-N15) and P33530.

### A new polarized gluonic Feynman rule

The gluonic OME  $\Delta A_{gg}$  requires a new Feynman rule to calculate the anomalous dimensions in the massless case, containing a local operator with with five external gluon lines, extending the setting given in Refs. [21, 40]. The operator insertion reads in the polarized case

$$p_{1}, \nu_{1}, c_{1} \underbrace{\text{correct}}_{p_{2}, \nu_{2}, c_{2}} \underbrace{f_{2}, \nu_{3}, c_{3}}_{p_{3}, \nu_{4}, c_{4}} = -ig^{3} \frac{1 - (-1)^{N}}{2} \left( f^{c_{1}c_{2}c_{x}} f^{c_{3}c_{x}c_{y}} f^{c_{4}c_{5}c_{y}} O_{\nu_{1}\nu_{2}\nu_{3}\nu_{4}\nu_{5}}(p_{1}, p_{2}, p_{3}, p_{4}, p_{5}) \right) \\ + f^{c_{1}c_{2}c_{x}} f^{c_{3}c_{5}c_{y}} O_{\nu_{1}\nu_{2}\nu_{4}\nu_{3}\nu_{5}}(p_{1}, p_{2}, p_{4}, p_{3}, p_{5}) + f^{c_{1}c_{2}c_{x}} f^{c_{3}c_{x}c_{y}} f^{c_{3}c_{5}c_{y}} O_{\nu_{1}\nu_{2}\nu_{5}\nu_{3}\nu_{4}}(p_{1}, p_{2}, p_{5}, p_{3}, p_{4}) \\ + f^{c_{1}c_{3}c_{x}} f^{c_{2}c_{x}c_{y}} f^{c_{4}c_{5}c_{y}} O_{\nu_{1}\nu_{3}\nu_{2}\nu_{4}\nu_{5}}(p_{1}, p_{3}, p_{2}, p_{4}, p_{5}) + f^{c_{1}c_{3}c_{x}} f^{c_{4}c_{x}c_{y}} f^{c_{2}c_{5}c_{y}} O_{\nu_{1}\nu_{3}\nu_{4}\nu_{2}\nu_{5}}(p_{1}, p_{3}, p_{4}, p_{5}) \\ + f^{c_{1}c_{3}c_{x}} f^{c_{2}c_{x}c_{y}} f^{c_{4}c_{5}c_{y}} O_{\nu_{1}\nu_{3}\nu_{2}\nu_{4}\nu_{4}}(p_{1}, p_{3}, p_{5}, p_{2}, p_{4}) + f^{c_{1}c_{3}c_{x}} f^{c_{4}c_{x}c_{y}} f^{c_{2}c_{5}c_{y}} O_{\nu_{1}\nu_{4}\nu_{2}\nu_{3}\nu_{5}}(p_{1}, p_{4}, p_{3}, p_{5}, p_{2}, p_{4}) \\ + f^{c_{1}c_{3}c_{x}} f^{c_{2}c_{x}c_{y}} f^{c_{2}c_{5}c_{y}} O_{\nu_{1}\nu_{4}\nu_{3}\nu_{2}\nu_{5}}(p_{1}, p_{4}, p_{3}, p_{2}, p_{5}) + f^{c_{1}c_{4}c_{x}} f^{c_{2}c_{x}c_{y}} f^{c_{2}c_{3}c_{y}} O_{\nu_{1}\nu_{4}\nu_{2}\nu_{3}\nu_{5}}(p_{1}, p_{4}, p_{3}, p_{2}, p_{5}) + f^{c_{1}c_{4}c_{x}} f^{c_{5}c_{x}c_{y}} f^{c_{2}c_{3}c_{y}} O_{\nu_{1}\nu_{4}\nu_{2}\nu_{3}\nu_{3}}(p_{1}, p_{5}, p_{2}, p_{3}, p_{4}) + f^{c_{1}c_{5}c_{x}} f^{c_{2}c_{x}c_{y}} f^{c_{2}c_{3}c_{y}} O_{\nu_{1}\nu_{5}\nu_{3}\nu_{2}\nu_{4}}(p_{1}, p_{5}, p_{3}, p_{2}, p_{4}) \\ + f^{c_{1}c_{5}c_{x}} f^{c_{4}c_{x}c_{y}} f^{c_{2}c_{3}c_{y}} O_{\nu_{1}\nu_{5}\nu_{4}\nu_{2}\nu_{3}}(p_{1}, p_{5}, p_{4}, p_{2}, p_{3}) + f^{c_{2}c_{3}c_{x}} f^{c_{1}c_{x}c_{y}} f^{c_{4}c_{5}c_{y}} O_{\nu_{2}\nu_{3}\nu_{1}\nu_{4}\nu_{5}}(p_{2}, p_{3}, p_{1}, p_{4}, p_{5}) \\ + f^{c_{2}c_{4}c_{x}} f^{c_{1}c_{x}c_{y}} f^{c_{3}c_{5}c_{y}} O_{\nu_{2}\nu_{4}\nu_{1}\nu_{3}\nu_{5}}(p_{2}, p_{4}, p_{1}, p_{3}, p_{5}) + f^{c_{2}c_{5}c_{x}} f^{c_{1}c_{x}c_{y}} f^{c_{2}c_{4}c_{y}} O_{\nu_{2}\nu_{3}\nu_{1}\nu_{4}\nu_{5}}(p_{2}, p_{3}, p_{1}, p_{4}, p_{5}) \\ + f^{c_{2}$$

with

$$O_{\nu_{1}\nu_{2}\nu_{3}\nu_{4}\nu_{5}}(p_{1}, p_{2}, p_{3}, p_{4}, p_{5}) = \Delta_{\nu_{3}} \left[ \Delta_{\nu_{5}} \varepsilon_{\Delta\nu_{1}\nu_{2}\nu_{4}} - \Delta_{\nu_{4}} \varepsilon_{\Delta\nu_{1}\nu_{2}\nu_{5}} \right] \sum_{m=2}^{N-1} (\Delta . p_{1} + \Delta . p_{2})^{m-2} (-\Delta . p_{4} - \Delta . p_{5})^{N-m-1} + \tilde{O}_{\nu_{1}\nu_{2}\nu_{3}\nu_{4}\nu_{5}}(p_{1}, p_{2}, p_{3}, p_{4}, p_{5}) + \tilde{O}_{\nu_{2}\nu_{1}\nu_{3}\nu_{5}\nu_{4}}(p_{2}, p_{1}, p_{3}, p_{5}, p_{4}) + \tilde{O}_{\nu_{4}\nu_{5}\nu_{3}\nu_{2}\nu_{1}}(p_{4}, p_{5}, p_{3}, p_{2}, p_{1}) + \tilde{O}_{\nu_{5}\nu_{4}\nu_{3}\nu_{2}\nu_{1}}(p_{5}, p_{4}, p_{3}, p_{2}, p_{1}) \right]$$

$$(147)$$

and

$$\dot{O}_{\nu_{1}\nu_{2}\nu_{3}\nu_{4}\nu_{5}}(p_{1}, p_{2}, p_{3}, p_{4}, p_{5}) = -\Delta_{\nu_{2}}\Delta_{\nu_{3}}\Delta_{\nu_{4}}\left[\Delta_{\nu_{5}}\varepsilon_{\Delta p_{5}p_{1}\nu_{1}} + p_{5}.\Delta\varepsilon_{\Delta\nu_{5}\nu_{1}p_{1}}\right]\sum_{m=2}^{N-3}\sum_{n=m+1}^{N-2}\sum_{o=n+1}^{N-1} \left(\Delta.p_{1}\right)^{m-2} \left(\Delta.p_{1} + \Delta.p_{2}\right)^{n-m-1} \times \left(-\Delta.p_{4} - \Delta.p_{5}\right)^{o-n-1} \left(-\Delta.p_{5}\right)^{N-o-1} + \Delta_{\nu_{3}}\Delta_{\nu_{4}}\left[\Delta_{\nu_{2}}\varepsilon_{\Delta p_{5}\nu_{1}\nu_{5}} - \Delta_{\nu_{1}}\varepsilon_{\Delta p_{5}\nu_{2}\nu_{5}}\right] \times \sum_{m=2}^{N-2}\sum_{n=m+1}^{N-1} \left(\Delta.p_{1} + \Delta.p_{2}\right)^{m-2} \left(-\Delta.p_{4} - \Delta.p_{5}\right)^{n-m-1} \left(-\Delta.p_{5}\right)^{N-n-1}.$$
(148)

All momenta are inflowing and the symbols  $f^{abc}$  denote the structure constants of  $SU(N_c)$ . For the sums in the Feynman rule it is understood that the upper summation bound is larger or equal than the lower bound.

#### References

- [1] B. Lampe and E. Reya, Phys. Rept. **332** (2000) 1–163 [arXiv:hep-ph/9810270 [hep-ph]].
- [2] S.D. Drell and T.M. Yan, Annals Phys. 66 (1971) 578–623.
- J. Blümlein and H. Böttcher, Nucl. Phys. B 841 (2010) 205–230 [arXiv:1005.3113 [hep-ph]];
   Proc. of DIS 2012, Bonn, 2012, 237–241, DESY-PROC-2012-02, [arXiv:1207.3170 [hep-ph]].
- [4] A. Piccione and G. Ridolfi, Nucl. Phys. B 513 (1998) 301–316 [arXiv:hep-ph/9707478 [hep-ph]];
   J. Blümlein and A. Tkabladze, Nucl. Phys. B 553 (1999) 427–464 [arXiv:hep-ph/9812478 [hep-ph]].
- [5] D.Y. Bardin, J. Blümlein, P. Christova and L. Kalinovskaya, Nucl. Phys. B 506 (1997) 295–328 [arXiv:hep-ph/9612435 [hep-ph]];
  A. Arbuzov, D.Y. Bardin, J. Blümlein, L. Kalinovskaya and T. Riemann, Comput. Phys. Commun. 94 (1996) 128–184 [arXiv:hep-ph/9511434 [hep-ph]];
  I. Akushevich, A. Ilyichev, N. Shumeiko, A. Soroko and A. Tolkachev, Comput. Phys. Commun. 104 (1997) 201–244 [arXiv:hep-ph/9706516 [hep-ph]];
  J. Blümlein and H. Kawamura, Nucl. Phys. B 708 (2005) 467–510 [arXiv:hep-ph/0409289 [hep-ph]].
- [6] H.D. Politzer, Phys. Rept. **14** (1974) 129–180.

- [7] A.J. Buras, Rev. Mod. Phys. 52 (1980) 199–276;
  E. Reya, Phys. Rept. 69 (1981) 195–353;
  J. Blümlein, Prog. Part. Nucl. Phys. 69 (2013) 28–84 [arXiv:1208.6087 [hep-ph]].
- [8] D. Boer, M. Diehl, R. Milner, R. Venugopalan, W. Vogelsang, D. Kaplan, H. Montgomery, S. Vigdor, A. Accardi and E. C. Aschenauer, et al. Gluons and the quark sea at high energies: Distributions, polarization, tomography, [arXiv:1108.1713 [nucl-th]].
- [9] S. Bethke et al., Workshop on Precision Measurements of α<sub>s</sub>, arXiv:1110.0016 [hep-ph];
  S. Moch et al., High precision fundamental constants at the TeV scale, arXiv:1405.4781 [hep-ph];
  S. Alekhin, J. Blümlein and S.O. Moch, Mod. Phys. Lett. A **31** (2016) no.25, 1630023.
- [10] S. Moch, J.A.M. Vermaseren and A. Vogt, Nucl. Phys. B 688 (2004) 101–134 [hepph/0403192].
- [11] J. Blümlein, P. Marquard, C. Schneider and K. Schönwald, Nucl. Phys. B 971 (2021) 115542 [arXiv:2107.06267 [hep-ph]].
- [12] D.J. Gross and F. Wilczek, Phys. Rev. D 8 (1973) 3633–3652.
- [13] H. Georgi and H.D. Politzer, Phys. Rev. D 9 (1974) 416–420.
- [14] K. Sasaki, Prog. Theor. Phys. 54 (1975) 1816–1827.
- [15] M.A. Ahmed and G.G. Ross, Phys. Lett. B 56 (1975) 385–390.
- [16] G. Altarelli and G. Parisi, Nucl. Phys. B **126** (1977) 298–318
- [17] R. Mertig and W.L. van Neerven, Z. Phys. C 70 (1996) 637–654 [hep-ph/9506451v2].
- [18] W. Vogelsang, Phys. Rev. D 54 (1996) 2023–2029 [hep-ph/9512218]; Nucl. Phys. B 475 (1996) 47–72 [hep-ph/9603366].
- [19] Y. Matiounine, J. Smith and W.L. van Neerven, Phys. Rev. D 58 (1998) 076002 [hepph/9803439].
- [20] S. Moch, J.A.M. Vermaseren and A. Vogt, Nucl. Phys. B 889 (2014) 351–400 [arXiv: 1409.5131 [hep-ph]].
- [21] A. Behring, J. Blümlein, A. De Freitas, A. Goedicke, S. Klein, A. von Manteuffel, C. Schneider and K. Schönwald, Nucl. Phys. B 948 (2019) 114753 [arXiv:1908.03779 [hep-ph]].
- [22] Y. Matiounine, J. Smith and W.L. van Neerven, Phys. Rev. D 57 (1998) 6701–6722 [arXiv: hep-ph/9801224 [hep-ph]].
- [23] J.A. Dixon and J.C. Taylor, Nucl. Phys. B 78 (1974) 552–560.
- [24] H. Kluberg-Stern and J. B. Zuber, Phys. Rev. D 12 (1975) 482–488; 3159–3180.
- [25] S. Sarkar, Nucl. Phys. B 82 (1974) 447–460.
- [26] S. Sarkar and H. Strubbe, Nucl. Phys. B **90** (1975) 45–51
- [27] S.D. Joglekar and B.W. Lee, Annals Phys. 97 (1976) 160–215.

- [28] S.D. Joglekar, Annals Phys. **108** (1977) 233–241; **109** (1977) 210–287.
- [29] R. Hamberg and W.L. van Neerven, Nucl. Phys. B **379** (1992) 143–171.
- [30] R. Hamberg, Second order gluonic contributions to physical quantities, PhD Thesis, U. Leiden, 1991.
- [31] J.C. Collins and R.J. Scalise, Phys. Rev. D 50 (1994) 4117–4136 [arXiv:hep-ph/9403231 [hep-ph]].
- [32] B.W. Harris and J. Smith, Phys. Rev. D 51 (1995) 4550–4560 [arXiv:hep-ph/9409405 [hep-ph]].
- [33] J. Blümlein, G. Falcioni and A. De Freitas, Nucl. Phys. B 910 (2016) 568–617 [arXiv: 1605.05541 [hep-ph]].
- [34] A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, Nucl. Phys. B 897 (2015) 612–644 [arXiv:1504.08217 [hep-ph]].
- [35] J. Blümlein and M. Saragnese, Phys. Lett. B 820 (2021) 136589 [arXiv:2107.01293 [hep-ph]].
- [36] S.A. Larin, Phys. Lett. B **303** (1993) 113–118 [arXiv:hep-ph/9302240 [hep-ph]].
- [37] J. Blümlein and C. Schneider, Phys. Lett. B 771 (2017) 31–36 [arXiv:1701.04614 [hep-ph]].
- [38] S. Moch, J.A.M. Vermaseren and A. Vogt, Phys. Lett. B 748 (2015) 432–438 [arXiv: 1506.04517 [hep-ph]].
- [39] F. Yndurain, The Theory of Quark and Gluon, (Springer, Berlin 2006).
- [40] I. Bierenbaum, J. Blümlein and S. Klein, Nucl. Phys. B 820 (2009) 417–482 [arXiv:0904.3563 [hep-ph]].
- [41] G. 't Hooft and M.J.G. Veltman, Nucl. Phys. B 44 (1972) 189–213;
  D.A. Akyeampong and R. Delbourgo, Nuovo Cim. A 17 (1973) 578–586; A 18 (1973) 94–104; A 19 (1974) 219–224;
  P. Breitenlohner and D. Maison, Commun. Math. Phys. 52 (1977) 39–54; 55–75.
- [42] D. Kreimer, Phys. Lett. B 237 (1990) 59-62;
  J.G. Körner, D. Kreimer and K. Schilcher, Z. Phys. C 54 (1992) 503-512;
  D. Kreimer, Dimensionale Regularisierung im Standardmodell, PhD Thesis, U. Mainz (1992).
  D. Kreimer, The Role of γ<sub>5</sub> in dimensional regularization, hep-ph/9401354;
  T L. Trueman, Z. Phys. C 69 (1996) 525-536 [arXiv:hep-ph/9504315 [hep-ph]];
  E. Kraus, Annals Phys. 262 (1998) 155-259 [hep-th/9709154];
  S. Weinzierl, Equivariant dimensional regularization, hep-ph/9903380;
  D. Stöckinger, Methoden zur Renormierung supersymmetrischer Eichtheorien in der Wess-Zumino-Eichung und deren Anwendung, PhD Thesis, TU Karlsruhe (2001);
  F. Jegerlehner, Eur. Phys. J. C 18 (2001) 673-679 [hep-th/0005255];
  H. Bélusca-Maïto, A. Ilakovac, P. Kühler, M. Mađor-Božinović and D. Stöckinger, [arXiv: 2109.11042 [hep-ph]].

- [43] D.J. Gross and F. Wilczek, Phys. Rev. Lett. **30** (1973) 1343–1346;
  H.D. Politzer, Phys. Rev. Lett. **30** (1973) 1346–1349;
  W.E. Caswell, Phys. Rev. Lett. **33** (1974) 244–246;
  D.R.T. Jones, Nucl. Phys. B **75** (1974) 531–538;
  O.V. Tarasov, A.A. Vladimirov and A.Y. Zharkov, Phys. Lett. B **93** (1980) 429–432;
  S.A. Larin and J.A.M. Vermaseren, Phys. Lett. B **303** (1993) 334–336 [hep-ph/9302208];
  T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin, Phys. Lett. B **400** (1997) 379–384 [hep-ph/9701390];
  M. Czakon, Nucl. Phys. B **710** (2005) 485–498 [hep-ph/0411261];
  P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, Phys. Rev. Lett. **118** (2017) no.8, 082002 [arXiv:1606.08659 [hep-ph]];
  F. Herzog, B. Ruijl, T. Ueda, J.A.M. Vermaseren and A. Vogt, JHEP **02** (2017) 090 [arXiv:1701.01404 [hep-ph]];
  T. Luthe, A. Maier, P. Marquard and Y. Schröder, JHEP **10** (2017) 166 [arXiv:1709.07718 [hep-ph]].
- [44] K.G. Chetyrkin, G. Falcioni, F. Herzog and J.A.M. Vermaseren, JHEP **1710** (2017) 179
   Addendum: [JHEP **1712** (2017) 006] [arXiv:1709.08541 [hep-ph]].
- [45] K.G. Chetyrkin, Nucl. Phys. B **710** (2005) 499–510 [arXiv:hep-ph/0405193 [hep-ph]].
- [46] E. Egorian and O.V. Tarasov, Teor. Mat. Fiz. 41 (1979) 26–32 [Theor. Math. Phys. 41 (1979) 863–867].
- [47] T. Luthe, A. Maier, P. Marquard and Y. Schröder, JHEP **1701** (2017) 081 [arXiv:1612.05512 [hep-ph]].
- [48] J. Blümlein, P. Marquard, C. Schneider, and K. Schönwald, DESY 21–028.
- [49] J. Blümlein and N. Kochelev, Nucl. Phys. B 498 (1997) 285–309 [arXiv:hep-ph/9612318 [hep-ph]].
- [50] P. Nogueira, J. Comput. Phys. **105** (1993) 279–289.
- [51] J.A.M. Vermaseren, New features of FORM, math-ph/0010025;
   M. Tentyukov and J.A.M. Vermaseren, Comput. Phys. Commun. 181 (2010) 1419–1427 [hep-ph/0702279].
- [52] T. van Ritbergen, A.N. Schellekens and J.A.M. Vermaseren, Int. J. Mod. Phys. A 14 (1999) 41–96 [arXiv:hep-ph/9802376 [hep-ph]].
- [53] J. Ablinger, J. Blümlein, S. Klein and C. Schneider, Nucl. Phys. Proc. Suppl. 205-206 (2010) 110–115 [arXiv:1006.4797 [math-ph]];
  J. Blümlein, A. Hasselhuhn and C. Schneider, PoS (RADCOR 2011) 032 [arXiv:1202.4303 [math-ph]];
  C. Schneider, Computer Algebra Rundbrief 53 (2013) 8–12;
  C. Schneider, J. Phys. Conf. Ser. 523 (2014) 012037 [arXiv:1310.0160 [cs.SC]].
- [54] P. Marquard and D. Seidel, *The* Crusher *algorithm*, unpublished.
- [55] J. Blümlein, P. Marquard and C. Schneider, PoS (RADCOR2019) 078 [arXiv:1912.04390 [cs.SC]].

- [56] M. Kauers, *Guessing Handbook*, JKU Linz, Technical Report RISC 09–07.
- [57] J. Blümlein, M. Kauers, S. Klein and C. Schneider, Comput. Phys. Commun. 180 (2009) 2143–2165 [arXiv:0902.4091 [hep-ph]].
- [58] Sage, http://www.sagemath.org/
- [59] M. Kauers, M. Jaroschek, and F. Johansson, in: Computer Algebra and Polynomials, Editors: J. Gutierrez, J. Schicho, Josef, M. Weimann, Eds.. Lecture Notes in Computer Science 8942 (Springer, Berlin, 2015) 105–125 [arXiv:1306.4263 [cs.SC]].
- [60] C. Schneider, Sém. Lothar. Combin. 56 (2007) 1–36 article B56b.
- [61] C. Schneider, Simplifying Multiple Sums in Difference Fields, in: Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions Texts and Monographs in Symbolic Computation eds. C. Schneider and J. Blümlein (Springer, Wien, 2013) 325–360 [arXiv:1304.4134 [cs.SC]].
- [62] J.A.M. Vermaseren, Int. J. Mod. Phys. A 14 (1999) 2037–2076 [hep-ph/9806280].
- [63] J. Blümlein and S. Kurth, Phys. Rev. D 60 (1999) 014018 [hep-ph/9810241].
- [64] E. Remiddi and J.A.M. Vermaseren, Int. J. Mod. Phys. A 15 (2000) 725–754 [hep-ph/9905237].
- [65] J. Ablinger, J. Blümlein and C. Schneider, J. Phys. Conf. Ser. 523 (2014) 012060 [arXiv:1310.5645 [math-ph]];
  J. Ablinger, PoS (LL2014) 019 [arXiv:1407.6180[cs.SC]]; A Computer Algebra Toolbox for Harmonic Sums Related to Particle Physics, Diploma Thesis, JKU Linz, 2009, arXiv:1011.1176[math-ph]; Computer Algebra Algorithms for Special Functions in Particle Physics, Ph.D. Thesis, Linz U. (2012) arXiv:1305.0687[math-ph]; PoS (LL2016) 067; Experimental Mathematics 26 (2017) [arXiv:1507.01703 [math.CO]]; PoS (RADCOR2017) 001 [arXiv:1801.01039 [cs.SC]]; arXiv:1902.11001 [math.CO]; PoS (LL2018) 063.
- [66] J. Ablinger, J. Blümlein and C. Schneider, J. Math. Phys. 52 (2011) 102301 [arXiv:1105.6063 [math-ph]]. [67]
- [67] J. Ablinger, J. Blümlein and C. Schneider, J. Math. Phys. 54 (2013) 082301 [arXiv:1302.0378 [math-ph]].
- [68] J. Ablinger, J. Blümlein, C.G. Raab and C. Schneider, J. Math. Phys. 55 (2014), 112301 [arXiv:1407.1822 [hep-th]].
- [69] J. Blümlein, Comput. Phys. Commun. 159 (2004) 19–54 [arXiv:hep-ph/0311046 [hep-ph]].
- [70] J. Blümlein, D.J. Broadhurst and J.A.M. Vermaseren, Comput. Phys. Commun. 181 (2010) 582–625 [arXiv:0907.2557 [math-ph]].
- [71] J. Blümlein, Comput. Phys. Commun. 180 (2009) 2218–2249 [arXiv:0901.3106 [hep-ph]].
- [72] J. Lagrange, Nouvelles recherches sur la nature et la propagation du son, Miscellanea Taurinensis, t. II, 1760-61; Oeuvres t. I, p. 263;
   C.F. Gauß, Theoria attractionis corporum sphaeroidicorum ellipticorum homogeneorum methodo novo tractate, Commentationes societas scientiarum Gottingensis recentiores, Vol

III, 1813, Werke Bd. V pp. 5–7;

G. Green, Essay on the Mathematical Theory of Electricity and Magnetism, Nottingham, 1828 [Green Papers, pp. 1–115];

- M. Ostrogradski, Mem. Ac. Sci. St. Peters., 6, (1831) 39–53;
- S. Laporta, Int. J. Mod. Phys. A 15 (2000) 5087–5159 [hep-ph/0102033].
- [73] M. Karr, J. ACM **28** (1981) 305–350;
  - M. Bronstein, J. Symbolic Comput. 29 (2000) no. 6 841-877;
  - C. Schneider, *Symbolic Summation in Difference Fields*, Ph.D. Thesis RISC, Johannes Kepler University, Linz technical report 01–17 (2001);
  - C. Schneider, An. Univ. Timisoara Ser. Mat.-Inform. 42 (2004) 163–179;
  - C. Schneider, J. Differ. Equations Appl. 11 (2005) 799-821;
  - C. Schneider, Appl. Algebra Engrg. Comm. Comput. 16 (2005) 1–32;
  - C. Schneider, J. Algebra Appl. 6 (2007) 415-441;
  - C. Schneider, Clay Math. Proc. 12 (2010) 285–308 [arXiv:0904.2323 [cs.SC]];
  - C. Schneider, Ann. Comb. 14 (2010) 533–552 [arXiv:0808.2596];

C. Schneider, in: Computer Algebra and Polynomials, Applications of Algebra and Number Theory, J. Gutierrez, J. Schicho, M. Weimann (ed.), Lecture Notes in Computer Science (LNCS) 8942 (2015) 157–191 [arXiv:1307.7887 [cs.SC]];

C. Schneider, J. Symb. Comput. **72** (2016) 82–127 [arXiv:1408.2776 [cs.SC]]. C. Schneider, J. Symb. Comput. **80** (2017) 616–664 [arXiv:1603.04285 [cs.SC]];

S.A. Abramov, M. Bronstein, M. Petkovšek, Carsten Schneider, J. Symb. Comput. 107 (2021) 23–66 [arXiv:2005.04944 [cs.SC]].

- [74] K.G. Chetyrkin and F.V. Tkachov, Nucl. Phys. B **192** (1981) 159–204.
- [75] F.V. Tkachov, Phys. Lett. B 100 (1981) 65–68; Theor. Math. Phys. 56 (1983) 866–870; Teor. Mat. Fiz. 56 (1983) 350–356;
  See also: T. Gehrmann, E.W.N. Glover, T. Huber, N. Ikizlerli and C. Studerus, JHEP 06 (2010) 094 [arXiv:1004.3653 [hep-ph]];
  R.N. Lee, A.V. Smirnov and V.A. Smirnov, Nucl. Phys. B Proc. Suppl. 205-206 (2010) 308–313 [arXiv:1005.0362 [hep-ph]];
  R.N. Lee and V.A. Smirnov, JHEP 02 (2011) 102 [arXiv:1010.1334 [hep-ph]].
- [76] T. Gehrmann and E. Remiddi, Comput. Phys. Commun. 144 (2002) 200–223 [arXiv:hep-ph/0111255 [hep-ph]];
  J. Vollinga and S. Weinzierl, Comput. Phys. Commun. 167 (2005) 177–194 [arXiv:hep-ph/0410259 [hep-ph]];
  J. Ablinger, J. Blümlein, M. Round and C. Schneider, Comput. Phys. Commun. 240 (2019) 189–201 [arXiv:1809.07084 [hep-ph]].
- [77] J. Ablinger, J. Blümlein, P. Marquard, N. Rana and C. Schneider, Nucl. Phys. B 939 (2019) 253–291 [arXiv:1810.12261 [hep-ph]].
- [78] J. Ablinger and J. Blümlein, in: Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions, (Springer, Wien, 2013), eds. C. Schneider and J. Blümlein, 1–32, [arXiv:1304.7071 [math-ph]].
- [79] J. Blümlein, Comput. Phys. Commun. 133 (2000) 76–104 [arXiv:hep-ph/0003100 [hep-ph]];
   J. Blümlein and S. O. Moch, Phys. Lett. B 614 (2005) 53–61 [arXiv:hep-ph/0503188 [hep-ph]];

A.V. Kotikov and V.N. Velizhanin, [arXiv:hep-ph/0501274 [hep-ph]];
J. Blümlein, Clay Math. Proc. 12 (2010) 167–188 [arXiv:0901.0837 [math-ph]].

- [80] E.B. Zijlstra and W.L. van Neerven, Nucl. Phys. B 417 (1994), 61–100 [Erratum: Nucl. Phys. B 426 (1994), 245; Erratum: Nucl. Phys. B 773 (2007), 105–106; Erratum: Nucl. Phys. B 501 (1997), 599–599].
- [81] W.L. van Neerven, Fortran-code for the Wilson coefficients of the structure functions  $g_1(x, Q^2)$  to two-loop order, unpublished.
- [82] I. Bierenbaum et al., DESY 15–004.
- [83] A. D. Watson, Z. Phys. C 12 (1982) 123–125;
   M. Glück, E. Reya and W. Vogelsang, Nucl. Phys. B 351 (1991) 579–592;
   W. Vogelsang, Z. Phys. C 50 (1991) 275–284.
- [84] J.A. Gracey, Nucl. Phys. B 480 (1996) 73–98 [arXiv:hep-ph/9609301 [hep-ph]].
- [85] J.F. Bennett and J.A. Gracey, Phys. Lett. B 432 (1998) 209–214 [arXiv:hep-ph/9803446 [hep-ph]].
- [86] J. Blümlein, Lect. Notes Phys. 546 (2000) 42–57 [arXiv:hep-ph/9909449 [hep-ph]].
- [87] J. Bartels, B.I. Ermolaev and M.G. Ryskin, Z. Phys. C 72 (1996), 627-635 [arXiv:hep-ph/9603204 [hep-ph]].
- [88] J. Blümlein and A. Vogt, Phys. Lett. B 386 (1996), 350-358 [arXiv:hep-ph/9606254 [hep-ph]].
- [89] A. Vogt, S. Moch, M. Rogal and J.A.M. Vermaseren, Nucl. Phys. B Proc. Suppl. 183 (2008) 155–161 [arXiv:0807.1238 [hep-ph]].
- [90] Y. Kiyo, J. Kodaira and H. Tochimura, Z. Phys. C 74 (1997) 631–639 [arXiv:hep-ph/9701365 [hep-ph]].
- [91] J. Bartels, B.I. Ermolaev and M.G. Ryskin, Z. Phys. C 70 (1996) 273–280 [hep-ph/9507271].
- [92] R. Kirschner and L.N. Lipatov, Nucl. Phys. B **213** (1983) 122–148.
- [93] J. Blümlein and A. Vogt, Phys. Lett. B **370** (1996) 149–155 [hep-ph/9510410].
- [94] S. Wandzura and F. Wilczek, Phys. Lett. B 72 (1977) 195–198.
- [95] J. Blümlein, V. Ravindran and W.L. van Neerven, Nucl. Phys. B 586 (2000) 349–381 [arXiv:hep-ph/0004172 [hep-ph]];
  J. Blümlein and H. Böttcher, Nucl. Phys. B 636 (2002) 225–263 [arXiv:hep-ph/0203155 [hep-ph]].
- [96] J. Ablinger, J. Blümlein, A. De Freitas, M. Saragnese, C. Schneider and K. Schönwald, Nucl. Phys. B 952 (2020) 114916 [arXiv:1911.11630 [hep-ph]];
  J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel, C. Schneider and K. Schönwald, Nucl. Phys. B 953 (2020) 114945 [arXiv:1912.02536 [hep-ph]];
  A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel, K. Schönwald and C. Schneider, Nucl. Phys. B 964 (2021) 115331 [arXiv:2101.05733 [hep-ph]];

J. Blümlein, A. De Freitas, M. Saragnese, C. Schneider and K. Schönwald, Phys. Rev. D **104** (2021) no.3, 034030 [arXiv:2105.09572 [hep-ph]];

J. Ablinger, J. Blümlein, A. De Freitas, A. Goedicke, M. Saragnese, C. Schneider and K. Schönwald, Nucl. Phys. B **955** (2020) 115059 [arXiv:2004.08916 [hep-ph]];

J. Ablinger et al., DESY 15–112; DESY 20–053;

J. Blümlein, J. Ablinger, A. Behring, A. De Freitas, A. von Manteuffel, and C. Schneider, PoS (QCDEV2017) 031 [arXiv:1711.07957 [hep-ph]].